FIPRONIX 67mg SPOT ON SOLUTION FOR SMALL DOGS
Norbrook USA
6x Carton
80 x 45 x 133 mm
Text - ≥ 7.5pt

TRIAL

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2015-2017 LIC. NO.

7018.1

000000C0?

6 APPLICATORS

Fipronix™ for DOGS

For use ONLY on Small DOGS & PUPPIES 8 weeks and older and

5 to 22 lbs

NET CONTENTS: 0.138 fl. oz. (4.02 mL)
(Six applicators containing 0.023 fl. oz. each)

6 APPLICATORS

KEEP OUT OF REACH OF CHILDREN
CAUTION See Package Insert for First Aid and Precautionary Statements

Fipronix™ for DOGS

Contains the same active ingredient as in FRONTLINE® Top Spot® for dogs

FRONTLINE® and Top Spot® are registered trademarks of Merial

For use ONLY on Small DOGS & PUPPIES 8 weeks and older and

5 to 22 lbs

DO NOT USE ON CATS
Keep cats away from treated dogs for 24 hours. If applied to or ingested by a cat, contact your veterinarian.

Long-lasting
Waterproof
Kills fleas
Kills ticks
including those that may transmit Lyme disease
Kills mosquitoes
Kills chewing lice

ACTIVE INGREDIENT:
fipronil.....9.7%
OTHER INGREDIENTS: 90.3%
TOTAL: 100.0%

Norbrook®

NET CONTENTS: 0.138 fl. oz. (4.02 mL)
(Six applicators containing 0.023 fl. oz. each)

Long-lasting
Waterproof
6 Doses

For Use Only on Small DOGS & PUPPIES 8 weeks and older and 5 to 22 lbs
READ ENTIRE LABEL AND ENCLOSED DIRECTIONS BEFORE EACH USE.

Fipronix for Dogs acts quickly, lasts up to 30 days and does not wash off treated dogs. Controls fleas, ticks, chewing lice, mosquitoes and sarcoptic mange mites.
Best if used year round. Fipronix for Dogs keeps working even after shampooing and rinsing with water, and continues to work if dogs are outdoors and exposed to the sun or rain, or even after swimming.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not allow children to apply product. TO PREVENT HARM TO YOU AND YOUR DOG, READ THE ENTIRE LABEL AND DIRECTIONS BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY. FOR EXTERNAL USE ON DOGS ONLY. DO NOT USE ON RABBITS. DO NOT USE ON CATS OR OTHER ANIMALS.

Side Effects: After application, check your dog for side effects such as redness from skin irritation, scratching or other signs of discomfort. In some cases, vomiting or diarrhea may occur. If you observe these signs in your treated dog, contact your veterinarian. If your dog has an unusual reaction to the initial application, consult a veterinarian before repeating treatment. Call 1-866-591-5777 for customer service.

DIRECTIONS FOR APPLICATION:
See enclosed package insert for directions for application.

FREQUENCY OF APPLICATION:
Monthly application kills fleas, ticks, chewing lice and mosquitoes and controls flea, tick, chewing lice and sarcoptic mange mite infestations. You can reapply Fipronix for Dogs once every 30 days.

LIMITATION OF WARRANTY AND LIABILITY: Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the Conditions, Disclaimer of Warranties and Limitations of Liability herein.

Made in Ireland

Manufactured by:
Norbrook Laboratories Limited, Newry BT35 6PU, Co. Down, Northern Ireland

EPA Reg. No. 86223-3
EPA Establishment No. 86223-IRL-001

Fipronix for Dogs is not manufactured or distributed by Merial.

Fipronix™ is a trademark of Norbrook Laboratories Limited.

DO NOT USE ON CATS
Keep cats away from treated dogs for 24 hours. If applied to or ingested by a cat, contact your veterinarian.

0 000000 000000 >

NORB401 - 80 x 45 x 133

Batch No:

TRIAL

FIPRONIX DOGS
Small Dogs Insert
148 x 210 mm
Text - ≥ 8pt

Fipronix™ for DOGS

For use ONLY on Small DOGS & PUPPIES 8 weeks and older and 5 to 22 lbs.

ACTIVE INGREDIENT:
fipronil 9.7%
OTHER INGREDIENTS: 90.3%
TOTAL: 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

Long-lasting. Waterproof.
For monthly control of fleas, ticks, chewing lice, mosquitoes and aids in the control of sarcoptic mange mites
Contains the same active ingredient as in FRONTLINE® Top Spot® for dogs
Kills fleas
Kills newly emerged adult fleas prior to egg-laying
Stops existing flea infestations and prevents establishment of new infestations
Kills adult fleas which may cause flea allergy dermatitis
Kills all life stages of ticks, including brown dog ticks (*Rhipicephalus sanguineus*), American dog ticks (*Dermacentor variabilis*), lone star ticks (*Amblyomma americanum*) and deer ticks (*Ixodes scapularis*)
Kills ticks that may transmit Lyme disease, Rocky Mountain Spotted Fever, ehrlichiosis, babesiosis and anaplasmosis
Kills mosquitoes
Kills chewing lice
Aids in control of sarcoptic mange mite infestations

FRONTLINE® and Top Spot® are registered trademarks of Merial.

FIRST AID	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
If on skin or clothing:	Take off contaminated clothing. Wash skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor immediately for treatment advice.
If swallowed:	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If in eyes:	Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
For additional information on this pesticide product (including health concerns, medical emergencies, or pesticide concerns), you may call 1-866-591-5777, twenty-four (24) hours per day, seven (7) days per week.	

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

HAZARDS TO HUMANS

CAUTION. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

HAZARDS TO DOMESTIC ANIMALS

CAUTION. FOR EXTERNAL USE ON DOGS ONLY. DO NOT USE ON CATS OR OTHER ANIMALS. Keep cats away from treated dogs for 24 hours. If applied to a cat or ingested by a cat, contact your veterinarian. Do not use on puppies under 8 weeks of age or 5 lbs. Individual sensitivities, while rare, may occur after using ANY pesticide product. Some dogs may experience temporary irritation at the site of product application. If signs persist, or become more severe within a few days of application, consult a veterinarian immediately. Certain medications can interact with pesticides. Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing dogs.

Side Effects. After application, check your dog for side effects such as redness from skin irritation, scratching or other signs of discomfort. In some cases, vomiting or diarrhea may occur. If you observe these signs in your treated dog, contact your veterinarian. If your dog has an unusual reaction to the initial application, consult a veterinarian before repeating treatment. Call 1-866-591-5777 for customer service.

DIRECTIONS FOR USE
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not allow children to apply product. TO PREVENT HARM TO YOU AND YOUR DOG, READ THE ENTIRE LABEL AND DIRECTIONS BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY. FOR EXTERNAL USE ON DOGS ONLY. DO NOT USE ON RABBITS. DO NOT USE ON CATS OR OTHER ANIMALS.

Fipronix for Dogs acts quickly, lasts up to 30 days and does not wash off treated dogs. Fipronix for Dogs controls fleas, ticks, chewing lice, mosquitoes and aids in the control of sarcoptic mange mites. For use on dogs and puppies 8 weeks of age or older (including small dogs and puppies weighing 5 to 22 lbs., medium dogs weighing 23-44 lbs., large dogs weighing 45-88 lbs., and extra-large dogs weighing 89-132 lbs.). **Use the appropriate product size based on the weight of your dog.** Best if used year round. Fipronix for Dogs keeps working even after shampooing and rinsing with water. Fipronix for Dogs continues to work if dogs are outdoors and exposed to the sun or rain, or even after swimming.

A one-spot application is convenient and easy for effective treatment of fleas, ticks, chewing lice, and mosquitoes. Fipronix for Dogs also aids in the control of sarcoptic mange mite infestations. Apply Fipronix for Dogs when your dog is dry. Fipronix for Dogs is applied at a single point on the surface of the skin above the dog's shoulder blades at the base of the neck and rapidly covers the entire dog by localizing in the hair, on the surface of the skin and in the sebaceous glands. Soon after coming into contact with a treated dog, fleas, ticks, mosquitoes and chewing lice are killed. Allow treated area to dry thoroughly after application.

TRIAL

Dogs can be bathed 24 hours after treatment. Do not reapply for 30 days.

FREQUENCY OF APPLICATION
Monthly application of Fipronix for Dogs controls flea, tick and chewing lice infestations. You can reapply Fipronix for Dogs once every 30 days.

Fleas: Fipronix for Dogs kills adult fleas for up to three months unless the risk is high for reinfestation of fleas. If your dog becomes reinfested with fleas, or if your dog is in situations where reinfestation may occur, you may apply Fipronix for Dogs every 30 days.

Ticks: Fipronix for Dogs kills ticks for at least one month. Apply Fipronix for Dogs every 30 days to control all life stages of ticks that may transmit Lyme disease, Rocky Mountain Spotted Fever, ehrlichiosis, babesiosis and anaplasmosis.

Chewing Lice: Fipronix for Dogs kills chewing lice for at least one month. Apply Fipronix for Dogs every 30 days to control chewing lice.

Mosquitoes: Apply Fipronix for Dogs every 30 days to kill mosquitoes. Mosquitoes are killed within 24 hours for up to 7 days, and within 48 hours for up to 28 days.

Sarcoptic Mange Mites: Fipronix for Dogs aids in the control of sarcoptic mange mite infestations. Apply Fipronix for Dogs every 30 days until mites are eliminated.

DIRECTIONS FOR APPLICATION
Apply Fipronix for Dogs as follows:

- **How to open:** Follow the instructions on the pouch to remove the applicator. Hold applicator upright away from face/body to allow product to drain from tip. Then cut with scissors or twist applicator to open it.
- **How to apply**: Part the dog's hair above the shoulder blades, at the base of the neck. Place the applicator tip just above the skin and squeeze. Apply entire contents of the applicator in a single spot directly onto the dog's skin. Do not apply on top of the hair coat. Avoid contact with treated area until dry.



DO NOT USE ON CATS.

DO NOT USE ON CATS. Keep cats away from treated dogs for 24 hours. If applied to or ingested by a cat, contact your veterinarian.

PHYSICAL OR CHEMICAL HAZARDS
COMBUSTIBLE. Do not use or store near heat or open flame.

000000I0?

STORAGE AND DISPOSAL
Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE
Store in orignal container in a cool dry (preferably locked) place that is inaccessible to children and animals (pets).

CONTAINER DISPOSAL
Nonrefillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP (1-800-253-2687) for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITATION OF WARRANTY AND LIABILITY
IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all which are beyond the control of the manufacturer and sellers. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, the manufacturer and sellers make no warranty, expressed or implied, of merchantability or of fitness for a particular purpose, or otherwise, concerning the use of this product other than as indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling are contrary to label instructions.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, neither the manufacturer nor the sellers shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

Made in Ireland

Manufactured by:
Norbrook Laboratories Limited, Newry, BT35 6PU,
Co. Down, Northern Ireland.

EPA Reg. No. 86223-3
EPA Establishment No. 86223-IRL-001

Fipronix for Dogs is not manufactured or distributed by Merial. Fipronix™ is a trademark of Norbrook Laboratories Limited.

Norbrook®