

APPROVED

By Lisa Adamson at 10:30 am, Nov 24, 2014

ZoGuard plus FOR DOGS

ZoGuard Plus for Dogs contains Fipronil and S-Methoprene (insect growth regulator) to effectively kill fleas and ticks (including those that may transmit Lyme disease). The active ingredients in ZoGuard Plus for Dogs feature a sustained release formula developed for fast-acting and long-lasting control of fleas, ticks, and chewing lice for up to 1 month on dogs.

DUAL ACTION! Fipronil with (S)-methoprene IGR Effectively Breaks the Flea Life Cycle



- Use ZoGuard Plus for Dogs monthly to break the flea life cycle. Kills fleas, ticks, chewing lice, flea eggs, flea larvae & prevents the development of flea pupae. Kills Ticks including those that may transmit Lyme disease. Kills brown dog ticks, American dog ticks, lone star ticks, and deer ticks for up to 1 month. Rapid elimination of chewing lice infestation. Controls mites that may cause sarcoptic mange. Fast-acting. Long-lasting. Waterproof. Can be used on breeding, pregnant and lactating dogs.

STORAGE AND DISPOSAL: Do not remove tube from the pack until ready to use. Store in a cool, dry place (below 77°F/25°C) away from children and pets. Do not refrigerate. Protect from direct sunlight. Do not use or refill this container. Empty, resealable container. Do not use or refill this container. Offer for recycling if available. If ready filled, call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

ZoGuard plus FOR DOGS ONLY For Dogs 45-88 lbs

In Case of Emergency, call: 1-844-685-9173. EPA REG. NO. 53883-351-91300. EPA EST. NO. 2517-NE-001. Made in USA.

READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON DOGS AND PUPPIES OVER 8 WEEKS OF AGE.

PRECAUTIONARY STATEMENTS. HAZARDS TO HUMANS. CAUTION. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. HAZARDS TO DOMESTIC ANIMALS. For external use on dogs only. Individual sensitivities, while rare, may occur after using any pesticide product on dogs. Certain medications can interact with pesticides. Do not treat your dog with more than one pesticide product at a time. Over dosing your dog can result in serious illness and even death. Consult a veterinarian before using on medicated, debilitated, aged, pregnant or nursing dogs. Consult a veterinarian before using on dogs with known organ dysfunction. If your dog is exhibiting signs of and/or is being treated for skin dermatitis, talk to your veterinarian before applying any topical flea and tick control product.

Table with 2 columns: Condition (If swallowed, If in eyes, If on skin) and Action (Call a poison control center or a doctor immediately for treatment advice, Flush eyes with plenty of water, Wash with plenty of soap and water).

DIRECTIONS FOR USE. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT ALLOW CHILDREN TO APPLY PRODUCT. TO PREVENT HARM TO YOU AND YOUR DOG, READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY. Do not use on cats, rabbits or animals other than dogs. Do not allow your dog to ingest this product. Do not use on puppies under 8 weeks of age or less than 45 lbs. Use entire contents of tube val on each dog. Do not use multiple tubes on one dog. Weigh your dog to be sure you are applying the right dose formulated for the weight of your dog. Separate the treated dog from all other dogs and cats for 24 hours after treatment has been applied.

How to open: Remove product tubes from the package. Separate one tube from the others. Hold the tube with notched end pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the lines. How to apply: Invert tube over dog and use open end to part dog's hair. Squeeze tube firmly to apply all of the solution to the dog's skin, as directed below. Do not apply more often than once every 4 weeks.

Only For Dogs Weighing 45 to 88 lbs. Apply one tube (2.68 mL) (0.091 fl oz) as a spot on the dog's back between the shoulder blades. Side Effects: Monitor your dog after application. Dogs may experience some temporary irritation at the site of product application such as redness, scratching or other signs of discomfort. If signs of sensitivity occur, bathe your dog with a mild soap and rinse with large amounts of water. If these or other side effects occur, consult your veterinarian or call 1-844-685-9173. If your dog has an unusual reaction to the initial application, consult a veterinarian before repeating application. The most common signs of incidental ingestion are excessive salivation and foaming at the mouth. If these symptoms occur, immediately feed your dog and continue to monitor your dog for the next 24 hours.

FREQUENCY OF APPLICATION. Use ZoGuard Plus for Dogs monthly for complete control of flea, tick, and chewing lice infestations. Studies show that ZoGuard Plus for Dogs kills adult fleas for up to three months. Apply monthly if your dog is at high risk for flea reinfestation. Apply monthly to control ticks and chewing lice. ZoGuard Plus for Dogs remains effective even after bathing, water immersion, or exposure to sunlight. Allow treated area to dry thoroughly. Do not reapply for 30 days. If you have questions or comments about this product, please write: Promika, LLC 1204 Village Market Pl #273, Morrisville, NC 27560. Distributed by: Promika, LLC 1204 Village Market Place #273 Morrisville, NC 27560.



ZoGuard plus FOR DOGS ONLY For Dogs 45-88 lbs P005922 8 59837005029



LICENSED 7020.3 PERIOD 2015-2017 LIC. NO.

3 MONTH SUPPLY ONLY For Dogs 45-88 lbs

ZoGuard plus FOR DOGS



Contains Fipronil and S-Methoprene, the active ingredients used in FRONTLINE Plus for Dogs

- KILLS FLEAS, TICKS, FLEA EGGS, FLEA LARVAE, CHEWING LICE

KEEP OUT OF REACH OF CHILDREN CAUTION. See Back Panel For Additional Precautionary Statements. NET CONTENTS: 0.273 fl oz Three 0.091 fl oz (2.68 mL) Applicators. Frontline is a registered trademark of Merial. ZoGuard Plus for Dogs is not manufactured, or distributed by Merial, seller of Frontline Plus for Dogs.

Table with 2 columns: Active Ingredients (Fipronil: 9.80%, (S)-methoprene: 8.80%) and Other Ingredients (81.40%, TOTAL: 100.0%).



DIRECTIONS FOR USE

How to Open. 1. Remove product tubes from the package. 2. Separate one tube from the others. Hold the tube with notched end pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the line.

How to Apply. 1. Invert tube over dog and use open end to part dog's hair. 2. Squeeze tube firmly to apply all of the solution to the dog's skin as a spot on the dog's back between the shoulder blades.

6 STEP CHECKLIST

- 1. Read the label completely and follow directions. 2. Weigh your dog. 3. Do NOT use dog products on cats. 4. Do NOT treat with more than one pesticide product at a time. 5. Separate animals after treatment to avoid chance of ingestion. 6. Do NOT split tubes between pets.

3 MONTH TRACKER

Table for 3-month tracker with columns for Treatment 1, 2, and 3, and rows for MM, DD, and YY.

APPROVED

By Lisa Adamson at 10:36 am, Nov 24, 2014

ZoGuard™ Plus for Dogs
For Dogs 4-22 lbs ≥8 wks
fipronil 9.8%, s-methoprene 8.8%
0.023 fl oz P005925



ZoGuard™ Plus for Dogs
For Dogs 23-44 lbs ≥8 wks
fipronil 9.8%, s-methoprene 8.8%
0.045 fl oz P005926



KEEP OUT OF REACH OF CHILDREN
CAUTION: Read directions and precautions
before using. Use scissors to open.
EPA REG. No. 53883-351-91300 P005941



ZoGuard™ Plus for Dogs
For Dogs 45-88 lbs ≥8 wks
fipronil 9.8%, s-methoprene 8.8%
0.091 fl oz P005927



ZoGuard™ Plus for Dogs
For Dogs 89-132 lbs ≥8 wks
fipronil 9.8%, s-methoprene 8.8%
0.136 fl oz P005928

