



advecta II FOR cats

For use ONLY on cats 8 weeks and Older and Weighing over 9 lbs.

4 MONTH SUPPLY

4 MONTH SUPPLY

For use ONLY on cats 8 weeks and Older and Weighing OVER 9 LBS



Effectively breaks the flea life cycle

4 MONTH SUPPLY

For use ONLY on cats 8 weeks and Older and Weighing over 9 lbs.

advecta II FOR cats

Effectively breaks the flea life cycle

advecta II FOR cats



CONTAINS THE SAME ACTIVE INGREDIENTS USED IN ADVANTAGE® II

Advecta™ is not manufactured by or distributed by Bayer. Advantage® is a registered trademark of Bayer Healthcare LLC.

For use ONLY on cats 8 weeks and Older and Weighing over 9 lbs.

advecta II FOR cats

Effectively breaks the flea life cycle

Convenient, easy-to-apply topical solution. For use ONLY on cats 8 weeks and Older and Weighing over 9 lbs.

ACTIVE INGREDIENTS:

Imidacloprid	9.10%
Pyriproxyfen	0.46%
OTHER INGREDIENTS	90.44%
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE BACK PANEL FOR PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE.

Net Contents: 0.108 fl. oz (3.2ml), Contains 4 tubes, each 0.027 fl.oz (0.8ml)

STATE OF HAWAII
Department of Agriculture

ACCEPTED
LICENSE NO. **7072.6**



8 18145 01522 9

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies or consumer questions call 1-855-844-4375.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment or advice.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling. Keep out of reach of children. Do not contaminate feed or food.

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not apply to cats or kittens under 8 weeks of age or weighing less than 9 lbs. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing cats. Individual sensitivities, while rare, may occur after using ANY pesticide product for cats. If signs persist, or become more severe consult a veterinarian immediately. If your cat is on medication, consult your veterinarian before using this or any other product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not allow children to apply product.

READ ENTIRE LABEL BEFORE EACH USE

RESTRICTIONS

USE ONLY ON CATS OR KITTENS 8 WEEKS AND OLDER WEIGHING OVER 9 lbs.
DO NOT USE ON OTHER ANIMALS
DO NOT EXCEED LABELED DOSAGE AMOUNT FOR CATS
DO NOT APPLY MORE THAN ONE (1) TUBE PER TREATMENT ON ANY CAT

EPA Reg. No. 2596-181-91384
EPA Est. No. 2596-OH-1

RESTRICTIONS (Cont.)

DO NOT HAVE CONTACT OR ALLOW CHILDREN TO HAVE CONTACT WITH TREATED AREA UNTIL COMPLETELY DRY.

SIDE EFFECTS (CATS): Monitor your cat after application. Side effects, although very rare, may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as hypersalivation, vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call, 1-855-844-4375.

HOW TO APPLY

1. Remove one applicator tube from the package.
2. Hold applicator tube in an upright position **facing away from you and your pet's face and eyes.** Pull cap off tube.
3. Turn the cap around and place other end of cap back on tube.
4. Twist cap to break seal, then remove cap from tube.
5. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. **Do not get this product in your pet's eyes, or allow your pet to ingest this product.** The product is bitter tasting and salivation may occur for a short time if the pet licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat to lick the product. Do not allow the product to run off.
6. Discard empty tube as described in the Storage and Disposal section.
7. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. On cats, do not retreat more often than once every seven (7) days. After flea control is attained, return to a monthly treatment schedule.


STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry place inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Non-refillable container. If empty: Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-888-846-4231 for disposal instructions. Never place unused product down any indoor or outdoor drain.

For more information call our experts at 1-888-846-4231

Manufactured for: CAP IM Supply, Inc.
303 Perimeter Center North, Suite 300
Atlanta, GA 30346
Made in the USA



113053

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies or consumer questions call 1-855-844-4375.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment or advice.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling. Keep out of reach of children. Do not contaminate feed or food.

HAZARDS TO DOMESTIC ANIMALS

For external use only. Do not apply to cats or kittens under 8 weeks of age or weighing less than **9 lbs.** As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing cats. Individual sensitivities, while rare, may occur after using ANY pesticide product for cats. If signs persist, or become more severe consult a veterinarian immediately. If your cat is on medication, consult your veterinarian before using this or any other product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not allow children to apply product.

READ ENTIRE LABEL BEFORE EACH USE

RESTRICTIONS

USE ONLY ON CATS OR KITTENS 8 WEEKS AND OLDER WEIGHING OVER 9 lbs.

DO NOT USE ON OTHER ANIMALS

DO NOT EXCEED LABELED DOSAGE AMOUNT FOR CATS

DO NOT APPLY MORE THAN ONE (1) TUBE PER TREATMENT ON ANY CAT

RESTRICTIONS (Cont.)

DO NOT HAVE CONTACT OR ALLOW CHILDREN TO HAVE CONTACT WITH TREATED AREA UNTIL COMPLETELY DRY.

SIDE EFFECTS (CATS): Monitor your cat after application. Side effects, although very rare, may include signs of skin irritation such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as hypersalivation, vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call, 1-855-844-4375.

HOW TO APPLY

1. Remove one applicator tube from the package.
2. Hold applicator tube in an upright position **facing away from you and your pet's face and eyes.** Pull cap off tube.
3. Turn the cap around and place other end of cap back on tube.
4. Twist cap to break seal, then remove cap from tube.
5. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. *Do not get this product in your pet's eyes, or allow your pet to ingest this product.* The product is bitter tasting and salivation may occur for a short time if the pet licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat to lick the product. Do not allow the product to run off.
6. Discard empty tube as described in the Storage and Disposal section.
7. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four (4) weeks. On cats, do not retreat more often than once every seven (7) days. After flea control is attained, return to a monthly treatment schedule.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry place inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Non-refillable container. **If empty:** Do not reuse or refill this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-888-846-4231 for disposal instructions. Never place unused product down any indoor or outdoor drain.

For more information call our experts at 1-888-846-4231

Manufactured for: CAP IM Supply, Inc.
303 Perimeter Center North, Suite 300
Atlanta, GA 30346
Made in the USA

EPA Reg. No. 2596-181-91384
EPA Est. No. 2596-OH-1

EPA Reg. No. 2596-181-91384
EPA Est. No. 2596-OH-1

**READ ENTIRE LABEL
BEFORE USE**

CAUTION

**KEEP OUT OF REACH
OF CHILDREN**

9.10% IMIDACLOPRID
0.46% PYRIPROXYFEN

advecta[™] II
for cats

ONLY FOR USE ON CATS AND
KITTENS 8 WEEKS OR
OLDER WEIGHING
9 lbs. and Over-0.027 fl. oz. (0.8 ml)

RM113041

advecta^{II}

FOR

cats

REMINDER STICKERS

The first time you treat your cat, place the first month sticker on your calendar. Apply Advecta^{II} for Cats monthly, every 30 days. Use these calendar stickers to remind you when your cat is due for the next application.



APPLY FIRST MONTH

advecta^{II}
FOR **cats**



APPLY SECOND MONTH

advecta^{II}
FOR **cats**



APPLY THIRD MONTH

advecta^{II}
FOR **cats**



APPLY FOURTH MONTH

advecta^{II}
FOR **cats**
REPURCHASE

For more information call our experts at 1-888-846-4231

Manufactured for: CAP IM Supply, Inc.
303 Perimeter Center North, Suite 300 Atlanta, GA 30346

Made in the USA