

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2014-2016** LIC. NO.

8012.25

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2011-2013** LIC. NO.

Parastar™ Plus contains fipronil and cyphenothrin to effectively start killing ticks and fleas in as little as 1 hour. Kills ticks, including those

PARASTAR
(fipronil/cyphenothrin) plus

For dogs & puppies 12 weeks or older and up to **22** lbs.

NOVARTIS
ANIMAL HEALTH



For dogs & puppies 12 weeks or older and up to **22** lbs.

NET CONTENTS: (3) 0.023 fl oz (0.67 mL) applicators

PARASTARTM

(fipronil/cyphenothrin) plus



Fast-acting. Long-lasting. Waterproof.

Kills Ticks, Fleas & Chewing Lice

See Back for full Directions for Use, First Aid and Precautionary Statements.

KEEP OUT OF REACH OF CHILDREN
WARNING

ACTIVE INGREDIENTS:
Fipronil.....9.8%
Cyphenothrin.....5.2%
OTHER INGREDIENTS:.....85.0%
TOTAL:.....100.0%



Starts killing ticks in as little as 1 hour

NOVARTIS
ANIMAL HEALTH



© 2011 Novartis Animal Health US, Inc.
Parastar is a trademark of Novartis AG.

NAH/PARA-PLUS/0.67/UDC/2
Product #609087



LOT NO:

NOVARTIS
ANIMAL HEALTH

PARASTARTM
(fipronil/cyphenothrin) plus

For dogs & puppies 12 weeks or older and up to **22** lbs.

PARASTAR[®] *plus*

(fipronil/cyphenothrin)

Parastar[™] Plus contains fipronil and cyphenothrin to effectively start killing ticks and fleas in as little as 1 hour. Kills ticks, including those that may transmit Lyme disease, Rocky Mountain spotted fever, ehrlichiosis, babesiosis, anaplasmosis and hepatozoonosis.



Parastar[™] Plus

- Kills Deer Ticks
- Kills Brown Dog Ticks
- Kills American Dog Ticks
- Kills Lone Star Ticks
- Kills fleas
- Kills chewing lice
- Controls mites that may cause Sarcosis Mange



READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON DOGS AND PUPPIES 12 WEEKS OF AGE OR OLDER.

For Technical Information, call 1-800-637-0281
Made in USA

Manufactured for
Novartis Animal Health US, Inc.
3200 Northline Avenue, Suite 300
Greensboro, NC 27408

EPA Registration No. 70585-13
EPA Est. No. 2517-TN-1
P003275

How to open:

1. Remove product tubes from the package.
2. Separate one tube from the others.
Holding the tube with notched end pointing up and away from the face and body, cut off the narrow end at the notches with scissors.

How to apply:

1. Invert tube over dog and use open end to part dog's hair.
2. Squeeze tube firmly to apply all of the solution to the dog's skin as a spot to the dog's back between the shoulder blades.



PARASTAR[®] *plus*

(fipronil/cyphenothrin)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE EACH USE.

Do not allow children to apply product. TO PREVENT HARM TO YOU AND YOUR DOG, READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY. DO NOT USE ON RABBITS. DO NOT USE ON OTHER ANIMALS. DO NOT USE ON CATS. May be toxic and potentially fatal if applied to or ingested by cats. FOR EXTERNAL USE ON DOGS ONLY. Do not use on puppies under 12 weeks of age. Weigh your dog to be sure you are applying the right dose formulated for the weight of your pet. Do not treat your pet with more than one pesticide product at a time. Over dosing your animal can result in serious illness and even death. **Best if used year round.** Certain medications can interact with pesticides. Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant, or nursing dogs. Consult a veterinarian before using on dogs with known organ dysfunction. DO NOT USE ON CATS or animals other than dogs. Separate the treated dog from all other dogs and cats for 24 hours after application. Cats and dogs that actively groom or engage in close physical contact with treated dog may be at risk for serious harmful effects. The most common signs of ingestion are excessive salivation and foaming at the mouth. If these symptoms occur immediately feed your dog and continue to monitor your dog for the next 24 hours. If symptoms persist, contact a veterinarian. Cats exhibiting these same symptoms from ingestion should be taken to a veterinarian immediately. If your dog is exhibiting signs of and/or is being treated for skin dermatitis, talk to your vet before applying any topical flea and tick control product.

Sensitivities may occur after using ANY pesticide product on dogs. Dogs may experience some temporary irritation at the site of product application. If signs of sensitivity occur bathe your dog with a mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

How to open: Remove product tubes from the package. Separate one tube from the others. Holding the tube with notched end pointing up and away from the face and body, cut off the narrow end at the notches with scissors.

How to apply: While holding the dog with one hand, use the other hand to apply the solution. Invert the tube and use open end to part dog's hair. Apply contents of one tube of Parastar[™] Plus as a spot to the dog's back between the shoulder blades. Hold the dog for a few seconds to give the solution time to be absorbed into the dog's coat. When finished, wrap the tube in paper and put into trash. Only one applicator per treatment is needed. Repeat applications may be made if necessary, but do not apply more than once every 4 weeks.

FREQUENCY OF APPLICATION

Use Parastar[™] Plus monthly for complete control of flea, tick and chewing lice infestations. Studies show that Parastar[™] Plus kills fleas for up to 30 days. If your pet is at high risk for flea reinfestation, or in a highly infested environment, apply monthly. Apply monthly to control ticks and chewing lice. Parastar[™] Plus remains effective, even after bathing, water immersion, or exposure to sunlight. Allow treated area to dry thoroughly. Do not reapply for 30 days.

For Dogs Weighing up to 22 lbs.

Apply one tube 0.023 fl. oz. (0.67 mL) as a spot on the dog's back between the shoulder blades.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Do not remove tube from the pack until ready to use. Store in a cool (below 77°F) dry place inaccessible to children and pets. Do not refrigerate. Protect from direct sunlight.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container.

If empty: Place in the trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

FIRST AID

- | | |
|-------------------------------|---|
| If in eyes | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15–20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for further treatment advice. |
| If on skin or clothing | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15–20 minutes.• Call a poison control center or doctor for treatment advice. |
| If swallowed | <ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person. |

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For more information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-800-637-0281.

NOTE TO PHYSICIAN OR VETERINARIAN

Treat patient symptomatically.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS. WARNING. Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Harmful if swallowed. Harmful if absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

HAZARDS TO DOMESTIC ANIMALS. WARNING. For external use only. Do not use on puppies under 12 weeks of age. Individual sensitivities, while rare, may occur after using any pesticide product. Dogs may experience some temporary irritation at the site of product application. If signs persist, or become more severe within a few days of application, consult a veterinarian immediately. If your dog has an unusual reaction to the initial application, consult a veterinarian before repeating treatment. Certain medications can interact with pesticides. Consult a veterinarian before using on medicated, debilitated or aged dogs.

WARRANTY

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of NOVARTIS ANIMAL HEALTH US, Inc. (NAHUS) or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold NAHUS and Seller harmless for any claims relating to such factors.

NAHUS warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or NAHUS, and Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NAHUS MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with State law, in no event shall NAHUS or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF NAHUS AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF NAHUS OR SELLER, THE REPLACEMENT OF THE PRODUCT.



NAH/PARA-PLUS/067/UDC/2
Product #609087

© 2011 Novartis Animal Health US, Inc.
Parastar is a trademark of Novartis AG.




4.6346



0.9167

4.6346



0.9167

PARASTAR 
(fipronil/cyphenothrin) **plus**



CALENDAR
STICKERS



CALENDAR
STICKERS



© 2011 Novartis Animal Health US, Inc.
Parastar is a trademark of Novartis AG.
NAH/PARA-PLUS/SC/1
P003312