

Artwork shown at approximately 110% of actual size

Kills Pandemic 2009 H1N1 influenza A virus.



- Bactericidal
- Fungicidal
- Germicidal
- Pseudomonacidal
- Staphylocidal
- Streptocidal
- Tuberculocidal
- Virucidal*

ELIMINATES ODORS

Multi-Purpose Disinfectant

Fresh Scent

ACTIVE INGREDIENTS:

| | |
|---|-----------|
| Octyl decyl dimethyl ammonium chloride..... | 0.1890% |
| Dioctyl dimethyl ammonium chloride..... | 0.0945% |
| Didecyl dimethyl ammonium chloride..... | 0.0945% |
| Alkyl (50% C14, 40% C12, 10% C16) dimethyl benzyl ammonium chlorides..... | 0.2520% |
| Ethanol..... | 65.0000% |
| INERT INGREDIENTS..... | 34.3700% |
| TOTAL..... | 100.0000% |

KEEP OUT OF REACH OF CHILDREN

WARNING: See back panel for additional precautionary statements.



NET WT. 19 OZ. (1.18 LB) 538g

Federal Regulations Prohibit CFC Propellants in Aerosols



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2014-2016** LIC. NO.

8013.20



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2011-2013** LIC. NO.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2008-2010** LIC. NO.

Artwork shown at approximately 110% of actual size

- This product meets AOAC Germicidal Spray Product Test efficacy standards for hospital disinfectants.
- Kills and prevents the growth of mold.
- Deodorizes by killing the germs that cause odors.
- Does not contain bleach.
- Kills Pandemic 2009 H1N1 influenza A virus.

Use on hard, nonporous surfaces in:

Restrooms • Hotels • Motels • Offices • Military Installations • Schools • Day Care Centers • Nurseries • Dorms • Shelters • Laboratories • Health Clubs • School Buses • Ambulances • Bowling Alleys • Play Areas • Convenience Stores • Locker Room Facilities • Storage Areas • Kennels

For use on:

Garbage Cans • Waste Baskets • Diaper Pails • Diaper Changing Tables • Toilet Seats • Faucets • Doorknobs • Telephones • Showers • Plastic Shower Curtains • Counter Tops • Desks • Metal Work Benches • Handles

Use on non-critical surfaces in:

Hospitals • Patient Rooms • Nursing Homes • Medical Clinics • Veterinary Offices.

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS & DOMESTIC ANIMALS


WARNING: Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear protective eye wear (safety glasses). Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. If skin contact with product occurs, wash thoroughly with soap and water, especially prior to food handling and preparation.

FIRST AID: IF IN EYES: Hold eyelids open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment. Questions? Call 1-800-797-7225.

PHYSICAL HAZARDS: Flammable: Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130 Fahrenheit may cause bursting.

STORAGE/DISPOSAL: Pesticide Storage and Disposal: Do not contaminate water, food, or feed by storage and disposal. Store at temperatures below 130 Fahrenheit. **Container Disposal:** Do not puncture or incinerate. Do not reuse empty container. Recycle empty container or discard in trash.

- This can is made of an average of 25% recycled steel (10% post-consumer) • Recyclable • Contains no phosphorus • Avoid use on polished wood, painted surfaces or acrylic plastics.



Questions? Comments?
Call toll-free
1-888-797-7225

Mfd. for Clorox Professional
Products Company, Oakland,
CA 94612
© 2000

The Clorox Company
Made in USA or Argentina
EPA Reg. No. 67619-3
EPA Est. No. 11525-IL-1
5813-ARG-001

See code on bottom of can.
Patent Pending

Disinfects against the following bacteria, viruses* and mold:

- Campylobacter jejuni
- Corynebacterium diphtheriae
- Enterobacter aerogenes
- Enterococcus faecalis (Vancomycin resistant)
- Escherichia coli O157:H7
- Klebsiella pneumoniae
- Listeria monocytogenes
- Mycobacterium bovis (Tuberculosis)
- Mycobacterium smegmatis
- Proteus mirabilis
- Proteus vulgaris
- Pseudomonas aeruginosa
- Pseudomonas cepacia (Burkholderia cepacia)
- Pseudomonas putida
- Salmonella choleraesuis
- Salmonella choleraesuis paratyphi B1 (schottmuelleri)
- Salmonella choleraesuis serotype enteritidis
- Serratia marcescens
- Shigella dysenteriae
- Staphylococcus aureus
- Staphylococcus aureus (Methicillin & Gentamicin resistant)
- Streptococcus pyogenes
- *Adenovirus type 2
- *Cytomegalovirus
- *Echovirus
- *Hepatitis A virus
- *Herpes simplex virus type 1
- *Herpes simplex virus 2
- *Human Immunodeficiency Virus Type 1 (HIV-1).
- *Influenza A2 virus (Hong Kong) (Flu virus)
- *Influenza virus type B
- *Polio virus
- *Respiratory syncytial virus (leading cause of lower respiratory infection in children)
- *Rhinovirus (cold virus)
- *Rhinovirus 39
- *Rotavirus (leading cause of infectious diarrhea in children)
- *Vaccinia virus
- Alternaria alternata
- Candida albicans
- Cladosporium herbarum
- Trichophyton mentagrophytes (Athlete's foot fungus)

Sanitizes in 30 seconds against: Klebsiella pneumoniae, Staphylococcus aureus.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For use on non-food contact surfaces only. A potable water rinse is required for surfaces which may be in direct contact with food. **This product must not result in the direct or indirect contamination of food products.**

SPECIFIC INSTRUCTIONS FOR *HIV-1: This product kills *HIV-1 on precleaned surfaces/objects previously soiled with blood/body fluids in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (*HIV-1) (associated with AIDS).

Special Instructions for Using this Product to Clean and Decontaminate Against *HIV-1 on Surfaces/Objects Soiled with Blood/Body Fluids.

Personal Protection: When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks, and eye coverings.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and other objects before applying this product.

Contact Time: Spray 6 to 10 inches from precleaned surface for 3-4 seconds until thoroughly wet. Surface must remain wet for 10 minutes before wiping or air drying.

Disposal of Infectious Materials: Use disposable latex gloves, gowns, masks, and eye coverings. Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

To Disinfect: Spray 6 to 10 inches from precleaned surface for 3-4 seconds until thoroughly wet. Surface must remain wet for 10 minutes before wiping.

To Sanitize Non-food Contact Surfaces: Spray 6-10 inches from precleaned surface for 3-4 seconds or until thoroughly wet. Surface must remain wet for 30 seconds before air drying.

To Control and Prevent the Growth of Mold and Mildew: Spray precleaned surface until thoroughly wet. Surface must remain wet for 10 minutes before wiping or air drying. Respray product as necessary for ongoing control.

To Deodorize: Spray on precleaned surfaces as needed.

R0526-1.1