

5 1/4

18 1/16

29 3/32

3 7/8

FIRST AID	
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

HOTLINE NUMBER
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For more information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-800-637-0281.

NOTE TO PHYSICIAN OR VETERINARIAN
Treat patient symptomatically.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS. CAUTION. Causes moderate eye irritation. Do not get in eyes or on clothing. Harmful if swallowed. Harmful if absorbed through skin. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.
HAZARDS TO DOMESTIC ANIMALS. For external use only. Individual sensitivities, while rare, may occur after using any pesticide product. Certain medications can interact with pesticides. Do not treat your dog with more than one topically applied pesticide product at a time. Overdosing your dog can result in serious illness and even death. Consult a veterinarian before using on medicated, pregnant, nursing, debilitated or aged dogs.

WARRANTY
The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of NOVARTIS ANIMAL HEALTH US, Inc. (NAHUS) or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold NAHUS and Seller harmless for any claims relating to such factors. NAHUS warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or NAHUS, and Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NAHUS MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE. To the extent consistent with State law, in no event shall NAHUS or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF NAHUS AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF NAHUS OR SELLER, THE REPLACEMENT OF THE PRODUCT.



© 2013 Novartis Animal Health US, Inc.
Parastar is a registered trademark of Novartis AG. NAH/PARA-PLUS/4.02/UDC/3
Product #609090

CAUTION
KEEP OUT OF REACH OF CHILDREN
Starts Killing Fleas and Ticks in Just 5 Minutes
Fast-acting. Long-lasting. Waterproof.
(fipronil/cyphenothrin)
PARASTAR plus for dogs
Apply only to puppies over 12 weeks old and over 89 lbs.
NET CONTENTS: (3) 0.136 fl oz (4.02 mL) applicators
EASY-TO-USE 3 APPLICATIONS
NOVARTIS ANIMAL HEALTH

PARASTAR plus for dogs
(fipronil/cyphenothrin)
Apply only to puppies over 12 weeks old and over 89 lbs.

PARASTAR plus for dogs
(fipronil/cyphenothrin)
Parastar Plus for Dogs

- Kills fleas
- Kills deer ticks
- Kills brown dog ticks
- Kills American dog ticks
- Kills lone star ticks
- Kills chewing lice
- Controls mites that may cause sarcoptic mange

How to Open:
1. Remove product tubes from the package.
2. Separate one tube from the others. Holding the tube with notched end pointing up and away from the face and body, cut off the narrow end at the notches with scissors.

How to Apply:
1. Invert tube over dog and use open end to part dog's hair.
2. Squeeze tube firmly to apply all of the solution to the dog's skin as a spot to the dog's back between the shoulder blades.

Side Effects
Monitor your dog after application. Dogs may experience some temporary irritation at the site of product application, such as redness, scratching or other signs of discomfort. If signs of sensitivity occur, bathe your dog with a mild soap and rinse with large amounts of water. If these or other side effects occur, consult your veterinarian or Novartis Animal Health at 1-800-637-0281. If your dog has an unusual reaction to the initial application, consult a veterinarian before repeating treatment. The most common signs of ingestion are excessive salivation and foaming at the mouth. If these symptoms occur, immediately feed your dog and continue to monitor your dog for the next 24 hours.

DO NOT USE ON CATS—MAY BE FATAL
Keep cats away from treated dogs for 24 hours. If applied to a cat or ingested by a cat, contact your veterinarian immediately.

READ ENTIRE LABEL BEFORE EACH USE. USE ONLY ON DOGS AND PUPPIES 12 WEEKS OF AGE.

For Technical Information, call 1-800-637-0281
Made in USA
Manufactured for Novartis Animal Health US, Inc., 3200 Northline Avenue, Suite 300 Greensboro, NC 27408
EPA Registration No. 70585-13
EPA Est. No. 2517-NE-02
P003278-4 RARC

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE EACH USE.
Do not allow children to apply product. TO PREVENT HARM TO YOU AND YOUR DOG, READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY. DO NOT USE ON RABBITS. DO NOT USE ON OTHER ANIMALS. DO NOT USE ON CATS—MAY BE FATAL. Keep cats away from treated dogs for 24 hours. If applied to a cat or ingested by a cat, contact your veterinarian immediately. FOR EXTERNAL USE ON DOGS ONLY. Do not allow your dog to ingest this product. Do not apply to puppies less than 12 weeks of age. Do not apply to dogs or puppies weighing less than 4 lbs. For use only on dogs weighing 89-132 lbs. Weigh your dog to be sure you are applying the right dose formulated for the weight of your dog. Do not double-dose dogs weighing greater than 132 lbs.
How to open: Remove product tubes from the package. Separate one tube from the others. Holding the tube with notched end pointing up and away from the face and body, cut off the narrow end at the notches with scissors.
How to apply: While holding the dog with one hand, use the other hand to apply the solution. Invert the tube and use open end to part dog's hair. Apply contents of one tube of Parastar Plus for Dogs as a spot to the dog's back between the shoulder blades. Hold the dog for a few seconds to give the solution time to be absorbed into the dog's coat. When finished, wrap the tube in paper and put into trash. Only one applicator per treatment is needed. Repeat applications may be made if necessary, but do not apply more than once every 4 weeks.

FREQUENCY OF APPLICATION
Use Parastar Plus for Dogs monthly for complete control of flea, tick and chewing lice infestations. Studies show that Parastar Plus for Dogs kills adult fleas for up to three months. If your dog is at high risk of flea reinfestation, or in a highly infested environment, apply monthly. Apply monthly to control ticks and chewing lice. Parastar Plus for Dogs remains effective, even after bathing, water immersion, or exposure to sunlight. Allow treated area to dry thoroughly. Do not reapply for 30 days.
For use only on dogs weighing 89-132 lbs.
Apply one tube 0.136 fl. oz. (4.02 mL) as a spot on the dog's back between the shoulder blades.

STORAGE AND DISPOSAL
Do not contaminate water, food, or feed by storage and disposal.
PESTICIDE STORAGE: Do not remove tube from the pack until ready to use. Store in a cool (below 77 F) dry place inaccessible to children and pets. Do not refrigerate. Protect from direct sunlight.
CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. If empty: Place in the trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.



8013.39

GSG GRAPHIC SYSTEMS GROUP
33 EAST 17TH STREET, NEW YORK, NY 10003
TEL: 646-230-1900 FAX: 212-228-9898
File Name: P003278-4 RARC, ParastarPlus, Ors, 89-132lbs, r13a1 IWO: 400053 (prv)288284

GSG Approvals
Proofreader Traffic QC

NOVARTIS
PRODUCT DESCRIPTION
Parastar Plus 89-132 lbs. Carton
ITEM CODE NAH/PARA PLUS/4.02/UDC/3 VERSION NO. 13
REVISION DATE 1/29/2014 REVERSED BY GL
MECHANICAL DUE DUE TO VENDOR
UPC NO. 716688090904 MAG. 81% HT. .63"
DIELINE NO.
DESIGN MANAGER

Printing Inks
PMS BLACK 5 C PMS 325 C PMS 2945 C
CMYK PROCESS

Printing Info
Printer: XX
Printing Method: xxxx
Substrate: Substrate Here

Printer Notes:

6 3/4