

LICENSED

PERIOD 2014-2016 LIC. NO.

8013.45

Front panel shown at approximately 155% of actual size

LICENSED

PERIOD 2011-2013 LIC. NO.



Contains
the power of
Clorox® Bleach
Foaming Action



LICENSED

PERIOD 2008-2010 LIC. NO.

Germicidal Spray Atomizador Germicida

ACTIVE INGREDIENT:
Sodium Hypochlorite 2.40%
OTHER INGREDIENTS: 97.60%
TOTAL: 100.00%
(Available Chlorine 2.28%)

KEEP OUT OF REACH OF CHILDREN
WARNING: EYE IRRITANT. USE
ONLY IN WELL-VENTILATED AREAS.
SEE BACK LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

1 QT (32 FL OZ) 946 mL

INGREDIENTE ACTIVO:
Hipoclorito de Sodio 2,40%
OTROS INGREDIENTES: 97,60%
TOTAL: 100,00%
(Cloro Disponible 2,28%)

MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS
ADVERTENCIA: IRRITANTE DE OJOS. UTILICE EL
PRODUCTO SOLAMENTE EN ÁREAS BIEN VENTILADAS.
VEA LA ETIQUETA POSTERIOR PARA DECLARACIONES PREVENTIVAS ADICIONALES.
Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

1 CUARTO DE GALÓN (32 FL OZ) 946 mL

R0642-8

Artcode 120364.021 (tc)

Back panel shown at approximately 155% of actual size

Clorox Commercial Solutions®
Clorox® Germicidal Spray
Atomizador Germicida

PEEL/DESPEGUE

- Ready-to-Use Hospital-Grade Disinfectant

- Reduces exposure to Methicillin resistant *Staphylococcus aureus* (MRSA) from treated surfaces

- Meets OSHA Bloodborne Pathogen Standards

PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS.

WARNING: Causes substantial, but temporary, eye injury. Do not get in eyes or on clothing. Avoid contact with skin. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove and wash contaminated clothing before reuse. Vapors may irritate. Use only in well-ventilated areas. Avoid prolonged breathing of vapors. **Not recommended for use by persons with heart conditions or chronic respiratory problems such as asthma, emphysema or obstructive lung disease.** **FIRST AID: IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control center or doctor for treatment advice. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. **IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glassful of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. **PHYSICAL AND CHEMICAL HAZARDS:** This product contains bleach. Do not use or mix this product with other chemicals such as products containing ammonia, toilet bowl cleaners, rust removers, vinegar or acid. To do so will release hazardous gases.


STORAGE AND DISPOSAL: Store this product in a cool dry area to avoid deterioration. Refill only with this product. Please recycle empty container or discard in trash.
CONTAINS NO PHOSPHORUS.

QUESTIONS OR COMMENTS? Call 1-888-797-7225. For MSDS information please go to www.cloroxprofessional.com.

**Mfd. for Clorox Professional Products Company,
1221 Broadway, Oakland, CA 94612.**

© 2007 Made in the U.S.A. Commercial Solutions is a registered trademark of Clorox Professional Products Company.

EPA Reg. No. 67619-13. EPA Est. No. 5813-CA-3, GA-1
(actual EPA Est. No. in code above or below).



Artcode 121569.004 (tc)

Hinged panel shown at approximately 155% of actual size

- Shelf-Stable Formula
- Eliminates Daily Mixing and Measuring
- Foam Trigger Helps to Minimize Potential Overspray
- One-Step Cleaner/Disinfectant

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DISINFECTION: To clean, disinfect and deodorize hard, nonporous surfaces: Set trigger to SPRAY or FOAM. Spray 6–8 inches from surface until surface is thoroughly wet. Let stand for contact time listed on label. Wipe excess. Allow to air dry. Rinse with potable water for food-contact surfaces. For all others, no rinsing is required. Gross filth should be removed prior to disinfecting. If streaking is observed, wipe with a clean, damp cloth or paper towel.

Specific Instructions for Use Against HIV-1: This product kills HIV-1 on precleaned environmental surfaces/objects previously soiled with blood/body fluids in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus (HIV-1) (associated with AIDS).

Special instructions for using this product to clean and decontaminate against HIV-1 on surfaces/objects soiled with blood/body fluids: Personal Protection: When handling items soiled with blood or body fluids, use disposable impervious gloves, gowns, masks and eye coverings. **Cleaning Procedure:** Blood and other body fluids must be thoroughly cleaned from surfaces and other objects before applying this product. **Contact Time:** Allow surfaces to remain wet for 30 seconds. Wipe excess. Allow to air dry. For all other organisms, see directions for contact times. **Disposal of Infectious Materials:** Use disposable impervious gloves, gowns, masks and eye coverings. Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that **(1)** is introduced directly into the human body, either into or in contact with the bloodstream, or normally sterile areas of the body, or **(2)** contacts intact mucous membranes, but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

HINGED AREA, NO PRINT ALLOWED

R0642-8

Artcode 121569.004 (tc)

Hinged panel shown at approximately 155% of actual size

HINGED AREA, NO PRINT ALLOWED

ORGANISMS:

Mycobacterium:

- Mycobacterium bovis BCG (Tuberculosis) OT 105401 **

Bacteria:

- Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA) (NARSA NRS123)(Genotype USA400)*
- Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA) (NARSA NRS384)(Genotype USA300)*
- †Enterococcus faecalis, vancomycin resistant (VRE) ATCC 51575*
- ESBL (Extended Spectrum Beta Lactamase) producing Escherichia coli ATCC BAA-196*
- Escherichia coli O157:H7 ATCC 35150*
- †Escherichia coli (E. coli) ATCC 11229*
- Klebsiella pneumoniae ATCC 4352**
- †Legionella pneumophila ATCC 33153*
- Listeria monocytogenes ATCC 19117**
- †Pseudomonas aeruginosa ATCC 15442*
- †Salmonella enterica ATCC 10708*
- †Shigella dysenteriae ATCC 11835*

- †Staphylococcus aureus ATCC 6538*
- †Staphylococcus aureus, methicillin resistant (MRSA) ATCC 33592*
- †Streptococcus pyogenes ATCC 12344*

Viruses Non-Enveloped:

- ‡Adenovirus type 2 ATCC VR-846 (Strain Adenoid 6) (common cause of respiratory infections)**
- ‡Human Hepatitis A (Strain HM-175)**
- ‡Poliovirus type 1 ATCC VR-1000 (Strain Brunhilde)**
- ‡Rhinovirus type 37 ATCC VR-1147 (Strain 151-1) (causative agent of the common cold)**
- ‡Rotavirus (Strain WA) (causative agent of viral diarrhea)**

Mold, Mildew, Fungi:

- Alternaria alternata ATCC 6663***
- Aspergillus niger ATCC 16404**
- Cladosporium herbarum ATCC 28987***
- Mucor bainieri ATCC 42642***
- Penicillium chrysogenum ATCC 10106***
- Trichophyton mentagrophytes ATCC 26323**

* 30 second contact time

** 1 minute contact time

*** 5 minute contact time

USE SITES:

Medical:

- Ambulances
- Anesthesia Rooms
- Central Service Areas
- Critical Care Units
- Dialysis Clinics
- Emergency Rooms
- Examination Rooms
- Health Care Settings
- Home Health Care
- Hospitals
- Intensive Care Units
- Laboratories
- Medical Facilities
- Neonatal Nurseries
- Nurses' Stations
- Nursing Homes
- Orthopedics
- Outpatient Clinics
- Patient Rooms
- Physical Therapy Rooms
- Radiology Rooms
- Recovery Rooms
- Respiratory Therapy Rooms
- Surgery Rooms
- Dental:
- Dentist's Offices
- Veterinary:
- Animal Housing Facilities
- Animal Life Science Laboratories

- Kennels
- Pet Shops
- Small Animal Facilities
- Veterinary Clinics/Hospitals

Miscellaneous:

- Blood Banks
- Correctional Facilities
- Day Care Centers
- Funeral Homes
- Health Club Facilities
- Military Installations
- Recreational Centers
- Restrooms
- Schools

TO DISINFECT HARD, NONPOROUS SURFACES: FOLLOW DIRECTIONS FOR USE.

| PATHOGEN | CONTACT TIME |
|---------------------|--------------|
| TB | 1 minute |
| Bacteria | 30 seconds† |
| ‡Viruses | 1 minute |
| Parvoviruses | 5 minutes |
| Mold, Mildew, Fungi | 5 minutes |

R0642-8

Artcode 121569.004 (tc)