

Front panel shown at approximately 120% of actual size



LICENSED

PERIOD **2014-2016** LIC. NO.

8013.60



LICENSED

PERIOD **2011-2013** LIC. NO.



Commercial SOLUTIONS®



Kills *C. difficile* spores

Kills *Clostridium difficile* spores

Kills *C. diff* spores



Germicidal Wipes, Toallitas Germicidas,

ACTIVE INGREDIENT:
Sodium Hypochlorite 0.65%
OTHER INGREDIENTS: 99.35%
TOTAL: 100.00%
(Yields 0.62% available chlorine)

KEEP OUT OF REACH OF CHILDREN
CAUTION: NOT TO BE USED AS
BABY WIPES. KEEP CONTAINER
TIGHTLY SEALED WHEN NOT IN USE.
READ COMPLETE DIRECTIONS AND PRECAUTIONS ON REAR PANELS.

NET CONTENTS: 150 — 6.75" x 8"
TOWELS; 1.78 LBS (28.5 OZ) 807 g

- **Kills 99.99% *bacteria, viruses*[†] & *fungi* in 1 MINUTE on hard, nonporous surfaces**
- **Kills *Mycobacterium bovis* - BCG, (TB) in 2 MINUTES**
- **Kills *Clostridium difficile* spores in 5 MINUTES***

SI USTED NO ENTIENDE LA ETIQUETA, BUSQUE A ALGUIEN PARA QUE SE LA EXPLIQUE A USTED EN DETALLE.

R0866-1

Front panel shown at approximately 95% of actual size

- Contains 6500 ppm Sodium Hypochlorite
- Cleans, Disinfects & Deodorizes in One Step

- Pre-Moistened Towels with 1:10 Dilution of Bleach & Detergent
- Kills Pandemic 2009 H1N1 Influenza A Virus

- Kills *Clostridium difficile* Spores in 5 Minutes on Precleaned, Hard, Nonporous Surfaces

NI-15268

Premoistened disposable towels for use in healthcare and other areas where control of cross contamination is required. Meets OSHA Bloodborne Pathogen Standards. **Surfaces:** May be used on hard, nonporous surfaces such as stainless steel, glazed ceramic tile, hard plastics, nonporous vinyl, painted surfaces, glazed porcelain, sealed fiberglass surfaces, Plexiglas, laminated plastic countertops and glass.

USE SITES: Hospital and Healthcare: clinics, dental offices, emergency rooms (ER), hospitals, hospital central service areas, intensive care units (ICU), laboratory, nursing homes, physical therapy, physician offices, radiology, recovery rooms, and surgery. **Critical Care Areas:** ER, ICU, neonatal intensive care units (NICU), pediatric intensive care units (PICU) and surgery.

APPLICATIONS: Hard, nonporous surfaces such as autoclaves, bed railings, blood glucose monitors, cabinets, carts, chairs, counters, exam tables, gurneys, infant incubators and care cribs, IV poles, phlebotomy trays, plastic mattress covers, PVC tubing, stretchers, tables, ambulance equipment, diagnostic equipment, dialysis machines, mammography equipment, patient monitoring equipment, respiratory equipment, ultrasound transducers and probes, and wheelchairs. **Nursery or Daycare Centers:** changing tables, cribs, trash cans, toys and other hard, nonporous surfaces listed on this label.

Efficacy tests have demonstrated that this product is an effective bactericide, virucide, tuberculocide and fungicide in the presence of organic soil (5% blood serum).

Kills bacteria, viruses* & fungi in 1 MINUTE:

- | | | |
|--------------------------------------|---------------------------------|---------------------------------------|
| • <i>Acinetobacter baumannii</i> | • *Hepatitis C Virus (HCV) | • *Rhinovirus |
| • *Avian Influenza A Virus | • *Herpes Simplex Virus (HSV-2) | • *Rotavirus |
| • *Canine Parvovirus | • *HIV-1 (AIDS Virus) | • <i>Salmonella enterica</i> |
| • <i>Enterobacter aerogenes</i> | • *Influenza A H1N1 | • <i>Staphylococcus aureus</i> |
| • <i>Enterococcus faecalis</i> (VRE) | • *Influenza A Virus | • <i>Staphylococcus aureus</i> (MRSA) |
| • <i>Escherichia coli</i> ESBL | • <i>Klebsiella pneumoniae</i> | • <i>Streptococcus pyogenes</i> |
| • *Feline Panleukopenia Virus | • *Norovirus | • <i>Trichophyton mentagrophytes</i> |
| • *Hepatitis A Virus (HAV) | • *Poliovirus Type 1 (Mahoney) | (Athlete's Foot Fungus) |
| • *Hepatitis B Virus (HBV) | • <i>Pseudomonas aeruginosa</i> | |

Kills *Mycobacterium bovis* - BCG, (TB) in 2 MINUTES. Kills *Clostridium difficile* spores in 5 MINUTES*

PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. **FIRST AID:** If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. If present, remove contact lenses after the first 5 minutes of rinsing, then continue rinsing eye. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **PHYSICAL OR CHEMICAL HAZARDS:** Do not use this product with ammonia or acids such as vinegar, rust removers or toilet bowl cleaners. This product not intended for use on fabrics.

STORAGE AND DISPOSAL: Store this product in a cool, dry area away from direct sunlight and heat. When not in use, keep center cap of lid closed to prevent moisture loss. Nonrefillable container. Do not reuse or refill this container. Discard used towels and container in trash or take container to recycling center. Do not flush towels down toilet.

R0866-1

DIRECTIONS FOR USE It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Dispenser Directions: To Open: Remove lid. Discard inner seal from canister. **To Thread:** Find center of wipe roll, twist corner of first wipe into a point and thread through the hole in the canister lid. Pull through about one inch. Replace lid. Dispense remaining wipes as necessary by pulling out and tearing off. When not in use, keep center cap of lid closed to prevent moisture loss. Gloves should be worn.

To Clean, Disinfect & Deodorize in One Step: Wipe surface with towel until completely wet. Let stand for one minute. To kill TB, let stand for two minutes at room temperature (20–25°C). Wipe dry or allow to air dry. Gross soil must be removed prior to disinfecting. Gloves should be worn.

To Clean: Follow directions above, but no standing time is required. If streaking is observed, wipe with a clean, damp cloth or paper towel.

***Special Instructions for Cleaning Prior to Disinfection Against *Clostridium difficile* Spores. Personal Protection:** Wear appropriate barrier protection such as gloves, gowns, masks or eye covering. **Cleaning Procedure:** Fecal matter/waste must be thoroughly cleaned from surfaces/objects before disinfection by application with this product. Cleaning is to include vigorous wiping and/or scrubbing, until all visible soil is removed. Special attention is needed for high-touch surfaces. Surfaces in patient rooms are to be cleaned in an appropriate manner, such as from right to left or left to right on horizontal surfaces, and top to bottom on vertical surfaces, to minimize spreading of the spores. Restrooms are to be cleaned last. Do not reuse soiled cloths. **Infectious Materials Disposal:** Materials used in the cleaning process that may contain feces/waste are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

Disinfection Procedure: Wipe surface with towel until completely wet. Let stand for five minutes. Wipe dry or allow to air dry. Gross soil must be removed prior to disinfecting. Gloves should be worn.

†Kills HIV, HBV and HCV on precleaned environmental surfaces/objects previously soiled with blood/body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV).

Special Instructions for Cleaning and Decontamination Against HIV-1 (AIDS Virus), HBV and HCV on Surfaces/Objects Soiled with Blood/Body Fluids. Personal Protection: When handling items soiled with blood/body fluids, personal protective equipment such as disposable gloves, gowns, masks and eye coverings should be used. **Cleaning Procedure:** Blood/body fluids must be thoroughly cleaned from surfaces and objects before disinfecting. **Contact Time:** Let stand for one minute. **Disposal of Infectious Materials:** Blood, body fluids and cleaning materials must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.

Ultrasound Transducers and Probes:** For use on external surfaces of ultrasound transducers and probes only.

To Disinfect: Preclean gel from external surface of ultrasound transducer or probe to be disinfected with a paper towel or clean cloth. Then thoroughly wet the probe with a Clorox Commercial Solutions Clorox® Germicidal Wipes, towel. Let stand for one minute. To kill TB, let stand for two minutes at room temperature.

**This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semicritical medical devices prior to sterilization or high-level disinfection.

EPA Reg. No. 56392-8-67619

EPA Est. Nos. 41925-WI-3 63650-WI-1 50757-WI-1 73027-WI-1 8251-OH-1 5813-GA-2 (Actual EPA Est. No. near batch code.)

QUESTIONS OR COMMENTS? Call 1-800-234-7700. For MSDS information, please go to: www.cloroxprofessional.com.

Mfd. for Clorox Professional Products Company, 1221 Broadway, Oakland, CA 94612.

© 2011 Made in the U.S.A. Commercial Solutions is a registered trademark of Clorox Professional Products Company. Clorox is a registered trademark of The Clorox Company.

