

Artwork shown at approximately 110%. PDF prints at approximately 100% from acrobat reader or when "Fit to print" settings are selected.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2014-2016** LIC. NO.

8013.64



- Bactericidal, Virucidal,* Tuberculocidal
- Compatible with Equipment Surfaces
- Formulated with Patented Hydrogen Peroxide Technology
- One-Step Disinfectant Cleaner
- No Harsh Chemical Fumes
- Non-Flammable


NON-BLEACH

ACTIVE INGREDIENT:
 Hydrogen Peroxide.....1.4%
 OTHER INGREDIENTS.....98.6%
 TOTAL.....100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION: SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

1 QT (32 FL OZ) 946 mL

RC162010



Artwork shown at approximately 110%. PDF prints at approximately 100% from acrobat reader or when "Fit to printable area" settings are selected.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

To Refill: Remove: Remove cap from empty bottle. **Pour:** Unscrew cap on refill and pour contents directly into empty bottle. Use a funnel to minimize product waste. **Use:** Replace cap and use as you normally would.

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream, or normally sterile areas of the body, or (2) contacts intact mucous membranes, but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

SPECIAL INSTRUCTIONS FOR USE AGAINST HIV-1, HBV, AND HCV: This product kills HIV-1, HBV and HCV on precleaned hard, nonporous surfaces/objects previously soiled with blood/body fluids in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus (HIV-1) (associated with AIDS), Human Hepatitis B Virus (HBV) and Human Hepatitis C Virus (HCV). **SPECIAL INSTRUCTIONS FOR USING THIS PRODUCT TO CLEAN AND DECONTAMINATE AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS: Personal Protection:** When handling items soiled with blood or body fluids, use disposable impervious gloves, gowns, masks and eye coverings. **Cleaning Procedure:** Blood and other body fluids must be thoroughly cleaned from surfaces and other objects before applying this product. **Contact Time:** Allow surface to remain wet for HBV, HCV and HIV-1 for 30 seconds. The contact times for other bacteria, viruses and fungi may differ. See product label for contact times.

DISPOSAL OF INFECTIOUS MATERIALS: Use disposable impervious gloves, gowns, masks and eye coverings. Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal. **TO PRECLEAN INSTRUMENTS PRIOR TO TERMINAL STERILIZATION/HIGH-LEVEL DISINFECTION: As a precleaning spray –** Place instruments into a suitable container. Spray Clorox Healthcare® Hydrogen Peroxide Cleaner Disinfectant onto instruments to thoroughly wet all surfaces. Let stand for up to 10 minutes. Rinse instruments. **A manual instrument cleaner –** Thoroughly pre-rinse dirty instruments under running water to remove visible, gross debris. Immerse pre-rinsed instruments into an appropriate size container filled with Clorox Healthcare® Hydrogen Peroxide Cleaner Disinfectant. Scrub instruments using a stiff bristle brush until visibly clean. Submerge instruments while scrubbing. Rinse instruments thoroughly. Change solution daily. Follow with an appropriate disinfection process. Cleaning of critical and semi-critical devices must be followed by an appropriate terminal sterilization/high-level disinfection process. **TO DISINFECT NON-CRITICAL, PRECLEANED INSTRUMENTS:** Instruments must be thoroughly precleaned to remove excess organic debris, rinsed and rough dried. (Clean and rinse lumens of hollow instruments before filling with Clorox Healthcare® Hydrogen Peroxide Cleaner Disinfectant.) Spray all surfaces of instruments with Clorox Healthcare® Hydrogen Peroxide Cleaner Disinfectant until thoroughly wet. Let stand for one minute. To kill TB, let stand for 4 minutes at room temperature. To kill mold and fungi, let stand for 3 minutes. Wipe with a clean, damp cloth or paper towel and allow to air dry.

DISINFECTION: This product cleans, disinfects and deodorizes hard, nonporous medical surfaces in one step with no rinsing required. **TO CLEAN, DISINFECT AND DEODORIZE HARD, NONPOROUS SURFACES:** Spray 6–8 inches from surface until surface is thoroughly wet. Allow this product to remain wet for contact time listed on label. Then wipe. For heavily soiled areas, a precleaning is required. Rinse with potable water for food-contact surfaces. For all others, no rinsing is required.

PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS. **CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. **FIRST AID: If in eyes:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call the poison control center at (800) 222-1222 or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

STORAGE AND DISPOSAL: Store this product in a cool, dry area, away from direct sunlight and heat to avoid deterioration. Do not reuse or refill this container except as described in the directions for use. Refill only with this product. Offer for recycling or place in trash. **QUESTIONS OR COMMENTS?** Call (800) 234-7700. For MSDS information go to www.cloroxhealthcare.com.

based on aseptix technology

Efficacy tests have demonstrated that this product is an effective bactericide, fungicide, and virucide* in the presence of organic soil (5% blood serum).

ORGANISMS:

BACTERIA: 30 second contact time

- *Acinetobacter baumannii*
- *Campylobacter jejuni*
- Carbapenem Resistant *Klebsiella pneumoniae*
- *Enterobacter aerogenes*
- *Escherichia coli (E. coli)*
- *Escherichia coli* New Delhi Metallo-Beta Lactamase-1 (NDM-1)
- *Escherichia coli* O157:H7
- Extended Spectrum Beta Lactamase producing *Escherichia coli* (ESBL producing *E. coli*)
- Extended Spectrum Beta Lactamase producing *Klebsiella pneumoniae* (ESBL producing *Klebsiella pneumoniae*)
- *Klebsiella oxytoca*
- *Klebsiella pneumoniae* New Delhi Metallo-Beta Lactamase-1 (NDM-1)
- Multidrug Resistant *Enterococcus faecium*
- Multidrug Resistant *Klebsiella pneumoniae*
- Extended Spectrum Beta Lactamase producing *Klebsiella pneumoniae* (ESBL producing *Klebsiella pneumoniae*)
- *Pseudomonas aeruginosa*
- *Salmonella enterica*
- *Stenotrophomonas maltophilia*
- Penicillin-Resistant *Streptococcus pneumoniae*
- *Streptococcus pyogenes*
- Vancomycin Resistant *Enterococcus faecalis* (VRE)

BACTERIA: 1 minute contact time

- *Staphylococcus epidermidis* (Coagulase-negative staphylococci) (CoNS)
- Community Acquired Methicillin Resistant *Staphylococcus aureus* (CA-MRSA)
- Linezolid Resistant *Staphylococcus aureus*
- Methicillin Resistant *Staphylococcus aureus* (MRSA) (ATCC 33591)
- Methicillin Resistant *Staphylococcus aureus* (MRSA) (ATCC 33592)
- *Staphylococcus aureus*

BACTERIA: 4 minute contact time

- *Mycobacterium bovis* (TB)

MOLD, FUNGI: 3 minute contact time

- *Candida albicans*
- *Trichophyton mentagrophytes*

***VIRUSES: 30 second contact time**

- *Avian Influenza A virus (H5N1)
- *Cytomegalovirus
- *Hepatitis B virus
- *Hepatitis C virus
- *Herpes simplex virus type 1
- *Herpes simplex virus type 2
- *Human Coronavirus
- *Human Immunodeficiency Virus type 1 (HIV-1)
- *Influenza A virus (H3N2)
- *Influenza A virus (2009 H1N1)
- *Influenza B virus
- *Respiratory syncytial virus (RSV)

***VIRUSES: 1 minute contact time**

- *Norovirus
- *Rhinovirus (type 37)
- *Rotavirus

Avoid Contact With:

- Aluminum
- Copper
- Galvanized steel
- Silver


Mfd. for Clorox Professional Products Company, 1221 Broadway, Oakland, CA 94612. Clorox Healthcare is a trademark of The Clorox Company.

© 2015 Made in the U.S.A.

EPA Reg. No. 67619-24. EPA Est. No. 5813-GA-1 (A4), GA-2 (VG); 8251-OH-1 (O6), WI-3 (KV), WI-5 (UQ). Beginning of batch code indicates Est. No. Visit us at www.cloroxhealthcare.com.

U.S. Patent No. 7,658,953.

RC162010 NI-28784 0



Artwork shown at approximately 110%. PDF prints at approximately 100% from acrobat reader or when "Fit to printable area" settings are selected.

Now kills microorganisms on curtains[‡] and fabric[‡]



Hydrogen Peroxide

Cleaner Disinfectant

Limpiador Desinfectante
con Peróxido de Hidrógeno

- Bactericidal, Virucidal,[‡] Tuberculocidal
- Compatible with Equipment Surfaces
- Formulated with Patented Hydrogen Peroxide Technology

- One-Step Disinfectant Cleaner
- Soft Surface[‡] Sanitizer
- No Harsh Chemical Fumes
- Non-Flammable

NON-BLEACH

ACTIVE INGREDIENT:
Hydrogen Peroxide.....1.4%
OTHER INGREDIENTS.....98.6%
TOTAL.....100.0%

INGREDIENTE ACTIVO:
Peróxido de Hidrógeno.....1.4%
OTROS INGREDIENTES.....98.6%
TOTAL.....100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION: SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.
MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS
PRECAUCIÓN: VEA EL PANEL POSTERIOR PARA DECLARACIONES PREVENTIVAS ADICIONALES.
SI USTED NO ENTIENDE LA ETIQUETA, BUSQUE A ALGUIEN PARA QUE SE LA EXPLIQUE A USTED EN DETALLE.

1 QT (32 FL OZ) 946 mL

NI-211930

PEEL/DESPEGUE

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TO OPERATE: To open, turn nozzle. Spray 6–8 inches from surface. Wipe with cloth. Do not breathe spray or mist.

TO REFILL: **Remove:** Remove trigger sprayer from empty bottle. **Pour:** Unscrew cap on refill and pour contents directly into empty bottle. Use a funnel to minimize product waste. **Use:** Replace trigger sprayer and use as you normally would.

PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS.


CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. **FIRST AID: If in eyes:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call the poison control center at (800) 222-1222 or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

DECLARACIONES PREVENTIVAS: RIESGOS PARA LOS SERES HUMANOS Y ANIMALES DOMÉSTICOS.

PRECAUCIÓN: Causa irritación moderada a los ojos. Evite el contacto con los ojos o la ropa. Lávese bien con agua y jabón después de usar y antes de comer, beber, mascar goma, usar tabaco o usar el inodoro. **PRIMEROS AUXILIOS: Contacto con los ojos:** Mantenga los ojos abiertos y enjuague con agua lentamente, con cuidado, durante 15–20 minutos. Si hay lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando. Llame al centro de control de envenenamiento al (800) 222-1222 ó al médico para obtener asesoría sobre el tratamiento. Cuando llame a un centro de control de envenenamiento o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto.

STORAGE AND DISPOSAL: Store this product in a cool, dry area, away from direct sunlight and heat to avoid deterioration. Do not reuse or refill this container except as described in the directions for use. Refill only with this product. Offer for recycling or place in trash.

QUESTIONS OR COMMENTS? Call (800) 234-7700. For MSDS information, please go to www.cloroxhealthcare.com.
Mfd. for & © 2013 Clorox Professional Products Company, 1221 Broadway, Oakland, CA 94612.
Clorox Healthcare is a registered trademark of The Clorox Company.
Made in the USA.
EPA Reg. No. 67619-24. EPA Est. No. 5813-GA-1 (A4), GA-2 (VG); 8251-OH-1 (Q6), WI-3 (KV), WI-5 (26).
Beginning of batch code indicates Est. No.
Visit us at www.cloroxhealthcare.com.
U.S. Patent No. 7,658,953.

R0903050
NI-211930



LICENSED

PERIOD **2014-2016** LIC. NO.

8013.64

Artwork shown at approximately 110%. PDF prints at approximately 100% from acrobat reader or when "Fit to printable area" settings are selected.

HINGED AREA, NO PRINT ALLOWED

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream, or normally sterile areas of the body, or (2) contacts intact mucous membranes, but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

SPECIAL INSTRUCTIONS FOR USE AGAINST HIV-1, HBV, AND HCV: This product kills HIV-1, HBV and HCV on precleaned hard, nonporous surfaces/objects previously soiled with blood/body fluids in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus (HIV-1) (associated with AIDS), Human Hepatitis B Virus (HBV) and Human Hepatitis C Virus (HCV). **SPECIAL INSTRUCTIONS FOR USING THIS PRODUCT TO CLEAN AND DECONTAMINATE AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:** **Personal Protection:** When handling items soiled with blood or body fluids, use disposable impervious gloves, gowns, masks and eye coverings. **Cleaning Procedure:** Blood and other body fluids must be thoroughly cleaned from surfaces and other objects before applying this product. **Contact Time:** Allow surface to remain wet for HBV, HCV and HIV-1 for 30 seconds. The contact times for other bacteria, viruses and fungi may differ. See product label for contact times. **DISPOSAL OF INFECTIOUS MATERIALS:** Use disposable impervious gloves, gowns, masks and eye coverings. Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal. **TO PRECLEAN INSTRUMENTS PRIOR TO TERMINAL STERILIZATION/HIGH-LEVEL DISINFECTION:** As a precleaning spray – Place instruments into a suitable container. Spray Clorox Healthcare® Hydrogen Peroxide Cleaner Disinfectant onto instruments to thoroughly wet all surfaces. Let stand for up to 10 minutes. Rinse instruments. **A manual instrument cleaner** – Thoroughly pre-rinse dirty instruments under running water to remove visible, gross debris. Immerse pre-rinsed instruments into an appropriate size container filled with Clorox Healthcare® Hydrogen Peroxide Cleaner Disinfectant. Scrub instruments using a stiff bristle brush until visibly clean. Submerge instruments while scrubbing. Rinse instruments thoroughly. Change solution daily. Follow with an appropriate disinfection process. Cleaning of critical and

HINGED AREA, NO PRINT ALLOWED

semi-critical devices must be followed by an appropriate terminal sterilization/high-level disinfection process. **TO DISINFECT NON-CRITICAL, PRECLEANED INSTRUMENTS:** Instruments must be thoroughly precleaned to remove excess organic debris, rinsed and rough dried. (Clean and rinse lumens of hollow instruments before filling with Clorox Healthcare® Hydrogen Peroxide Cleaner Disinfectant.) Spray all surfaces of instruments with Clorox Healthcare® Hydrogen Peroxide Cleaner Disinfectant until thoroughly wet. Let stand for one minute. To kill TB, let stand for 4 minutes at room temperature. To kill mold and fungi, let stand for 3 minutes. Wipe with a clean, damp cloth or paper towel and allow to air dry.

DISINFECTION / PARA DESINFECTAR

This product cleans, disinfects and deodorizes hard, nonporous medical surfaces in one step with no rinsing required.

TO CLEAN, DISINFECT AND DEODORIZE HARD, NONPOROUS SURFACES: Spray 6–8 inches from surface until surface is thoroughly wet. Allow this product to remain wet for contact time listed on label. Then wipe. For heavily soiled areas, a precleaning is required. Rinse with potable water for food-contact surfaces. For all others, no rinsing is required.

PARA LIMPIAR, DESINFECTAR Y DESODORIZAR SUPERFICIES DURAS Y SIN POROS: Rocíe a una distancia de 6–8 pulgadas de la superficie hasta que esté bien mojada. Deje que permanezca mojada con este producto por el tiempo de contacto que se lista en la etiqueta. Luego limpie. Para áreas muy sucias, se requiere una limpieza previa. Para superficies que entren en contacto con los alimentos, enjuague con agua potable. Para todas las demás, no se requiere enjuague.

SOFT SURFACE SANITIZATION/PARA SANITIZAR SUPERFICIES BLANDAS

TO SANITIZE SOFT SURFACES: Hold bottle upright 6–8 inches from surface. Spray until fabric is wet. DO NOT SATURATE. Let stand for 30 seconds. Allow to air dry. Heavy soil must be removed prior to sanitizing. For first time use on soft surfaces, spot test in a small, inconspicuous area.

PARA SANITIZAR SUPERFICIES BLANDAS: Mantenga la botella en posición vertical a una distancia de 6–8 pulgadas de la superficie. Rocíe hasta que la tela esté mojada. NO LA SATURE. Deje reposar 30 segundos. Deje secar al aire. La mugre difícil se debe quitar antes de sanitizar. Para el primer uso en superficies blandas, haga una prueba en un área pequeña y desapercibida.

HINGED AREA, NO PRINT ALLOWED

R0903050

PREL/DESPEQUE

Artwork shown at approximately 110%. PDF prints at approximately 100% from acrobat reader or when "Fit to printable area" settings are selected.

Efficacy tests have demonstrated that this product is an effective bactericide, fungicide, and virucide* in the presence of organic soil (5% blood serum).

ORGANISMS		
BACTERIA: 30 second contact time		
<ul style="list-style-type: none"> • <i>Acinetobacter baumannii</i> • <i>Campylobacter jejuni</i> • Carbapenem Resistant <i>Klebsiella pneumoniae</i> • <i>Enterobacter aerogenes</i> • <i>Escherichia coli</i> (<i>E. coli</i>) • <i>Escherichia coli</i> New Delhi Metallo-Beta Lactamase-1 (NDM-1) • <i>Escherichia coli</i> O157:H7 • Extended Spectrum Beta Lactamase producing <i>Escherichia coli</i> (ESBL producing <i>E. coli</i>) 	<ul style="list-style-type: none"> • Extended Spectrum Beta Lactamase producing <i>Klebsiella pneumoniae</i> (ESBL producing <i>Klebsiella pneumoniae</i>) • <i>Klebsiella oxytoca</i> • <i>Klebsiella pneumoniae</i> New Delhi Metallo-Beta Lactamase-1 (NDM-1) • Multidrug Resistant <i>Enterococcus faecium</i> • Multidrug Resistant <i>Klebsiella pneumoniae</i> 	<ul style="list-style-type: none"> • <i>Pseudomonas aeruginosa</i> • <i>Salmonella enterica</i> • <i>Stenotrophomonas maltophilia</i> • Penicillin-Resistant <i>Streptococcus pneumoniae</i> • <i>Streptococcus pyogenes</i> • Vancomycin Resistant <i>Enterococcus faecalis</i> (VRE)
BACTERIA: 1 minute contact time		
<ul style="list-style-type: none"> • <i>Staphylococcus epidermidis</i> (Coagulase-negative staphylococci) (CoNS) • Community Acquired Methicillin Resistant <i>Staphylococcus aureus</i> (CA-MRSA) 	<ul style="list-style-type: none"> • Linezolid Resistant <i>Staphylococcus aureus</i> • Methicillin Resistant <i>Staphylococcus aureus</i> (MRSA) (ATCC 33591) 	<ul style="list-style-type: none"> • Methicillin Resistant <i>Staphylococcus aureus</i> (MRSA) (ATCC 33592) • <i>Staphylococcus aureus</i>
BACTERIA: 4 minute contact time	MOLD, FUNGI: 3 minute contact time	
<ul style="list-style-type: none"> • <i>Mycobacterium bovis</i> (TB) 	<ul style="list-style-type: none"> • <i>Candida albicans</i> • <i>Trichophyton mentagrophytes</i> 	
*VIRUSES: 30 second contact time		
<ul style="list-style-type: none"> • *Avian Influenza A virus (H5N1) • *Cytomegalovirus • *Hepatitis B virus • *Hepatitis C virus • *Herpes simplex virus type 1 	<ul style="list-style-type: none"> • *Herpes simplex virus type 2 • *Human Coronavirus • *Human Immunodeficiency Virus type 1 (HIV-1) • *Influenza A virus (H3N2) 	<ul style="list-style-type: none"> • *Influenza A virus (2009 H1N1) • *Influenza B virus • *Respiratory syncytial virus (RSV)
*VIRUSES: 1 minute contact time		
<ul style="list-style-type: none"> • *Norovirus 	<ul style="list-style-type: none"> • *Rhinovirus (type 37) 	<ul style="list-style-type: none"> • *Rotavirus

Soft Surface Sanitization: 30 second contact time		
<ul style="list-style-type: none"> • <i>Enterobacter aerogenes</i> • <i>Staphylococcus aureus</i> 		
Use Sites:		
<ul style="list-style-type: none"> • Ambulances • Ambulatory Surgical Centers (ASC) • CCUs • Dialysis clinics • Emergency rooms • Garbage cans • Hospitals • ICUs 	<ul style="list-style-type: none"> • Laboratories • Long term care facilities • Clinics • Newborn Intensive Care Units (NICU) • Nursing homes • Nurses' stations • Operating rooms • Patient restrooms 	<ul style="list-style-type: none"> • Patient rooms • Pediatric examination areas • Pediatric Intensive Care Units (PICU) • Physical therapy areas • Physicians' offices • Public areas • Radiology rooms
Medical Surfaces:		
<ul style="list-style-type: none"> • Bedrails • Bedside tables • Carts • Counters 	<ul style="list-style-type: none"> • High touch surfaces[†] • IV Stands, pumps, poles • Large surfaces • Shower fixtures 	<ul style="list-style-type: none"> • Stretchers • Wheelchairs
Surface Materials:		
<ul style="list-style-type: none"> • Baked enamel • Chrome • Glass 	<ul style="list-style-type: none"> • Glazed ceramic • Glazed porcelain • Laminated surfaces 	<ul style="list-style-type: none"> • Plastic • Stainless steel • Vinyl
Soft Surface:		
<ul style="list-style-type: none"> • Chairs[‡] • Couches[‡] 	<ul style="list-style-type: none"> • Curtains[‡] • Fabrics[‡] 	<ul style="list-style-type: none"> • Upholstered furniture[‡]
Avoid Contact With:		
<ul style="list-style-type: none"> • Aluminum • Copper 	<ul style="list-style-type: none"> • Galvanized steel • Silver 	<p>[‡]composed of 100% cotton or 100% polyester</p> <p>[‡]compuestas 100% de algodón ó 100% de poliéster</p>
[†] hard, nonporous surfaces of		

HINGED AREA, NO PRINT ALLOWED

HINGED AREA, NO PRINT ALLOWED

R0903050

Artwork shown at approximately 110%. PDF prints at approximately 100% from acrobat reader or when "Fit to printable area" settings are selected.

Now kills microorganisms on curtains[‡] and fabric[‡]



Hydrogen Peroxide Cleaner Disinfectant

- Bactericidal, Virucidal*, Tuberculocidal
- Compatible with Equipment Surfaces
- Formulated with Patented Hydrogen Peroxide Technology
- One-Step Disinfectant Cleaner
- Soft Surface[‡] Sanitizer
- No Harsh Chemical Fumes
- Non-Flammable

NON-BLEACH

ACTIVE INGREDIENT:
 Hydrogen Peroxide.....1.4%
 OTHER INGREDIENTS:.....98.6%
 TOTAL:.....100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION:
 SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

1 GALLON (128 FL OZ) 3.78 L

R090-4040
NI-21831



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2014-2016** LIC. NO.

8013.64

Artwork shown at approximately 110%. PDF prints at approximately 100% from acrobat reader or when "Fit to printable area" settings are selected.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TO REFILL SPRAY BOTTLES: Remove: Remove trigger sprayer from empty bottle. Unscrew cap on refill and pour contents directly into empty bottle. Use a funnel to minimize product waste. **Use:** Replace trigger sprayer and use as you normally would. This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream, or normally sterile areas of the body, or (2) contacts intact mucous membranes, but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

SPECIAL INSTRUCTIONS FOR USE AGAINST HIV-1, HBV AND HCV: This product kills HIV-1, HBV and HCV on precleaned hard, nonporous surfaces/objects previously soiled with blood/body fluids in healthcare settings (hospitals, nursing homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus (HIV-1) (associated with AIDS), Human Hepatitis B Virus (HBV) and Human Hepatitis C Virus (HCV). **SPECIAL INSTRUCTIONS FOR USING THIS PRODUCT TO CLEAN AND DECONTAMINATE AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS: Personal Protection:** When handling items soiled with blood or body fluids, use disposable impervious gloves, gowns, masks and eye coverings. **Cleaning Procedure:** Blood and other body fluids must be thoroughly cleaned from surfaces and other objects before applying this product. **Contact Time:** Allow surface to remain wet for HBV, HCV and HIV-1 for 30 seconds. The contact times for other bacteria, viruses and fungi may differ. See product label for contact times. **DISPOSAL OF INFECTIOUS MATERIALS:** Use disposable impervious gloves, gowns, masks and eye coverings. Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

TO PRECLEAN INSTRUMENTS PRIOR TO TERMINAL STERILIZATION/HIGH-LEVEL DISINFECTION: As a precleaning spray – Place instruments into a suitable container. Spray Clorox Healthcare® Hydrogen Peroxide Cleaner Disinfectant onto instruments to thoroughly wet all surfaces. Let stand for up to 10 minutes. Rinse instruments. **A manual instrument cleaner** – Thoroughly pre-rinse dirty instruments under running water to remove visible, gross debris. Immerse pre-rinsed instruments into an appropriate size container filled with Clorox Healthcare® Hydrogen Peroxide Cleaner Disinfectant. Scrub instruments using a stiff bristle brush until visibly clean. Submerge instruments while scrubbing. Rinse instruments thoroughly. Change solution daily. Follow with an appropriate disinfection process. Cleaning of critical and semi-critical devices must be followed by an appropriate terminal sterilization/high-level disinfection process. **TO DISINFECT NON-CRITICAL, PRECLEANED INSTRUMENTS:** Instruments must be thoroughly precleaned to remove excess organic debris, rinsed and rough dried. (Clean and rinse lumens of hollow instruments before filling with Clorox Healthcare® Hydrogen Peroxide Cleaner Disinfectant.) Spray all surfaces of instruments with Clorox Healthcare® Hydrogen Peroxide Cleaner Disinfectant until thoroughly wet. Let stand for one minute. To kill TB, let stand for 4 minutes at room temperature. To kill mold and fungi, let stand for 3 minutes. Wipe with a clean, damp cloth or paper towel and allow to air dry.

DISINFECTION

This product cleans, disinfects and deodorizes hard, nonporous medical surfaces in one step with no rinsing required.

TO CLEAN, DISINFECT AND DEODORIZE HARD, NONPOROUS SURFACES: Spray 6–8 inches from surface until surface is thoroughly wet. Allow this product to remain wet for contact time listed on label. Then wipe. For heavily soiled areas, a precleaning is required. Rinse with potable water for food-contact surfaces. For all others, no rinsing is required.

SOFT SURFACE SANITIZATION

TO SANITIZE SOFT SURFACES: Hold bottle upright 6–8 inches from surface. Spray until fabric is wet. DO NOT SATURATE. Let stand for 30 seconds. Allow to air dry. Heavy soil must be removed prior to sanitizing. For first time use on soft surfaces, spot test in a small, inconspicuous area.

Efficacy tests have demonstrated that this product is an effective bactericide, fungicide, and virucide* in the presence of organic soil (5% blood serum).



ORGANISMS

BACTERIA: 30 second contact time

- *Acinetobacter baumannii*
- *Campylobacter jejuni*
- Carbapenem Resistant *Klebsiella pneumoniae*
- *Enterobacter aerogenes*
- *Escherichia coli* (E. coli)
- *Escherichia coli* New Delhi Metallo-Beta Lactamase-1 (NDM-1)
- *Escherichia coli* O157:H7
- Extended Spectrum Beta Lactamase producing *Escherichia coli* (ESBL producing E. coli)
- Extended Spectrum Beta Lactamase producing *Klebsiella pneumoniae* (ESBL producing *Klebsiella pneumoniae*)
- *Klebsiella oxytoca*
- *Klebsiella pneumoniae* New Delhi Metallo-Beta Lactamase-1 (NDM-1)
- Multidrug Resistant *Enterococcus faecium*
- Extended Spectrum Beta Lactamase producing *Escherichia coli* (ESBL producing E. coli)
- *Pseudomonas aeruginosa*
- *Salmonella enterica*
- *Stenotrophomonas maltophilia*
- Penicillin-Resistant *Streptococcus pneumoniae*
- *Streptococcus pyogenes*
- Vancomycin Resistant *Enterococcus faecalis* (VRE)

BACTERIA: 1 minute contact time

- *Staphylococcus epidermidis* (Coagulase-negative staphylococci) (CoNS)
- Community Acquired Methicillin Resistant *Staphylococcus aureus* (CA-MRSA)
- Linezolid Resistant *Staphylococcus aureus* (MRSA) (ATCC 33592)
- Methicillin Resistant *Staphylococcus aureus* (MRSA) (ATCC 33591)
- Methicillin Resistant *Staphylococcus aureus* (MRSA) (ATCC 33592)
- *Staphylococcus aureus*

BACTERIA: 4 minute contact time

- *Mycobacterium bovis* (TB)

VIRUSES: 30 second contact time

- *Avian Influenza A virus (H5N1)
- *Cytomegalovirus
- *Hepatitis B virus
- *Hepatitis C virus
- *Herpes simplex virus type 1
- *Herpes simplex virus type 2
- *Human Coronavirus
- *Human Immunodeficiency Virus type 1 (HIV-1)
- *Influenza A virus (H3N2)
- *Influenza A virus (2009 H1N1)
- *Influenza B virus
- *Respiratory syncytial virus (RSV)

VIRUSES: 1 minute contact time

- *Norovirus
- *Rhinovirus (type 37)
- *Rotavirus

MOLD, FUNGI: 3 minute contact time

- *Candida albicans*
- *Trichophyton mentagrophytes*

Soft Surface Sanitization: 30 second contact time

- *Enterobacter aerogenes*
- *Staphylococcus aureus*

Use Sites: Ambulances, Ambulatory Surgical Centers (ASC), CCUs, Dialysis clinics, Emergency rooms, Home Health Care, ICUs, Laboratories, Long-term care facilities, Clinics, Newborn Intensive Care Units (NICU), Nursing homes, Nurses' stations, Operating rooms, Patient restrooms, Patient rooms, Pediatric examination areas, Pediatric Intensive Care Units, Physical therapy areas, Physicians' offices, Public areas, Radiology rooms. **Medical Surfaces:** Bedrails, Bedside tables, Carts, Counters, High-touch surfaces¹, IV stands/pumps/poles, Large surfaces, Shower fixtures, Stretchers, Wheelchairs. **Surface Materials:** Baked enamel, Chrome, Glass, Glazed ceramic, Glazed porcelain, Laminated surfaces, Plastic, Stainless steel, Vinyl. **Soft Surfaces:** Use on Chairs², Couches³, Curtains⁴, Fabrics⁵, Upholstered furniture⁶, and other Soft surfaces⁷. **Avoid Contact With:** Aluminum, Copper, Galvanized steel, Silver. ¹Hard, nonporous surfaces of ²composed of 100% cotton or 100% polyester

CAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS.

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. **FIRST AID: If in eyes:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call the poison control center at (800) 222-1222 or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

STORAGE AND DISPOSAL: Store this product in a cool, dry area, away from direct sunlight and heat to avoid deterioration. Nonrefillable container. Do not reuse or refill empty container. Recycle empty container or discard in trash. **QUESTIONS OR COMMENTS?** Call (800) 234-7700. For MSDS information, please go to www.cloroxhealthcare.com.

Mid. for & © 2013 Clorox Professional Products Company, 1221 Broadway, Oakland, CA 94612.

Clorox Healthcare is a registered trademark of The Clorox Company. Made in the USA. EPA Reg. No. 67619-24, EPA Est. No. 5813-GA-1 (A4), GA-2 (VG); 8251-OH-1 (O6), WI-3 (KV), WI-5 (UQ) Beginning of batch code indicates Est. No. Visit us at www.cloroxhealthcare.com. U.S. Patent No. 7,658,953.



MI-21931
R0904040