

Oxivir® Tb Wipes

*Virucidal • Bactericidal • Fungicidal • Tuberculocidal

FOR INSTITUTIONAL USE

ACTIVE INGREDIENT:

Hydrogen Peroxide.....0.5%

OTHER INGREDIENTS:99.5%

TOTAL:.....100.0%

This is **not** a baby wipe.

Not for personal cleansing. Do not flush in toilet.

KEEP OUT OF REACH OF CHILDREN

Net Contents: 6.2 lbs. (2.8 kg)

160 Wipes, 11 in. x 12 in. (28 cm x 30 cm)



Sealed Air
Diversey Care

Diversey™



5627427

This product is a one-step disinfectant cleaner and odor neutralizer designed for general cleaning, disinfecting, and deodorizing of hard, non-porous environmental surfaces. It cleans quickly by removing dirt, grime, body oils and other common soils found in hospitals, nursing homes, office buildings, schools, day care centers, airports, cruise ships, correctional facilities, athletic facilities, health clubs, veterinary clinics, pet shops, public areas and restrooms.

This product's advanced quick cleaning formula eliminates the time and labor normally required for rinsing. It is designed for use on the following hard, non-porous environmental surfaces: vinyl, painted surfaces, plastic surfaces, glass, mirrors, glazed ceramic, glazed porcelain, chrome, aluminum, laminated surfaces and baked enamel surfaces associated with floors, walls, ceilings, tables, chairs, non-food contact countertops, telephones, fixtures, athletic equipment, exercise equipment, glazed tile, exterior surfaces of toilets, exterior surfaces of urinals, sinks, sink basins, faucets, tubs, shower rooms and locker room areas. Do not use on glasses, dishes and utensils. Not recommended for use on brass, copper or marble.

When used as directed, this product is highly effective against a broad-spectrum of pathogenic microorganisms.

BACTERICIDAL in the presence of 5% serum load and 1 minute contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

Bactericidal Activity: *Pseudomonas aeruginosa* (ATCC 15442), *Staphylococcus aureus* (ATCC 6538), *Salmonella enterica* (ATCC 10708) (formerly known as *Salmonella choleraesuis*), *Acinetobacter baumannii* (ATCC 19606), *Escherichia coli* O157:H7 (ATCC 35150), *Klebsiella pneumoniae*, (ATCC 4352), *Escherichia coli* with extended beta-lactamase resistance (ESBL), (ATCC BAA-196), *Shigella dysenteriae*, (ATCC 11835)

Antibiotic-Resistant Strains of Bacteria: *Enterococcus faecalis*, (ATCC 51575) (VRE), *Staphylococcus aureus*, (ATCC 33592) (MRSA), *Staphylococcus aureus* (CA-MRSA), (NARSA NRS 384)(Genotype USA300) Community Associated Methicillin Resistant, *Staphylococcus aureus* (CA-MRSA), (NARSA NRS 123)(Genotype USA400) Community Associated Methicillin Resistant

***VIRUCIDAL** in the presence of 5% serum load and 1 minute contact time at 20°C (68° F) on hard, non-porous environmental surfaces.

***Viruses:** *Hepatitis B Virus (HBV) (Acquired from Hepadnavirus Testing), *Hepatitis C Virus (HCV) (Acquired from National Veterinary Services Laboratories), *Herpes Simplex Virus, Type 1 (HSV-1) (ATCC VR-733),

*Herpes Simplex Virus, Type 2 (HSV-2) (ATCC VR-734), *HIV-1 (AIDS Virus), Strain HTLV-III_b (HIV-1) (Acquired from Adv. Biotechnologies, Inc.), *Human Coronavirus (ATCC VR-740), *Influenza A/Hong Kong (ATCC VR-544), *Norovirus (Feline Calicivirus, ATCC VR-782 as the surrogate), *Poliovirus Type 1, Strain Brunhilde (ATCC VR-1000), *Rotavirus WA (Acquired from University of Ottawa), *Adenovirus type 8, (ATCC VR-1368), *Rhinovirus Type 14, (ATCC VR-284)

*Kills Pandemic 2009 H1N1 Influenza A virus.

***Veterinary Virus:** *Avian Influenza A (H3N2)(ATCC VR-2072)

TUBERCULOCIDAL in the presence of a 5% serum load and 5 minute contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

Mycobacterium bovis (Tb) (OT 451C150)

FUNGICIDAL: This product is fungicidal against the pathogenic fungi, *Trichophyton mentagrophytes* (ATCC 9533) (the fungus which causes Athlete's Foot) in the presence of a 5% serum load and 10 minute contact time at 20°C (68°F) when used as directed on hard surfaces found in bathrooms, shower stalls, locker rooms, or other clean, non-porous, hard surfaces commonly contacted by bare feet.

See reference sheet for a complete list of organisms and additional directions for use.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for personal cleansing. **This is not a baby wipe.** Do not flush in toilet.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

For Use as a One-Step Cleaner/Disinfectant: Pre-clean heavily soiled areas. Pull towelette from dispenser (canister) and wipe hard, non-porous environmental surfaces. All surfaces must remain visibly wet for 1 minute. Use a 5 minute contact time for Tuberculosis. Use a 10 minute contact time for fungi. Allow to air dry or rinse with potable water if necessary.

NOTE: All food contact surfaces must be rinsed with potable water. Do not use on glassware, utensils, or dishes.

NOTE: When cleaning floors position wet-floor signs around area to be cleaned. Floors will be slippery when wet or contaminated with foreign materials. Promptly clean up spills and foreign materials.

For Use as a Restroom, Locker Room, Athletic Equipment One-Step

Cleaner/Disinfectant: Pre-clean heavily soiled areas. Pull towelette from dispenser (canister) and wipe hard, non-porous environmental surfaces. To disinfect, all surfaces must remain wet for 1 minute. Use a 5 minute contact time for Tuberculosis. Use a 10 minute contact time for fungi. Wipe surfaces dry or rinse or allow to air dry.

NOTE: When cleaning floors position wet-floor signs around area to be cleaned. Floors will be slippery when wet or contaminated with foreign materials. Promptly clean up spills and foreign materials.

***KILLS HIV AND HCV and HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (Hospitals, Nursing Homes) and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV), or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATING AGAINST HIV-1 (AIDS Virus), HBV OR HCV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood/body fluids must be thoroughly cleaned from surfaces/objects before application of this product.

Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1, HBV and HCV. Use a 5 minute contact time for Tuberculosis and a 10 minute contact for fungi.

Disposal of Infectious Material: Blood/body fluids must be autoclaved and disposed according to Federal, State and local regulations for infectious waste disposal.

©2015 Manufactured for:
Diversey, Inc., PO Box 19747, Charlotte, NC 28219-0747

For Use as a Tuberculocide: Pre-clean heavily soiled areas. Pull towelette from dispenser (canister) and wipe hard, non-porous environmental surfaces. Allow the surface to remain wet for 5 minutes. Use additional towelettes if needed to ensure the correct wet contact time. Wipe surfaces dry or rinse or allow to air dry. Change towelette after each use.

To Kill Fungi: Pre-clean heavily soiled areas. Pull towelette from dispenser (canister) and wipe hard, non-porous environmental surfaces. All surfaces must remain visibly wet for 10 minutes. Use additional towelettes if needed to ensure the correct wet contact time. Wipe surfaces dry or rinse or allow to air dry.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Store in original container in areas inaccessible to small children. Keep securely closed.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Wrap empty container and put in trash.

Towelette Storage/Disposal: Dispose of wipe in trash after use. Do not flush. Tightly close lid between uses to retain moisture.

ENVIRONMENTAL HAZARDS: This product is toxic to birds, fish and aquatic invertebrates. Caution should be used when applying indoors because pets may be at risk.

See SDS# MS0800545

EPA Reg. No.: 70627-60

EPA Est. No.: 74559-CAN-1 [VX]; 8251-WI-004 [KL]; 0312-WI-3 [NW]

Lot code letters indicate establishment number.

EMERGENCY PHONE : 1-800-851-7145

 **Questions? Comments**
1-800-558-2332

Product of Canada



5992643 (15/198)

Oxivir® Tb Wipes



BACTERICIDAL in the presence of 5% serum load and 1 minute contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

Bactericidal Activity:

Pseudomonas aeruginosa (ATCC 15442)

Salmonella enterica, (ATCC 10708)

(formerly known as *Salmonella choleraesuis*)

Staphylococcus aureus (ATCC 6538)

Acinetobacter baumannii (ATCC 19606)

Escherichia coli O157:H7 (ATCC 35150)

Klebsiella pneumoniae, (ATCC 4352),

Escherichia coli with extended beta-lactamase resistance (ESBL), (ATCC BAA-196)

Shigella dysenteriae, (ATCC 11835)

Antibiotic-Resistant Strains of Bacteria:

Enterococcus faecalis, (ATCC 51575) (Resistant to Vancomycin (VRE))

Staphylococcus aureus, (ATCC 33592) (Resistant to Methicillin (MRSA))

Staphylococcus aureus (CA-MRSA), (NARSA NRS 384)(Genotype USA300)

Community Associated Methicillin Resistant

Staphylococcus aureus (CA-MRSA), (NARSA NRS 123)(Genotype USA400)

Community Associated Methicillin Resistant

***VIRUCIDAL** in the presence of 5% serum load and 1 minute contact time at 20°C (68° F) on hard, non-porous environmental surfaces.

***Viruses:**

*Hepatitis B Virus (HBV) (Acquired from Hepadnavirus Testing)

*Hepatitis C Virus (HCV) (Acquired from National Veterinary Services Laboratories)

*Herpes Simplex Virus, Type 1 (HSV-1) (ATCC VR-733)

*Herpes Simplex Virus, Type 2 (HSV-2) (ATCC VR-734)

*Kills Pandemic 2009 H1N1 Influenza A virus.

*HIV-1 (AIDS Virus), Strain HTLV-III_B (HIV-1) (Acquired from Adv. Biotechnologies, Inc.)

*Human Coronavirus (ATCC VR-740)

*Influenza A/Hong Kong (ATCC VR-544)

*Norovirus (Feline Calicivirus, ATCC VR-782 as the surrogate)

*Poliovirus Type 1, Strain Brunhilde (ATCC VR-1000)

*Rotavirus WA (Acquired from University of Ottawa)

*Adenovirus type 8, (ATCC VR-1368)

*Rhinovirus Type 14, (ATCC VR-284)

***Veterinary viruses:**

*Avian Influenza A, (H3N2) (ATCC VR-2072)

*Feline Calicivirus, Strain F9 (ATCC VR-782)

TUBERCULOCIDAL in the presence of a 5% serum load and 5 minute contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

Mycobacterium bovis (Tb) (OT 451C150)

FUNGICIDAL: This product is fungicidal against the pathogenic fungi, *Trichophyton mentagrophytes* (ATCC 9533) (the fungus which causes Athlete's Foot) in the presence of a 5% serum load and 10 minute contact time at 20°C (68°F) when used as directed on hard surfaces found in bathrooms, shower stalls, locker rooms, or other clean, non-porous, hard surfaces commonly contacted by bare feet.

Trichophyton mentagrophytes (ATCC 9533)

Non-Food Contact Surface Sanitizer for hard non-porous surfaces – in the presence of a 5% serum load and 30 second contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

Klebsiella pneumoniae (ATCC 4352)

Staphylococcus aureus (ATCC 6538)

Escherichia coli O157:H7 (ATCC 35150)

Salmonella enterica, (ATCC 10708)

(formerly known as *Salmonella choleraesuis*)

Staphylococcus aureus MRSA (ATCC 33592)

Pseudomonas aeruginosa (ATCC 15442)

Enterococcus faecalis VRE (ATCC 51575)

Malodor Activity – eliminates odors and odor-causing bacteria on hard, non-porous surfaces in restroom areas, behind and under sinks and counters, and storage areas where bacterial growth can cause malodors.

For use as a Non-Food Contact Sanitizer:

1. Pre-clean heavily soiled areas.
2. Pull towelette from dispenser canister and wipe hard, non-porous environmental surfaces.
3. Allow the surface to remain wet for 30 seconds.
4. Wipe surfaces dry or rinse or allow to air dry.
5. Change towelette after each use.

For Use as a Cleaner and/or Deodorizer:

Pull towelette from dispenser (canister) and wipe hard, non-porous environmental surfaces. Allow to air dry or rinse with potable water if necessary.

Animal Housing Facilities Including Veterinary Clinics, Farms, Livestock, Swine and Poultry Houses: For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities (including pens, chutes, and barns) occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding or watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water. Pull towelette from dispenser (canister) and wipe hard, non-porous environmental surfaces.
5. All surfaces must remain visibly wet for 1 minute.
6. Allow to air dry or rinse with potable water if necessary.
7. Ventilate buildings and other closed spaces.
8. Do not house animals or employ equipment until treatment has been absorbed, set or dried.

NOTE: Not for use on handling and restraining equipment such as leashes, muzzles, halters or ropes, as well as forks, shovels, and scrapers used for removing litter and manure.