



STATE OF HAWAII
Department of Agriculture
ACCEPTED
LICENSE NO.
8017.160

Sealed Air
Food Care

Diversey™

Divosan® Hypochlorite

Liquid Chlorinated Sanitizer



Liquid Bactericide • Disinfectant • Sanitizer • Deodorizer

For Industry Use

ACTIVE INGREDIENT:

| | |
|---------------------------------|---------|
| Sodium Hypochlorite | 5.25% |
| OTHER INGREDIENTS: | 94.75% |
| TOTAL | 100.00% |

Available Chlorine - 5.0%

KEEP OUT OF REACH OF CHILDREN DANGER

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR
DOCTOR FOR TREATMENT ADVICE.
1-800-851-7145**

Have the product container or label with you when calling a
Poison Control Center or doctor or going in for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

See additional precautionary statements on side panel.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
Note: This product degrades with age. Use a chlorine test kit and increase/decrease dosage, as necessary to obtain the required level of available chlorine.

MANUAL CLEANING OF DAIRY EQUIPMENT, BULK TANKS AND UTENSILS.

Rinse equipment thoroughly, immediately after use or emptying of bulk tanks with warm (100°F) water. Drain. Wash equipment using an appropriate manual cleaner according to use directions. Brush all equipment thoroughly. Rinse suds away with cool water. Drain. Acid rinse equipment using an appropriate acid cleaner diluted 1 ounce per 10 gallons in tap water. Drain. After cleaning and just prior to use, sanitize equipment with 5 ounces of Divosan Hypochlorite per 10 gallons of tap water (200 ppm). Test solution periodically with a chlorine test kit. Expose equipment to sanitizing solution at least 2 minutes. Drain.

SANITIZATION OF CONVEYORS FOR MEAT, POULTRY, SEAFOOD, FRUITS, AND VEGETABLES

For use in the static or continuous washing, rinsing and sanitizing of conveyor equipment, peelers, collators, slicers, saws, etc. Remove all products from equipment if during treatment the sanitizer will directly contact the items.

Prepare 200 ppm solution of this product (5 oz / 10 gallons).

Apply sanitizer solution to the return portion of the conveyor or to the equipment using a coarse spray, other means of wetting the surfaces. Treat for at least one (1) minute. Control the volume of solution so as to permit maximum drainage and to prevent puddles. The conveyor may still be damp when food contact occurs.

Allow equipment to drain adequately before reusing; a dry surface is not required.

SANITIZATION OF NON-POROUS FOOD CONTACT SURFACES:

Dairy, Beverage, Meat, Poultry, Commissary and Food Processing Plants - RINSE, IMMERSER OR FLOOD APPLICATIONS. A solution of 100 ppm available chlorine may be used in the sanitizing solution if a test kit is available. Solutions containing an initial concentration of 100 ppm available chlorine must be tested and adjusted periodically to insure that the available chlorine does not drop below 50 ppm. Prepare a 100 ppm sanitizing solution by thoroughly mixing 2.5 oz of this product in 10 gallons of water. If no test kit is available, prepare a sanitizing solution by thoroughly mixing 5 oz of this product in 10 gallons of water to provide approximately 200 ppm available chlorine by weight.

Clean equipment surfaces in the normal manner. Prior to use, rinse, immerse or flood surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. If the solution contains less than 50 ppm available chlorine, as determined by a suitable test kit, either discard the solution or add sufficient product to reestablish 200 ppm residual. Do not rinse equipment with water after treatment and do not soak equipment overnight.

Sanitizers used in automated systems may be used for general cleaning but may not be re-used for sanitization purposes.

SANITIZATION OF NON-POROUS NON-FOOD CONTACT SURFACES:

Dairy, Beverage, Meat, Poultry, Commissary and Food Processing Plants - RINSE, IMMERSER OR FLOOD APPLICATIONS. Prepare a sanitizing solution by thoroughly mixing 5 oz of this product with 10 gallons of water to provide approximately 200 ppm available chlorine by weight. Clean equipment surfaces in the normal manner. Prior to use, rinse, immerse or flood surfaces thoroughly with the sanitizing solution, maintaining contact with the solution for at least 2 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight.

SANITIZING POROUS FOOD CONTACT EQUIPMENT:

Prepare a 600 ppm solution by thoroughly mixing 15 ounces of this product in 10 gallons of water. Clean surfaces in the normal manner. Rinse, immerse or flood all surfaces thoroughly with the 600 ppm solution, maintaining contact for at least 2 minutes. Prior to using equipment, rinse all surfaces with a 200 ppm available chlorine solution (5 oz / 10 gallons). Do not rinse and do not soak overnight.

SANITIZATION OF POROUS NON-FOOD CONTACT SURFACES:

Dairy, Beverage, Meat, Poultry, Commissary and Food Processing Plants - RINSE, IMMERSER OR FLOOD APPLICATIONS. Prepare a sanitizing solution by thoroughly mixing 7.5 oz of this product with 5 gallons of water to provide approximately 600 ppm available chlorine by weight. Clean surfaces in the normal manner. Prior to use, rinse, immerse or flood all surfaces thoroughly with the sanitizing solution, maintaining contact with the solution for at least 2 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight.

DISINFECTION OF NON-POROUS NON-FOOD CONTACT SURFACES:

Dairy, Beverage, Meat, Poultry, Commissary and Food Processing Plants - RINSE, IMMERSER OR FLOOD APPLICATIONS. Prepare a disinfecting solution by thoroughly mixing 7.5 oz of this product with 5 gallons of water to provide approximately 600 ppm available chlorine. Clean equipment in the normal manner. Prior to use, rinse, immerse or flood surfaces thoroughly with the disinfecting solution, maintaining contact with the solution for at least 10 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight.

SPRAY AND FOGGING SANITIZATION OF NON-POROUS FOOD CONTACT SURFACES:

Pre-clean all surfaces after use. Use a 200 ppm available chlorine solution to control bacteria, mold or fungi and a 600 ppm solution to control bacteriophage. Prepare a 200 ppm sanitizing solution of sufficient size by thoroughly mixing the production in a ratio of 5 oz. product with 10 gallons of water. Prepare a 600 ppm solution by thoroughly mixing the product in a ratio of 15 oz. per 10 gallons of water. Use spray or fogging equipment which can resist hypochlorite solutions. Always empty and rinse spray/fog equipment with potable water after use. Thoroughly spray or fog all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours. Prior to using equipment, rinse all surfaces treated with a 600 ppm solution with a 200 ppm solution.

FOOD EGG SANITIZING – Thoroughly clean all eggs. Dilute 5 oz. of product for each 10 gallons of warm water to produce 200 ppm available chlorine solution. The sanitizer temperature should not exceed 130° F. Spray the warm sanitizer so that all eggs are thoroughly wetted. Allow the eggs to completely dry before casing or breaking. Do not apply a potable water rinse. Do not reuse this solution for sanitizing eggs.

FRUIT AND VEGETABLE WASHING – Thoroughly clean all fruit and vegetables by immersion in recommended product solution in a sink or spray washer. Dilute 0.5 ounce of product in eight gallons of water to make a sanitizing solution containing 25 ppm available chlorine. Drain and rinse product with potable water. Immerse or spray in a separate sink with dilute sanitizer for 2 minutes. Check available chlorine content with test kit. Spray rinse vegetables with additional sanitizer solution prior to packaging. Rinse fruits and vegetables with potable water prior to use.


TREATMENT OF POULTRY PROCESSING WATER

Follow guidelines of local water authority for water potability treatment.

Continuous Feed: Using an automatic metering device, continuously feed this product into the water to obtain and/or maintain a level of 5-20 ppm available chlorine (1.25 oz product per 100 gal. water, to 5 oz product per 100 gals. water). Confirm target chlorine level with either a chlorine test kit or an automatic testing device. When the available chlorine level reaches 10 ppm, notify the USDA plant inspector.

Intermittent Feed: Start up by adding 2.5 oz of product per 1000 gallons of water for each 1 ppm of available chlorine needed. For subsequent doses, check chlorine level with a chlorine test kit. Add enough of this product to maintain the target chlorine level and confirm this level with a chlorine test kit. Do not pour this product directly on poultry product in the water.

HAND SANITIZING: Thoroughly wash hands with soap or detergent. Rinse with potable water. Prepare a solution containing 1 oz of this product per 8 gallons of water providing 50 ppm available chlorine (use test kit to determine exact available chlorine). Submerge hands. Let air dry.



SKU #: 57465100

Net Contents:

Product No:

18.9 L / 5 U.S. Gal.

57465100

LOT #:

COOLING TOWER/EVAPORATIVE CONDENSER WATER: CONTINUOUS FEED

METHOD:

Initial Dose: When system is noticeably fouled, apply 125 to 250 oz. of this product per 10,000 gallons of water in the system to obtain 5 to 10 ppm available chlorine.

Subsequent Dose: Maintain this treatment level by starting a continuous feed of 2.5 oz. of this product per 1,000 gallons of water lost by blowdown to maintain a 1 ppm residual. Badly fouled systems must be cleaned before treatment is begun.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store this product in a cool dry area, away from direct sunlight and heat to avoid deterioration.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic organisms. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting agency has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Causes irreversible eye damage and skin burns. Harmful if swallowed.

Wear chemical splash-proof goggles or face shield, rubber gloves and protective clothing when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until strong odors have dissipated.

PHYSICAL AND CHEMICAL HAZARDS

STRONG OXIDIZING AGENT. Mix only with water according to label directions. Mixing this product with chemicals (e.g. ammonia, acids, detergents, etc.) or organic matter (e.g. urine, feces, etc.) will release irritating chlorine gases which are harmful to eyes, lungs and mucous membranes.

NOTE: This product degrades with age. Use a chlorine test kit and increase dosage, as necessary, to obtain the required level of available chlorine.



EPA Registration No. 875-93

EPA Establishment No.

3635-OH-1 [SV]; 875-WI-1 [EC]; 875-CA-1 [MD]; 81642-OH-1 [HV];
13938-CAN-1 [LD]; 10508-TN-1 [RC]; 10508-TX-1 [RD]; 10508-OH-1 [RM];
10508-PA-1 [SB]; 0312-WI-3 [NW]; 7546-WI-1 [WT]; 34490-CA-1 [RH]

Lot code letters indicate establishment number

Divosan® Hypochlorite

INDIVIDUAL WATER SYSTEMS: DRILLED & BORED WELLS: Run pump until water is as free from turbidity as possible. Pour a 100 ppm available chlorine sanitizing solution into the well made by thoroughly mixing 2.5 ounces of Divosan Hypochlorite into 10 gallons of water. Add 5 to 10 gallons of clean, chlorinated water to the well in order to force the sanitizer into the rock formation. Wash the exterior of pump cylinder with the sanitizer. Drop pipeline into well, start pump and pump water until strong odor of chlorine in water is noted. Stop pump and wait at least 24 hours. After 24 hours, flush well until all traces of chlorine have been removed from the water. Deep wells with high water levels may necessitate the use of special methods for introduction of the sanitizer into the well. Consult your local Health Department or further details.

DISINFECTION OF DRINKING WATER (PUBLIC SYSTEMS):

Prepare a stock disinfectant solution by mixing 1-1/4 oz. of product with 5 gallons water to obtain 100 ppm available chlorine. Meter the stock solutions with hypochlorinator until a free residual chlorine content of at least 0.2 ppm but no more than 0.6 ppm is obtained throughout the distribution system. Check water frequently with a Chlorine Test Kit. Bacteriological sampling must be conducted at a frequency no less than that prescribed by the National Primary Drinking Water Regulation. Contact your local Health Department for further details.

COMMERCIAL LAUNDRY SANITIZERS: Wet fabrics or clothes should be spun dry prior to sanitization.

Thoroughly mix 5 oz. of this product with 10 gallons of water to yield 200 ppm available chlorine. Promptly after mixing the sanitizer, add the solution into the prewash prior to washing fabrics/clothes in the regular wash cycle with a good detergent. Test the level of available chlorine if solution has been allowed to stand. Add more of the product if the available chlorine level has dropped below 200 ppm.

FARM PREMISES: Remove all animals, poultry, and feed from premises, vehicles and enclosures. Remove all litter, and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities occupied or transversed by animals or poultry. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. To disinfect, saturate all surfaces with a solution of at least 1000 ppm available chlorine for a period of 10 minutes. A 1000 ppm solution can be made by thoroughly mixing 25 oz of this product with 10 gallons of water. Immerse all halters, ropes and other types of equipment used in handling and restraining animals or poultry, as well as the cleaned forks, shovels and scrapers used for removing litter and manure. Ventilate buildings, cars, boats and other closed spaces. Do not house livestock or poultry or employ equipment until chlorine has dissipated. All treated feed racks, mangers, troughs, automatic feeders, fountains and waterers must be rinsed with potable water before use.

See container label for First Aid, Precautionary Statements and complete Directions for Use.

EPA Registration No: 875-93