

# Avert™

# Diversey

## Sporicidal Disinfectant Cleaner

Effective in 4 minutes against *C. difficile* spores

**ACTIVE INGREDIENT:**

Sodium hypochlorite ..... 1.312%

**OTHER INGREDIENTS** ..... 98.688%

**TOTAL** ..... 100.000%

1.25% available Chlorine

### KEEP OUT OF REACH OF CHILDREN DANGER

See additional precautionary statements on back panel.

**FIRST AID**

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes.

Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center for treatment advice.

**IF SWALLOWED:** Call poison control center or doctor immediately for treatment advice.

Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

**IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR DOCTOR FOR FURTHER TREATMENT ADVICE.**

1-800-851-7145

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

Net Contents

946 mL / 1 U.S. Qt.

100842725

### Cleans, Deodorizes, Disinfects

\*Virucidal Fungicidal,

Sporicidal against *Clostridium difficile* spores

Cleans, disinfects, and deodorizes in one labor saving step.

See reference sheet for a complete list of claims and additional directions for use.

This product is a one-step, hospital disinfectant cleaner designed for general cleaning, disinfection and deodorizing of hard, nonporous surfaces.

**DIRECTIONS FOR USE:**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

Not recommended for use on: aluminum, brass, chipped enamel, clear plastic, copper, finished floors, natural marble, painted surfaces, and silver.

**Special Instructions for Cleaning Prior to Disinfection against *Clostridium difficile* spores**

**Personal Protection:** Wear appropriate barrier protection such as gloves, gowns, masks or eye covering.

**Cleaning Procedure:** Fecal matter/waste must be thoroughly cleaned from surfaces/objects before disinfection by application with a clean cloth, mop, and/or sponge saturated with the sporicidal product. Cleaning is to include vigorous wiping and/or scrubbing, until all visible soil is removed. Special attention is needed for high-touch surfaces. Surfaces in patient rooms are to be cleaned in an appropriate manner, such as from right to left or left to right, on horizontal surfaces, and top to bottom, on vertical surfaces, to minimize spreading of the spores. Restrooms are to be cleaned last. Do not reuse soiled cloths.

**Infectious Materials Disposal:** Materials used in the cleaning process that may contain feces/wastes are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

**Disinfection Procedure:** Apply product by coarse trigger sprayer to hard, nonporous inanimate surfaces. Spray 6 – 8 inches from surface, making sure to wet surfaces thoroughly. Surfaces must remain wet for 4 minutes. Wipe surfaces and let air dry.

EPA Reg. No. 70627-72

EPA Est. No. 0312-WI-3 (NW); 7546-WI-1 (WT)

Lot code letters indicate establishment number.

Not for sale or use on or after:



STATE OF HAWAII  
Department of Agriculture

**ACCEPTED**

LICENSE NO.

**8017.167**

**\*Kills HBV, HCV and HIV-1 on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids** in health care settings (Hospitals, Nursing Homes) and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Hepatitis B Virus, Hepatitis C Virus and Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.**

**Personal Protection:** Specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable latex gloves, gowns, masks and eye coverings.

**Cleaning Procedure:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of the disinfectant.

**Contact Time:** Allow surface to remain wet for 1 minute. For *Clostridium difficile* spores, allow to remain wet for 4 minutes.

**Disposal of Infectious Materials:** Blood and other body fluids, cleaning materials and clothing must be autoclaved and disposed of according to Federal, state and local regulations for infectious waste disposal.

### STORAGE AND DISPOSAL

Do not contaminate food or feed by storage, disposal, or cleaning of equipment.

**STORAGE:** Store this product in a cool dry area, away from direct sunlight and heat to avoid deterioration. In case of spill, flood areas with large quantities of water.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Wrap empty container and put in trash or offer for recycling if available.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**DANGER:** Corrosive. Causes skin burns and eye damage. Do not get in eyes, on skin, or on clothing. Wear chemical-splash goggles, chemical-resistant gloves and protective footwear. In case of inadequate ventilation wear respiratory protection. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

### PHYSICAL OR CHEMICAL HAZARDS

Keep away from clothing and other combustible materials. Do not mix with any other product or chemical.



8

07174 53060

5

300152457 (15/250)

# Avert™

## Sporicidal Disinfectant Cleaner



This product can be used in the following areas: alternative care facilities, ambulances, ambulatory surgery centers, anesthesia rooms, assisted living facilities, CAT laboratories, central service areas, clinics, critical care units, dental offices, exam rooms, emergency rooms, dialysis clinics, blood donation centers, home health care, hospices, hospitals, intensive care units, isolation rooms, laboratories, long-term care facilities, medical facilities, doctors' offices, nurseries, nursing homes, operating rooms, patient rooms, radiology rooms, rehabilitation centers, airports, animal hospitals, athletic facilities, bathrooms, correctional facilities, cruise ships, food service, funeral homes, health clubs, hotels, institutions, locker rooms, office buildings, recreational facilities, retail businesses, schools, transportation terminals, and veterinary clinics.

For Use On: acrylic, baked enamel surfaces, Corian®, glass, glazed ceramic, glazed porcelain, glazed tile, laminated surfaces, sealed mirror stone, painted surfaces, plastic surfaces, quarry tiles, sealed granite, sealed marble, stainless steel, finished terrazzo and vinyl.

Not recommended for use on: aluminum, brass, chipped enamel, clear plastic, copper, finished floors, natural marble, painted surfaces, and silver.

**Note:** This product is compatible with the listed materials. If product is intended to be used on other surfaces, it is recommended to test product by applying to a small discrete area to determine compatibility before proceeding with its use.

It can be used on: exterior surfaces of autoclaves, backboards, hospital beds, bedpans, bedside tables, blood pressure cuffs, blood pressure monitors, cabinets, call buttons, computerized axial tomography equipment (CATs), chemotherapy hoods, cellular phones, chairs, closets, coated covers for mattresses and pillows, remote controls, counters, desktops, diagnostic equipment, exterior surfaces of dialysis machines, docking stations, exterior surfaces of medical equipment, faucets, exterior of glucometers, gurneys, handheld electronic devices, handles, headboards, high touch surfaces, isolettes, IV stands and exterior surfaces of pumps, laptops, loupes, mammography equipment, exterior surfaces of MRIs, mobile workstations, operating room tables, orthopedics, pagers, patient care equipment, patient support and delivery equipment, phlebotomy trays, phones, hard nonporous edges of privacy curtains, pulse oximeters, scales, sequential compression devices, side rails, sinks, slit lamps, stethoscopes, stools, stretchers, support bars, tables, exterior surfaces of toilets, toilet grab bars, traction devices and patient slings, urinals, walls, wash basins, wheelchairs, weighing devices, and X-Ray equipment.

### DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Using approved Standard Quantitative Disk Carrier Test Method, in the presence of soil and 4 minute contact time, this product kills and inactivates spores of *Clostridium difficile* (ATCC 43598) on hard, nonporous surfaces.

When used as directed, this product is highly effective against a broad-spectrum of pathogenic microorganisms including bacteria, antibiotic-resistant bacteria, \*viruses, and fungi. Using approved AOAC test methods, in the presence of 5% serum load and 1 minute contact time, this product kills the following on hard nonporous inanimate surfaces:

#### Bacteria:

<i>Pseudomonas aeruginosa</i> , (ATCC 15442)	<i>Campylobacter jejuni</i> , (ATCC 29428)	<i>Listeria monocytogenes</i> , (ATCC 19117)
<i>Staphylococcus aureus</i> , (ATCC 6538)	<i>Escherichia coli</i> , (ATCC 11229)	<i>Stenotrophomonas maltophilia</i> , (ATCC 25596)
<i>Salmonella enterica</i> , (ATCC 10708)	<i>Escherichia coli</i> O157:H7, (ATCC 35150)	<i>Streptococcus pyogenes</i> , (ATCC 19615)

#### Antibiotic-Resistant Bacteria:

<i>Acinetobacter baumannii</i> , Multi-Drug Resistant (MDRO) (ATCC 19606)	<i>Escherichia coli</i> , Extended-Spectrum Beta-lactamase (ESBL) producing (ATCC BAA-196)	<i>Klebsiella pneumoniae</i> , NDM-1 positive (CDC 1000527)
<i>Enterobacter aerogenes</i> , Multi-Drug Resistant (MDRO) (ATCC 29751)	<i>Klebsiella pneumoniae</i> , Carbapenem Resistant (CRKP) (ATCC BAA-1705)	<i>Pseudomonas aeruginosa</i> , Multi-Drug Resistant Metallo-beta lactamase (MBL) positive (CDC 2012059)
<i>Enterococcus faecalis</i> , Vancomycin Resistant (VRE) (ATCC 51575)	<i>Klebsiella pneumoniae</i> , Extended-Spectrum Beta-lactamase (ESBL) producing (ATCC 700603)	<i>Staphylococcus aureus</i> , Methicillin Resistant (MRSA) (ATCC 33592)
<i>Enterococcus faecium</i> , Multi-Drug Resistant (ATCC 51559)	<i>Klebsiella pneumoniae</i> , Multi Drug Resistant (ATCC 51503)	<i>Staphylococcus aureus</i> , Community Acquired Methicillin Resistant (CA-MRSA) Genotype USA 300, (NRS 384)
<i>Escherichia coli</i> , Carbapenem Resistant (CRE) (CDC 81371)		

#### \*Viruses:

*Adenovirus Type 2, (ATCC VR-846, Strain Adenoid 6)	*Herpes Simplex Virus Type 2 (ATCC VR-734, Strain G)	*Norovirus, Feline Calicivirus as a Surrogate for
*Adenovirus Type 4, (ATCC VR-4, Strain RI-67)	*Human Coronavirus, (ATCC VR-740, Strain 229E)	*Poliovirus Type 1
*Hepatitis A Virus, (Strain HM-175)	*Human Immunodeficiency Virus (AIDS virus) (HTLV-III <sub>b</sub> )	*Respiratory Syncytial Virus (RSV) (ATCC VR-26, Strain Long)
*Hepatitis B Virus (HBV)	*Influenza A Virus, 2009-H1N1 (Novel H1N1), Strain A/Mexico/4108/2009 CDC# 2009712192	*Rhinovirus Type 37, (ATCC VR-1147, Strain 151-1)
*Hepatitis C Virus (HCV)		*Rotavirus, Strain WA

#### \*Veterinary Viruses:

*Avian Influenza A Virus, H5N1 (Strain VNHSN1-PR8/CDC-RG, CDC# 2006719965)	*Canine Parvovirus, (ATCC VR-2017, Strain Cornell-780916-80)	*Feline Calicivirus
--	--	---------------------

#### Fungi:

This product is fungicidal when used as directed on hard, nonporous inanimate surfaces in the presence of a 5% serum load and 1 minute contact time.

<i>Trichophyton mentagrophytes</i> , (ATCC 9533)	<i>Aspergillus niger</i> , (ATCC 6275)
--	--

#### Yeast:

*Candida albicans*, (ATCC 10231)

#### For Use as a One-Step Cleaner/Disinfectant, Deodorizer, and Fungicide:

1. Pre-clean heavily soiled areas.
2. Apply product by coarse trigger sprayer to hard, nonporous inanimate surfaces. Spray 6 – 8 inches from surface, making sure to wet surfaces thoroughly. Wipe or rub with cloth, disposable wiper, sponge, brush, or hand mop to spread product on surface.
3. Surfaces must remain wet for one (1) minute.
4. Wipe surfaces and let air dry or, if desired, remove solution and entrapped soil with a clean wet hand mop, cloth or wet vacuum pickup.

**Animal Housing Facilities:** For cleaning and disinfecting the following hard, nonporous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities including pens, chutes, and barns occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding or watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water. Saturate surfaces by spraying with product, allowing to remain wet for 1 (one) minute.
5. Spray handling and restraining equipment such as leashes, muzzles, halters or ropes, as well as forks, shovels, and scrapers used for removing litter and manure. Allow to remain wet for 1 minute.
6. Ventilate buildings and other closed spaces.
7. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub treated feed racks, troughs, mangers, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

See container label for First Aid, Precautionary Statements and complete Directions for Use.

EPA Reg. No. 70627-72

Manufactured for: ©2015 Diversey, Inc., PO Box 19747, Charlotte, NC 28219-0747