

***KILLS HIV, HCV and HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (Hospitals, Nursing Homes) and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis C Virus (HCV) or Hepatitis B Virus (HBV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS), HCV OR HBV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood/body fluids must be thoroughly cleaned from surfaces/objects before application of this product.

Contact Time: Allow surface to remain wet for 30 seconds to kill HIV-1, HCV and HBV, 4 minutes for mold, and 1 minute to kill all other organisms cited on label.

Disposal of Infectious Material: Blood/body fluids must be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

To Kill Fungi: Pre-clean heavily soiled areas. Spray 6 – 8 inches from hard, nonporous surface until completely wet. Allow to remain wet for 30 seconds. Wipe surfaces dry or allow to air dry.

To Use as a Non-Food Contact Sanitizer: Pre-clean heavily soiled areas, Apply product with trigger sprayer, cloth or disposable wipe to hard, non-porous environmental surfaces. Allow the surface to remain wet for 10 seconds. Wipe surfaces dry or allow to air dry.

To Sanitize Soft Surfaces: Remove gross filth prior to sanitization. Hold bottle upright 6 – 8 inches from surface. Spray until fabric is wet. DO NOT SATURATE, Allow surface to remain wet for 30 seconds. Let air dry.

†plain cotton weave and 100% polyester fabric

To Clean and Destroy Specified Allergens: Spray, wait 1 minute, and wipe excess. Allow to air dry or rinse. If streaking is observed, wipe with a clean, damp paper towel.

Allergens: Cockroach matter -or- particles, Dust mite matter -or- particles, Pet dander -or- dog and cat dander

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in areas inaccessible to small children. Keep securely closed. Store in a dry, well-ventilated area away from chemicals, direct light, heat or open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. For more information on the disposal of unused, unwanted product and the clean up of spills, contact the EPA Regional Office of the Manufacturer for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill empty container. Offer empty container for recycling, if available, or discard in trash.

ENVIRONMENTAL HAZARDS: This product is toxic to birds, fish and aquatic invertebrates. Caution should be used when applying indoors because pets may be at risk.

EPA Reg. No. 70627-74

EPA Est. No. 74559-CAN-1 (VX)

(Lot code letters indicate establishment number.)

Medical Emergency: 1-800-851-7145

Sealed Air
Diversey Care

Diversey™

Oxivir® 1

ONE-STEP READY TO USE HOSPITAL DISINFECTANT CLEANER

Active Ingredient:

Hydrogen Peroxide 0.5%

Other Ingredients: 99.5%

Total: 100.0%

BACTERICIDAL: 1 MINUTE

VIRUCIDAL: 30 SECONDS

ANIMAL PREMISE *VIRUCIDAL: 1 MINUTE

BROAD-SPECTRUM NON-FOOD CONTACT

SANITIZER: 10 SECONDS

SOFT SURFACE SANITIZATION: 30 SECONDS

FUNGICIDAL: 30 SECONDS

See reference sheet for a complete list of organisms and additional directions for use

KEEP OUT OF REACH OF CHILDREN



Department of Agriculture
STATE OF HAWAII

LICENSED

8017.170

PERIOD 2015-2017 LIC. NO.

Made in Canada

Net Contents/Contenu net:

946 mL / 1 U.S. Qt.



*AHP is a registered trademark of Virox Technologies Inc., used under license.

100850916

For Hospital, Health Care, Industrial, & Institutional Use

This product cleans, disinfects, and deodorizes hard, nonporous inanimate environmental surfaces and patient care equipment in one step with no rinsing required.

This product can be used in the following areas: long-term care facilities, medical facilities, operating rooms, animal treatment facilities, cruise ships, hotels, public buildings, restrooms, recreational facilities, retail businesses, and schools.

For use on medical, hard, nonporous inanimate surfaces: acrylic, chrome, glass, glazed ceramic tile, glazed porcelain laminated surfaces, and linoleum.

It can be used on the following hard, nonporous surfaces: hospital beds, blood pressure cuffs, exterior surfaces of medical equipment, patient care equipment, stethoscope, chairs, counters, floors, hand rails, high touch surfaces, showers, tables and walls. Do not use on glasses, dishes and utensils.

Eliminates the most common allergens**

Reduces allergens**

**This product removes the following non-living allergens: cat dander, cockroach matter, cockroach debris, cockroach particles, dog dander, dust mite debris, pet dander, pollen particles.

This product kills germs and destroys allergens** while cleaning hard, nonporous surfaces

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not recommended for use on: brass or copper.

When used as directed, this product is highly effective against a broad-spectrum of pathogenic microorganisms including bacteria, antibiotic-resistant bacteria, viruses, and fungi. Using approved AOAC test methods, in the presence of 5% serum load and 1 minute contact time, unless otherwise noted, this product kills the following on hard nonporous environmental surfaces: *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Staphylococcus aureus* Methicillin Resistant (MRSA), *Staphylococcus aureus* Vancomycin Intermediate Resistant (VISA), *Staphylococcus aureus* Vancomycin Resistant (VRSA).

*Virucidal, using approved AOAC methods, in the presence of 5% serum load with a 30 second contact time, unless otherwise noted. *Poliovirus Type 1, *Rhinovirus Type 37, *Norovirus (Feline Calicivirus, as the surrogate). *Animal Viruses with a 1 minute contact time: *Canine parvovirus

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

To Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Spray 6 – 8 inches from hard, non-porous inanimate surfaces, making sure to wet thoroughly.
3. Allow surface to remain wet for 1 minute.
4. Allow to air dry. If streaking is observed, wipe with a clean, damp cloth.

Note: Rinsing is not necessary unless floors are to be coated with finish or restorer. All food contact surfaces must be rinsed with potable water. Do not use on glassware, utensils, or dishes.

Manufactured for: ©2015 Diversey, Inc.,
PO Box 19747, Charlotte, NC 28219-0747

6085851 (15/230)



8

07 174 53 162

6

Oxivir[®] 1

When used as directed, this product is highly effective against a broad-spectrum of pathogenic microorganisms including bacteria, antibiotic-resistant bacteria,*viruses, and fungi. Using approved AOAC test methods, in the presence of 5% serum load and 1 minute contact time, unless otherwise noted, this product kills the following on hard nonporous inanimate environmental surfaces:

Bacteria:

Pseudomonas aeruginosa, (ATCC 15442)
Staphylococcus aureus, (ATCC 6538)
Salmonella enterica, (ATCC 10708)
 formerly known as *Salmonella choleraesuis*
Bordetella bronchiseptica, (ATCC 10580)
Burkholderia cepacia, (ATCC 25416)
Bordetella pertussis, (ATCC 12743)

Enterobacter aerogenes, (ATCC 13048)
Enterococcus faecium, (ATCC 51559)
Escherichia coli, (ATCC 11229)
Escherichia coli O157:H7, (ATCC 35150)
Klebsiella pneumoniae, (ATCC 4352)
Legionella pneumophila, (ATCC 33153)
Listeria monocytogenes, (ATCC 19117)

Pasteurella multocida, (ATCC 12947)
Staphylococcus pseudintermedius, (ATCC 49444)
Stenotrophomonas (Pseudomonas) maltophilia, (ATCC 25596)
Streptococcus pyogenes, (ATCC19615)

Antibiotic-Resistant (Strains of) Bacteria (Antibiotic-Resistant Bactericidal Activity):*Acinetobacter baumannii*,

Multi-Drug Resistant MDR (ATCC 19606)
 Ampicillin, Cefazolin, Cefepime, Gentamicin,
 Ertapenem, and Trimethoprim/Sulfa

Enterobacter cloacae, (New Delhi metallo-β-lactamase

1 positive (producing) (NDM 1)) (CDC 1000654)

Enterococcus faecalis, (ATCC 51575) (Vancomycin Resistant (VRE))

Escherichia coli, Extended Spectrum Beta-lactamase

producing (positive) (ESBL) (ATCC BAA-196)

Escherichia coli, New Delhi metallo-β-lactamase 1 (NDM-1) producing
 positive (CDC 1001728)

Klebsiella pneumoniae, (Carbapenem Resistant (CRE)) (ATCC BAA 1705)

Klebsiella pneumoniae, (Extended Spectrum beta-lactamase (ESBL) producing
 (positive)) (ATCC700603)

Klebsiella pneumoniae, (New Delhi metallo-β-lactamase 1(NDM-1)

producing (positive)) (CDC 1000527)

Staphylococcus aureus, Community Associated Methicillin Resistant (CA-MRSA)

(NARSA NRS 123) (Genotype USA400)

Staphylococcus aureus, (Methicillin Resistant (MRSA)) (ATCC 33591)

Staphylococcus aureus, (ATCC 33592) (Methicillin Resistant (MRSA))

Staphylococcus aureus, (Vancomycin Intermediate Resistant (VISA)) [HIP 5836]

Staphylococcus aureus, (Vancomycin Resistant (VISA))[VRS1]

Staphylococcus epidermidis, (Methicillin Resistant (MRSE)) (ATCC 51625)

Streptococcus pneumoniae, (Penicillin Resistant (PRSP)) (ATCC 700677)

***Virucidal** using approved AOAC methods, in the presence of 5% serum load with a 30 second contact time, unless otherwise noted.

***Viruses:**

*Adenovirus Type 8, (ATCC VR-1368)

*Cytomegalovirus, (ATCC VR-538)

*Hepatitis B Virus (HBV), Duck Hepatitis B Virus as surrogate
 (Acquired from Hepadnavirus Testing)

*Hepatitis C Virus (HCV), Bovine Viral Diarrhea

Virus as surrogate (Acquired from National Veterinary
 Services Laboratories)

*Herpes Simplex Virus, Type 1 (HSV-1) (ATCC VR-733)

*Herpes Simplex Virus, Type 2 (HSV-2), (ATCC VR-734)

*HIV-1 (AIDS Virus), Strain HTLV-IIIB (HIV-1),
 (Acquired from Adv. Biotechnologies, Inc.)

*Human Coronavirus, (ATCC VR-740)

*Influenza A Virus (Hong Kong Strain), (ATCC VR-544)

*Influenza B Virus, (ATCC VR-823)

*Norovirus, (Feline Calicivirus, ATCC VR-782 as the surrogate)

*Pandemic Influenza A Virus H1N1

*Parainfluenza Virus Type 3, (ATCC VR-93)

*Poliovirus Type 1, Strain Brunhilde, (ATCC VR-1562)

*Respiratory Syncytial Virus, (RSV) (ATCC VR-26)

*Rhinovirus Type 37, Strain 151-1, (ATCC VR-1147)

*Rotavirus, (WA strain) (Acquired from University of Ottawa)

*Vaccinia Virus, (ATCC VR-119)

***Animal Viruses:**

*Avian Influenza A Virus (H7N9)

*Avian Influenza A Virus (H3N2), (ATCC VR-2072)

*Bovine Viral Diarrhea Virus (BVDV)

*Duck Hepatitis B Virus

*Equine Influenza A Virus (H3N8) (EIV)

*Feline Calicivirus, Strain F9 (ATCC VR-782)

*Murine Norovirus

***Animal Viruses with a 1 minute contact time:**

*Canine Parvovirus, (ATCC VR-2017)

*Feline Infectious Peritonitis Virus, (FIPV) (ATCC VR-2022)

*Feline Herpes Virus, (Feline Rhinotracheitis)v (ATCC VR-636)

*Murine Hepatitis Virus, (ATCC VR-765)

*Minute Virus of Mouse, (Murine Parvovirus) (ATCC VR-1346)

TUBERCULOCIDAL against *Mycobacterium bovis* (Tb) (BCG) (OT 451C150) in the presence of a 5% serum load and 1 minute contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

FUNGICIDAL: This product is fungicidal against the pathogenic fungi *Trichophyton mentagrophytes* (ATCC 9533) the fungus which causes Athlete's Foot in the presence of a 5% soil and 30 second contact time when used as directed on hard surfaces found in bathrooms, shower stalls, locker rooms, or other clean, non-porous, hard surfaces commonly contacted by bare feet.

FUNGICIDAL against *Candida albicans* (ATCC 10231) and *Microsporium canis* (ATCC 10214) in the presence of 5% soil with a 1 minute contact time.

Mold/Mildew: Kills the growth of mold and mildew with a 4 minute contact time: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) (ATCC 16404) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Mold/Mildew: Controls and prevents the growth of mold and mildew: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Non-Food Contact Surface Sanitizer for hard non-porous surfaces in the presence of a 5% serum load and 10 second contact time on hard, non-porous environmental surfaces.

Acinetobacter baumannii, (ATCC 19606)

Campylobacter jejuni, (ATCC 29428)

Enterobacter aerogenes, (ATCC 13048)

Enterococcus faecalis, (Vancomycin Resistant (VRE)) (ATCC 51575)

Escherichia coli O157:H7, (ATCC 35150)

Klebsiella pneumoniae, (ATCC 4352)

Listeria monocytogenes, (ATCC 19111)

Pseudomonas aeruginosa, (ATCC 15442)

Salmonella enterica, (formally known as *Salmonella choleraesuis*) (ATCC 10708)

Staphylococcus aureus, (ATCC 6538)

Staphylococcus aureus, (Methicillin Resistant (MRSA)) (ATCC 33592)

Soft Surface Sanitizer in the presence of 5% soil with a 30 second contact time on cotton or polyester fabrics against the following bacteria:

Staphylococcus aureus, (ATCC 6538)

Enterobacter aerogenes, (ATCC 13048)

Enterococcus faecalis, (ATCC 51575) (Resistant to Vancomycin (VRE))

Staphylococcus aureus, (ATCC 33592) (Resistant to Methicillin (MRSA))

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Animal Housing Facilities (Including Veterinary Clinics, Farms, Livestock, Swine and Poultry Houses):

For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities (including barns, pens, stalls, chutes, and barns) occupied or traversed by animals.
3. Empty or cover all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water.
5. Saturate all hard, non-porous surfaces by spraying with product until thoroughly wet. To disinfect, all surfaces must remain wet for 1 minute.
6. Immerse or spray all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
8. Thoroughly scrub treated feed racks, troughs, mangers, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

To Kill Mold and Mildew: Pre-clean surfaces. Spray 6 – 8 inches from hard, nonporous surfaces until completely wet. Allow to remain wet for 4 minutes. Let air dry or wipe surface dry.

To Control Mold and Mildew: Spray 6 – 8 inches from hard, nonporous surfaces until completely wet. Let air dry or wipe surface dry. Repeat application weekly or when growth reappears.

See the container label for use directions and additional required information for use, storage and disposal.

EPA Reg. No. 70627-74