

Avert™ Sporicidal Disinfectant Cleaner Wipes

Sealed Air
Diversey Care

Diversey

Effective in 4 minutes against *C. difficile* spores

ACTIVE INGREDIENT:

Sodium hypochlorite..... 1.312%

OTHER INGREDIENTS..... 98.688%

TOTAL..... 100.000%

1.25% available Chlorine

**KEEP OUT OF
REACH
OF CHILDREN
DANGER**

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center for treatment advice.

IF SWALLOWED: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

**IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR DOCTOR FOR FURTHER TREATMENT ADVICE.
1-800-951-7145**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

See additional precautionary statements on side of label.

Net Contents: 6.2 lbs. (2.8 kg)

160 Wipes, 11 in. x 12 in. (28 cm x 30 cm)

100895932

Cleans • Deodorizes • Disinfects • *Virucidal • Fungicidal

Sporicidal against *Clostridium difficile* spores. Cleans, disinfects, and deodorizes in one labor saving step.

This product can be used in the following areas: assisted living facilities, hospitals, isolation rooms, and patient rooms.

This product has been developed to aid in the reduction of cross-contamination on treated surfaces not only in hospitals, but also in: bathrooms, health clubs, public buildings, and schools.

This product cleans, disinfects and deodorizes hard, nonporous hospital semi-critical medical surfaces in one step. It can be used on the following: patients beds, bedrails, bedframes, bedpans, bedside tables, coated covers for mattresses and pillows, diagnostics equipment, exterior surfaces of medical equipment, exercise equipment, floors, and walls.

May be used as part of a comprehensive approach to *Clostridium difficile* spore control. Using approved Standard Quantitative Disk Carrier Test Method under Good Laboratory Practices, in the presence of soil and 4 minute contact time, this product kills and inactivates spores of *Clostridium difficile* (ATCC 43598) on hard, nonporous surfaces.

Bacteria: *Pseudomonas aeruginosa*, *Staphylococcus aureus*

*Viruses: *Norovirus

See reference sheet for a complete list of claims and additional directions for use.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for personal cleansing. This is not a baby wipe. DO NOT FLUSH in toilet.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

Not recommended for use on or avoid contact with: aluminum, brass, chipped enamel, clear plastic, copper, finished floors, natural marble, painted surfaces, and silver.

Note: Prior to use on electronic equipment or components, unplug from electrical outlet, and ensure equipment is powered off.

Special Instructions for Cleaning Prior to Disinfection against *Clostridium difficile* spores

Personal Protection: Wear appropriate barrier protection such as gloves, gowns, masks or eye covering.

Cleaning Procedure: Fecal matter/waste must be thoroughly cleaned from surfaces/objects before disinfection by application with this product. Cleaning is to include vigorous wiping and/or scrubbing, until all visible soil is removed. Special attention is needed for high-touch surfaces. Surfaces in patient rooms are to be cleaned in an appropriate manner, such as from right to left or left to right, on horizontal surfaces, and top to bottom, on vertical surfaces, to minimize spreading of the spores. Restrooms are to be cleaned last. Do not reuse soiled cloths.

Infectious Materials Disposal: Materials used in the cleaning process that may contain feces/wastes are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

Disinfection Procedure: Wipe hard, nonporous, inanimate, environmental surfaces with towelette until thoroughly wet. Surfaces must remain wet for 4 minutes. Wipe surfaces and let air dry.

For Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Wipe hard, non-porous inanimate environmental surfaces with towelette.
3. All surfaces must remain visibly wet for 1 minute.
4. Wipe surfaces and let air dry.

***Kills HBV, HCV and HIV-1 on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids** in health care settings and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Hepatitis B Virus, Hepatitis C Virus and Human Immunodeficiency Virus Type 1 (HIV-1).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable latex gloves, gowns, masks and eye coverings.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of the disinfectant.

Contact Time: Allow surface to remain wet for 1 minute. For *Clostridium difficile* spores, allow to remain wet for 4 minutes.

Disposal of Infectious Materials: Blood and other body fluids, cleaning materials and clothing must be autoclaved and disposed of according to Federal, state and local regulations for infectious waste disposal.

STORAGE AND DISPOSAL

Do not contaminate food or feed by storage, disposal, or cleaning of equipment.

STORAGE: Store this product in a cool dry area, away from direct sunlight and heat to avoid deterioration. In case of spill, flood areas with large quantities of water. Keep container tightly closed between uses to retain moisture.

Container Disposal: Dispose of wipe in trash after use. DO NOT FLUSH. Nonrefillable container. Do not reuse or refill this container. Dispose of container in trash or offer for recycling if available.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Causes skin burns and eye damage. Do not get in eyes, on skin, or on clothing. Wear chemical-splash goggles, chemical-resistant gloves and protective footwear. In case of inadequate ventilation wear respiratory protection. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PHYSICAL OR CHEMICAL HAZARDS

Keep away from clothing and other combustible materials. Do not mix with any other product or chemical.

EPA Reg. No. 70627-75

EPA Est. No. 73027-WI-001

Lot code letters indicate establishment number.

Manufactured by:

©2015 Diversey, Inc.,

PO Box 19747, Charlotte, NC 28219-0747

Not for sale or use on or after: _____



STATE OF HAWAII
Department of Agriculture

ACCEPTED
LICENSE NO. **8017.172**

Avert™

Sporicidal Disinfectant Cleaner Wipes

Using approved Standard Quantitative Disk Carrier Test Method under Good Laboratory Practices, in the presence of soil and 4 minute contact time, this product kills and inactivates spores of *Clostridium difficile* (ATCC 43598) on hard, nonporous surfaces.

When used as directed, this product is highly effective against a broad-spectrum of pathogenic microorganisms including bacteria, antibiotic-resistant bacteria,*viruses, and fungi. Using approved AOAC test methods under Good Laboratory Practices, in the presence of 5% serum load and 1 minute contact time, this product kills the following on hard nonporous, inanimate, environmental surfaces:

Bacteria:

Pseudomonas aeruginosa, (ATCC 15442)
Staphylococcus aureus, (ATCC 6538)
Salmonella enterica, (ATCC 10708)

Campylobacter jejuni, (ATCC 29428)
Escherichia coli, (ATCC 11229)
Escherichia coli, coli O157:H7, (ATCC 35150)

Listeria monocytogenes, (ATCC 19117)
Stenotrophomonas maltophilia, (ATCC 25596)
Streptococcus pyogenes, (ATCC 19615)

Antibiotic-Resistant Bacteria:

Acinetobacter baumannii, Multi-Drug Resistant (MDRO) (ATCC 19606)
Enterobacter aerogenes, Multi-Drug Resistant (MDRO) (ATCC 29751)
Enterococcus faecalis, Vancomycin Resistant (VRE) (ATCC 51575)
Enterococcus faecium, Multi-Drug Resistant (ATCC 51559)
Escherichia coli, Carbapenem Resistant (CRE) (CDC 81371)

Escherichia coli, Extended-Spectrum Beta-lactamase (ESBL) producing (ATCC BAA-196)
Klebsiella pneumoniae, Carbapenem Resistant (CRKP) (ATCC BAA-1705)
Klebsiella pneumoniae, Extended-Spectrum Beta-lactamase (ESBL) producing (ATCC 700603)
Klebsiella pneumoniae, Multi Drug Resistant (ATCC 51503)

Klebsiella pneumoniae, NDM-1 positive (CDC 1000527)
Pseudomonas aeruginosa, Multi-Drug Resistant Metallo-beta lactamase (MBL) positive (CDC 2012059)
Staphylococcus aureus, Methicillin Resistant (MRSA) (ATCC 33592)
Staphylococcus aureus, Community Acquired Methicillin Resistant (CA-MRSA) Genotype USA 300, (NRS 384)

*Viruses:

*Adenovirus Type 2, (ATCC VR-846, Strain Adenoid 6)
*Adenovirus Type 4, (ATCC VR-4, Strain RI-67)
*Hepatitis A Virus, (Strain HM-175)
*Hepatitis B Virus (HBV)
*Hepatitis C Virus (HCV)

*Herpes Simplex Virus Type 2 (ATCC VR-734, Strain G)
*Human Coronavirus, (ATCC VR-740, Strain 229E)
*Human Immunodeficiency Virus (AIDS virus) (HTLV-IIIa)
*Influenza A Virus, 2009-H1N1 (Novel H1N1),
Strain A/Mexico/4108/2009 CDC# 2009712192

*Norovirus, Feline Calicivirus as a Surrogate for
*Poliovirus Type 1
*Respiratory Syncytial Virus (RSV) (ATCC VR-26, Strain Long)
*Rhinovirus Type 37, (ATCC VR-1147, Strain 151-1)
*Rotavirus, Strain WA

*Veterinary Viruses:

Avian Influenza A Virus, H5N1 (Strain VNHSN1-PR8/CDC-RG, CDC# 2006719965)

*Canine Parvovirus, (ATCC VR-2017, Strain Cornell-780916-80)

*Feline Calicivirus

*Animal Viruses with a 1 minute contact time:

*Canine Parvovirus, (ATCC VR-2017)
*Feline Infectious Peritonitis Virus, (FIPV) (ATCC VR-2022)

*Feline Herpes Virus, (Feline Rhinotracheitis)v (ATCC VR-636)
*Murine Hepatitis Virus, (ATCC VR-765)

*Minute Virus of Mouse, (Murine Parvovirus) (ATCC VR-1346)

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Animal Housing Facilities Including Veterinary Clinics, Farms, Livestock, Swine and Poultry Houses: For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc.

For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities (including pens, chutes, and barns) occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding or watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water. Wipe hard, non-porous inanimate environmental surfaces with towelette.
5. All surfaces must remain visibly wet for 1 minute.
6. Allow to air dry or rinse with potable water if necessary.
7. Ventilate buildings and other closed spaces.
8. Do not house animals or employ equipment until treatment has been absorbed, set or dried.

NOTE: Not for use on handling and restraining equipment such as leashes, muzzles, halters or ropes, as well as forks, shovels, and scrapers used for removing litter and manure.

For Use as a Cleaner and/or Deodorizer: Wipe hard, non-porous environmental surfaces. Allow to air dry or rinse with potable water if necessary.

See the container label for use directions and additional required information for First Aid, Precautionary Statements, and Storage and Disposal.

EPA Reg. No. 70627-75

Manufactured for: ©2015 Diversey, Inc., PO Box 19747, Charlotte, NC 28219-0747

Avert™ Sporicidal Disinfectant Cleaner Wipes

Effective in 4 minutes against *C. difficile* spores

ACTIVE INGREDIENT:

Sodium hypochlorite 1.312%

OTHER INGREDIENTS 98.688%

TOTAL 100.000%

1.25% available Chlorine

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center for treatment advice.

IF SWALLOWED: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR DOCTOR FOR FURTHER TREATMENT ADVICE.

1-800-851-7145

Have the product container or label with you when calling a poison control center or doctor,
or going for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

See additional precautionary statements on side of label.



STATE OF HAWAII
Department of Agriculture

ACCEPTED
LICENSE NO. **8017.172**

Sealed Air®
Diversey Care

Diversey®

**KEEP OUT OF REACH OF CHILDREN
DANGER**

**Net Contents: 6.2 lbs. (2.8 kg)
160 Wipes, 11 in. x 12 in. (28 cm x 30 cm)**

100895931

Cleans • Deodorizes • Disinfects • *Virucidal • Fungicidal

Sporicidal against *Clostridium difficile* spores

Cleans, disinfects, and deodorizes in one labor saving step.

This product can be used in the following areas: assisted living facilities, hospitals, isolation rooms, and patient rooms.

This product has been developed to aid in the reduction of cross-contamination on treated surfaces not only in hospitals, but also in: bathrooms, health clubs, public buildings, and schools.

This product cleans, disinfects and deodorizes hard, nonporous hospital semi-critical medical surfaces in one step. It can be used on the following: patients beds, bedrails, bedframes, bedpans, bedside tables, coated covers for mattresses and pillows, diagnostics equipment, exterior surfaces of medical equipment, exercise equipment, floors, and walls.

May be used as part of a comprehensive approach to *Clostridium difficile* spore control.

Using approved Standard Quantitative Disk Carrier Test Method under Good Laboratory Practices, in the presence of soil and 4 minute contact time, this product kills and inactivates spores of *Clostridium difficile* (ATCC 43598) on hard, nonporous surfaces.

Bacteria: *Pseudomonas aeruginosa*, *Staphylococcus aureus*

*Viruses: *Norovirus

See reference sheet for a complete list of claims and additional directions for use.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for personal cleansing. This is not a baby wipe. DO NOT FLUSH in toilet.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

Not recommended for use on or avoid contact with: aluminum, brass, chipped enamel, clear plastic, copper, finished floors, natural marble, painted surfaces, and silver.

Note: Prior to use on electronic equipment or components, unplug from electrical outlet, and ensure equipment is powered off.

Special Instructions for Cleaning Prior to Disinfection against *Clostridium difficile* spores

Personal Protection: Wear appropriate barrier protection such as gloves, gowns, masks or eye covering.

Cleaning Procedure: Fecal matter/waste must be thoroughly cleaned from surfaces/objects before disinfection by application with this product. Cleaning is to include vigorous wiping and/or scrubbing, until all visible soil is removed. Special attention is needed for high-touch surfaces. Surfaces in patient rooms are to be cleaned in an appropriate manner, such as from right to left or left to right, on horizontal surfaces, and top to bottom, on vertical surfaces, to minimize spreading of the spores. Restrooms are to be cleaned last. Do not reuse soiled cloths.

Infectious Materials Disposal: Materials used in the cleaning process that may contain feces/wastes are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

Disinfection Procedure: Wipe hard, nonporous, inanimate, environmental surfaces with towelette until thoroughly wet. Surfaces must remain wet for 4 minutes. Wipe surfaces and let air dry.

For Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Wipe hard, non-porous inanimate environmental surfaces with towelette.
3. All surfaces must remain visibly wet for 1 minute.
4. Wipe surfaces and let air dry.

Not for sale or use on or after: _____

***Kills HBV, HCV and HIV-1 on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids** in health care settings and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Hepatitis B Virus, Hepatitis C Virus and Human Immunodeficiency Virus Type 1 (HIV-1).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable latex gloves, gowns, masks and eye coverings.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of the disinfectant.

Contact Time: Allow surface to remain wet for 1 minute. For Clostridium difficile spores, allow to remain wet for 4 minutes.

Disposal of Infectious Materials: Blood and other body fluids, cleaning materials and clothing must be autoclaved and disposed of according to Federal, state and local regulations for infectious waste disposal.

STORAGE AND DISPOSAL

Do not contaminate food or feed by storage, disposal, or cleaning of equipment.

STORAGE: Store this product in a cool dry area, away from direct sunlight and heat to avoid deterioration. In case of spill, flood areas with large quantities of water. Keep container tightly closed between uses to retain moisture.

Container Disposal: Dispose of wipe in trash after use. DO NOT FLUSH. Refillable container. Refill this container only with this product. Do not reuse this container for any other purpose. Dispose of container in trash or offer for recycling if available after final use.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Causes skin burns and eye damage. Do not get in eyes, on skin, or on clothing. Wear chemical-splash goggles, chemical-resistant gloves and protective footwear. In case of inadequate ventilation wear respiratory protection. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PHYSICAL OR CHEMICAL HAZARDS

Keep away from clothing and other combustible materials. Do not mix with any other product or chemical.

EPA Reg. No. 70627-75

EPA Est. No. 73027-WI-001

Lot code letters indicate establishment number.

Manufactured for:

©2015 Diversey, Inc., PO Box 19747, Charlotte, NC 28219-0747



300151554 (15/254)

Avert™

Sporicidal Disinfectant Cleaner Wipes

Using approved Standard Quantitative Disk Carrier Test Method under Good Laboratory Practices, in the presence of soil and 4 minute contact time, this product kills and inactivates spores of *Clostridium difficile* (ATCC 43598) on hard, nonporous surfaces.

When used as directed, this product is highly effective against a broad-spectrum of pathogenic microorganisms including bacteria, antibiotic-resistant bacteria,*viruses, and fungi. Using approved AOAC test methods under Good Laboratory Practices, in the presence of 5% serum load and 1 minute contact time, this product kills the following on hard nonporous, inanimate, environmental surfaces:

Bacteria:

Pseudomonas aeruginosa, (ATCC 15442)
Staphylococcus aureus, (ATCC 6538)
Salmonella enterica, (ATCC 10708)

Campylobacter jejuni, (ATCC 29428)
Escherichia coli, (ATCC 11229)
Escherichia coli, coli O157:H7, (ATCC 35150)

Listeria monocytogenes, (ATCC 19117)
Stenotrophomonas maltophilia, (ATCC 25596)
Streptococcus pyogenes, (ATCC 19615)

Antibiotic-Resistant Bacteria:

Acinetobacter baumannii, Multi-Drug Resistant (MDRO) (ATCC 19606)
Enterobacter aerogenes, Multi-Drug Resistant (MDRO) (ATCC 29751)
Enterococcus faecalis, Vancomycin Resistant (VRE) (ATCC 51575)
Enterococcus faecium, Multi-Drug Resistant (ATCC 51559)
Escherichia coli, Carbapenem Resistant (CRE) (CDC 81371)

Escherichia coli, Extended-Spectrum Beta-lactamase (ESBL) producing (ATCC BAA-196)
Klebsiella pneumoniae, Carbapenem Resistant (CRKP) (ATCC BAA-1705)
Klebsiella pneumoniae, Extended-Spectrum Beta-lactamase (ESBL) producing (ATCC 700603)
Klebsiella pneumoniae, Multi Drug Resistant (ATCC 51503)

Klebsiella pneumoniae, NDM-1 positive (CDC 1000527)
Pseudomonas aeruginosa, Multi-Drug Resistant Metallo-beta lactamase (MBL) positive (CDC 2012059)
Staphylococcus aureus, Methicillin Resistant (MRSA) (ATCC 33592)
Staphylococcus aureus, Community Acquired Methicillin Resistant (CA-MRSA) Genotype USA 300, (NRS 384)

*Viruses:

*Adenovirus Type 2, (ATCC VR-846, Strain Adenoid 6)
*Adenovirus Type 4, (ATCC VR-4, Strain RI-67)
*Hepatitis A Virus, (Strain HM-175)
*Hepatitis B Virus (HBV)
*Hepatitis C Virus (HCV)

*Herpes Simplex Virus Type 2 (ATCC VR-734, Strain G)
*Human Coronavirus, (ATCC VR-740, Strain 229E)
*Human Immunodeficiency Virus (AIDS virus) (HTLV-IIIa)
*Influenza A Virus, 2009-H1N1 (Novel H1N1),
Strain A/Mexico/4108/2009 CDC# 2009712192

*Norovirus, Feline Calicivirus as a Surrogate for
*Poliovirus Type 1
*Respiratory Syncytial Virus (RSV) (ATCC VR-26, Strain Long)
*Rhinovirus Type 37, (ATCC VR-1147, Strain 151-1)
*Rotavirus, Strain WA

*Veterinary Viruses:

Avian Influenza A Virus, H5N1 (Strain VNHSN1-PR8/CDC-RG, CDC# 2006719965)

*Canine Parvovirus, (ATCC VR-2017, Strain Cornell-780916-80)

*Feline Calicivirus

*Animal Viruses with a 1 minute contact time:

*Canine Parvovirus, (ATCC VR-2017)
*Feline Infectious Peritonitis Virus, (FIPV) (ATCC VR-2022)

*Feline Herpes Virus, (Feline Rhinotracheitis)v (ATCC VR-636)
*Murine Hepatitis Virus, (ATCC VR-765)

*Minute Virus of Mouse, (Murine Parvovirus) (ATCC VR-1346)

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Animal Housing Facilities Including Veterinary Clinics, Farms, Livestock, Swine and Poultry Houses: For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc.

For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities (including pens, chutes, and barns) occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding or watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water. Wipe hard, non-porous inanimate environmental surfaces with towelette.
5. All surfaces must remain visibly wet for 1 minute.
6. Allow to air dry or rinse with potable water if necessary.
7. Ventilate buildings and other closed spaces.
8. Do not house animals or employ equipment until treatment has been absorbed, set or dried.

NOTE: Not for use on handling and restraining equipment such as leashes, muzzles, halters or ropes, as well as forks, shovels, and scrapers used for removing litter and manure.

For Use as a Cleaner and/or Deodorizer: Wipe hard, non-porous environmental surfaces. Allow to air dry or rinse with potable water if necessary.

See the container label for use directions and additional required information for First Aid, Precautionary Statements, and Storage and Disposal.

EPA Reg. No. 70627-75

Manufactured for: ©2015 Diversey, Inc., PO Box 19747, Charlotte, NC 28219-0747

STORAGE AND DISPOSAL

Do not contaminate food or feed by storage, disposal, or cleaning of equipment.

STORAGE: Store this product in a cool dry area, away from direct sunlight and heat to avoid deterioration. In case of spill, flood areas with large quantities of water. Keep container tightly closed between uses to retain moisture.

Container Disposal: Dispose of wipe in trash after use. DO NOT FLUSH. Nonrefillable container. Do not reuse or refill this container. Dispose of container in trash or offer for recycling if available.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Causes skin burns and eye damage. Do not get in eyes, on skin, or on clothing. Wear chemical-splash goggles, chemical-resistant gloves and protective footwear. In case of inadequate ventilation wear respiratory protection. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PHYSICAL OR CHEMICAL HAZARDS

Keep away from clothing and other combustible materials. Do not mix with any other product or chemical.

EPA Reg. No. 70627-75

EPA Est. No. 73027-WI-001

Lot code letters indicate establishment number.

©2015 Manufactured for Diversey, Inc.,
PO Box 19747, Charlotte, NC 28219-0747



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO. 8017.172

Avert™ Sporicidal Disinfectant Cleaner Wipes

Sealed Air
Diversey Care

Diversey®

Effective in 4 minutes against *Clostridium difficile* spores.

ACTIVE INGREDIENT:

Sodium hypochlorite 1.312%

OTHER INGREDIENTS 98.688%

TOTAL 100.000%

1.25% available Chlorine

KEEP OUT OF REACH OF CHILDREN DANGER

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center for treatment advice.

IF SWALLOWED: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR DOCTOR FOR FURTHER

TREATMENT ADVICE.

1-800-851-7145

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

See additional precautionary statements on side of label.

Net contents: 1.9 lb. (879 g),
160 Wipes, 6.1 in. x 6.8 in.
(15.5 cm x 17.3 cm)

100895790

Cleans • Deodorizes • Disinfects • *Virucidal • Fungicidal

Sporicidal against *Clostridium difficile* spores
Cleans, disinfects, and deodorizes in one labor saving step.

This product can be used in the following areas: assisted living facilities, hospitals, isolation rooms, and patient rooms.

This product has been developed to aid in the reduction of cross-contamination on treated surfaces not only in hospitals, but also in: bathrooms, health clubs, public buildings, and schools.

This product cleans, disinfects and deodorizes hard, nonporous hospital semi-critical medical surfaces in one step. It can be used on the following: patients beds, bedrails, bedframes, bedpans, bedside tables, coated covers for mattresses and pillows, diagnostics equipment, exterior surfaces of medical equipment, exercise equipment, floors, and walls.

May be used as part of a comprehensive approach to *Clostridium difficile* spore control.

Using approved Standard Quantitative Disk Carrier Test Method under Good Laboratory Practices, in the presence of soil and 4 minute contact time, this product kills and inactivates spores of *Clostridium difficile* (ATCC 43598) on hard, nonporous surfaces.

Bacteria: *Pseudomonas aeruginosa*, *Staphylococcus aureus*

*Viruses: *Norovirus

See reference sheet for a complete list of claims and additional directions for use.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for personal cleansing. This is not a baby wipe. DO NOT FLUSH in toilet.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

Not recommended for use on or avoid contact with: aluminum, brass, chipped enamel, clear plastic, copper, finished floors, natural marble, painted surfaces, and silver.

Note: Prior to use on electronic equipment or components, unplug from electrical outlet, and ensure equipment is powered off.

Special Instructions for Cleaning Prior to Disinfection against *Clostridium difficile* spores

Personal Protection: Wear appropriate barrier protection such as gloves, gowns, masks or eye covering.

Cleaning Procedure: Fecal matter/waste must be thoroughly cleaned from surfaces/objects before disinfection by application with this product. Cleaning is to include vigorous wiping and/or scrubbing, until all visible soil is removed. Special attention is needed for high-touch surfaces. Surfaces in patient rooms are to be cleaned in an appropriate manner, such as from right to left or left to right, on horizontal surfaces, and top to bottom, on vertical surfaces, to minimize spreading of the spores. Restrooms are to be cleaned last. Do not reuse soiled cloths.

Infectious Materials Disposal: Materials used in the cleaning process that may contain feces/wastes are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

Disinfection Procedure: Wipe hard, nonporous, inanimate, environmental surfaces with towelette until thoroughly wet. Surfaces must remain wet for 4 minutes. Wipe surfaces and let air dry.

For Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Wipe hard, non-porous inanimate environmental surfaces with towelette.
3. All surfaces must remain visibly wet for 1 minute.
4. Wipe surfaces and let air dry.

Not for sale or use on or after: _____



***Kills HBV, HCV and HIV-1 on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids** in health care settings and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Hepatitis B Virus, Hepatitis C Virus and Human Immunodeficiency Virus Type 1 (HIV-1).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable latex gloves, gowns, masks and eye coverings.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of the disinfectant.

Contact Time: Allow surface to remain wet for 1 minute. For *Clostridium difficile* spores, allow to remain wet for 4 minutes.

Disposal of Infectious Materials: Blood and other body fluids, cleaning materials and clothing must be autoclaved and disposed of according to Federal, state and local regulations for infectious waste disposal.



300151550 (15/254)

Avert™

Sporicidal Disinfectant Cleaner Wipes

Using approved Standard Quantitative Disk Carrier Test Method under Good Laboratory Practices, in the presence of soil and 4 minute contact time, this product kills and inactivates spores of *Clostridium difficile* (ATCC 43598) on hard, nonporous surfaces.

When used as directed, this product is highly effective against a broad-spectrum of pathogenic microorganisms including bacteria, antibiotic-resistant bacteria,*viruses, and fungi. Using approved AOAC test methods under Good Laboratory Practices, in the presence of 5% serum load and 1 minute contact time, this product kills the following on hard nonporous, inanimate, environmental surfaces:

Bacteria:

Pseudomonas aeruginosa, (ATCC 15442)
Staphylococcus aureus, (ATCC 6538)
Salmonella enterica, (ATCC 10708)

Campylobacter jejuni, (ATCC 29428)
Escherichia coli, (ATCC 11229)
Escherichia coli, coli O157:H7, (ATCC 35150)

Listeria monocytogenes, (ATCC 19117)
Stenotrophomonas maltophilia, (ATCC 25596)
Streptococcus pyogenes, (ATCC 19615)

Antibiotic-Resistant Bacteria:

Acinetobacter baumannii, Multi-Drug Resistant (MDRO) (ATCC 19606)
Enterobacter aerogenes, Multi-Drug Resistant (MDRO) (ATCC 29751)
Enterococcus faecalis, Vancomycin Resistant (VRE) (ATCC 51575)
Enterococcus faecium, Multi-Drug Resistant (ATCC 51559)
Escherichia coli, Carbapenem Resistant (CRE) (CDC 81371)

Escherichia coli, Extended-Spectrum Beta-lactamase (ESBL) producing (ATCC BAA-196)
Klebsiella pneumoniae, Carbapenem Resistant (CRKP) (ATCC BAA-1705)
Klebsiella pneumoniae, Extended-Spectrum Beta-lactamase (ESBL) producing (ATCC 700603)
Klebsiella pneumoniae, Multi Drug Resistant (ATCC 51503)

Klebsiella pneumoniae, NDM-1 positive (CDC 1000527)
Pseudomonas aeruginosa, Multi-Drug Resistant Metallo-beta lactamase (MBL) positive (CDC 2012059)
Staphylococcus aureus, Methicillin Resistant (MRSA) (ATCC 33592)
Staphylococcus aureus, Community Acquired Methicillin Resistant (CA-MRSA) Genotype USA 300, (NRS 384)

*Viruses:

*Adenovirus Type 2, (ATCC VR-846, Strain Adenoid 6)
*Adenovirus Type 4, (ATCC VR-4, Strain RI-67)
*Hepatitis A Virus, (Strain HM-175)
*Hepatitis B Virus (HBV)
*Hepatitis C Virus (HCV)

*Herpes Simplex Virus Type 2 (ATCC VR-734, Strain G)
*Human Coronavirus, (ATCC VR-740, Strain 229E)
*Human Immunodeficiency Virus (AIDS virus) (HTLV-IIIa)
*Influenza A Virus, 2009-H1N1 (Novel H1N1),
Strain A/Mexico/4108/2009 CDC# 2009712192

*Norovirus, Feline Calicivirus as a Surrogate for
*Poliovirus Type 1
*Respiratory Syncytial Virus (RSV) (ATCC VR-26, Strain Long)
*Rhinovirus Type 37, (ATCC VR-1147, Strain 151-1)
*Rotavirus, Strain WA

*Veterinary Viruses:

Avian Influenza A Virus, H5N1 (Strain VNHSN1-PR8/CDC-RG, CDC# 2006719965)

*Canine Parvovirus, (ATCC VR-2017, Strain Cornell-780916-80)

*Feline Calicivirus

*Animal Viruses with a 1 minute contact time:

*Canine Parvovirus, (ATCC VR-2017)
*Feline Infectious Peritonitis Virus, (FIPV) (ATCC VR-2022)

*Feline Herpes Virus, (Feline Rhinotracheitis)v (ATCC VR-636)
*Murine Hepatitis Virus, (ATCC VR-765)

*Minute Virus of Mouse, (Murine Parvovirus) (ATCC VR-1346)

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Animal Housing Facilities Including Veterinary Clinics, Farms, Livestock, Swine and Poultry Houses: For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc.

For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities (including pens, chutes, and barns) occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding or watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water. Wipe hard, non-porous inanimate environmental surfaces with towelette.
5. All surfaces must remain visibly wet for 1 minute.
6. Allow to air dry or rinse with potable water if necessary.
7. Ventilate buildings and other closed spaces.
8. Do not house animals or employ equipment until treatment has been absorbed, set or dried.

NOTE: Not for use on handling and restraining equipment such as leashes, muzzles, halters or ropes, as well as forks, shovels, and scrapers used for removing litter and manure.

For Use as a Cleaner and/or Deodorizer: Wipe hard, non-porous environmental surfaces. Allow to air dry or rinse with potable water if necessary.

See the container label for use directions and additional required information for First Aid, Precautionary Statements, and Storage and Disposal.

EPA Reg. No. 70627-75

Manufactured for: ©2015 Diversey, Inc., PO Box 19747, Charlotte, NC 28219-0747