

Sealed Air
Diversey Care

Diversey

Oxivir® 1 Wipes

***Virucide • Bactericide • Tuberculocide • Fungicide
For Institutional Use**

Active Ingredient:

Hydrogen Peroxide..... 0.5%

Other Ingredients:99.5%

Total: 100.0%



KEEP OUT OF REACH OF CHILDREN

BACTERICIDAL: 1 MINUTE

VIRUCIDAL: 30 SECONDS

ANIMAL PREMISE *VIRUCIDAL: 1 MINUTE

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER: 10 SECONDS

FUNGICIDAL: 30 SECONDS



AHP Technology and Logos are registered trademarks
of Virox Technologies Inc., used under license.

Net Contents: 60 Wipes

7 in. x 8 in. (17.8 cm x 20.3 cm)

1.11 lb. (0.5 kg)

100850922

This product has been developed to aid in the reduction of cross-contamination between treated surfaces in: airports, athletic facilities, correctional facilities, cruise ships, day care centers, health clubs. This product can be used in: clinics, hospitals, nursing homes, operating rooms, patient rooms, rehabilitation centers, office buildings, pet shops, public restrooms, retail businesses, schools, and veterinary clinics.

For daily cleaning and disinfecting of hard, nonporous inanimate surface materials: acrylic materials: Corian®, chrome, glass, glazed tile, laminated surfaces, linoleum, metal (except brass or copper), painted surfaces, and plastic.

Note: This product is compatible with the listed materials. If product is intended to be used on other surfaces, it is recommended to test product by applying to a small discrete area to determine compatibility before proceeding with its use.

This product cleans, disinfects and deodorizes hard, nonporous surfaces in one step. May be used to clean and disinfect sinks, faucets, fixtures, bathrooms, restrooms, tubs, and showers.

It can be used on the following hard, nonporous surfaces: chairs, counters, faucets, sinks, urinals, exercise equipment, floors, lockers, showers, tubs, and walls. Do not use on glasses, dishes and utensils.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not recommended for use on: brass or copper.

When used as directed, this product is highly effective against a broad-spectrum of pathogenic microorganisms. Using approved AOAC test methods, in the presence of 5% serum load and 1 minute contact time, unless otherwise noted, this product kills the following on hard nonporous environmental surfaces: **Bacteria:** *Pseudomonas aeruginosa*, *Staphylococcus aureus*, **Antibiotic-Resistant Bacteria:** *Staphylococcus aureus* Methicillin Resistant (MRSA), *Staphylococcus aureus* Vancomycin Intermediate Resistant (VISA), *Staphylococcus aureus* Vancomycin Resistant (VRSA). ***Virucidal**, in the presence of 5% serum load with a 30 second contact time, unless otherwise noted. ***Viruses:** *Rhinovirus Type 37, *Norovirus (Feline Calicivirus, as the surrogate) ***Animal Virus:** *Murine Norovirus.

See reference sheet for a complete list of pathogenic organisms

Not for personal cleansing. This is not a baby wipe. Do not flush in toilet.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

To Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Wipe hard, non-porous environmental surfaces with towelette.
3. Allow surface to remain wet for 1 minute. For certain listed viruses, allow to remain wet for 30 seconds.
4. Allow to air dry or rinse with potable water if necessary.

Note: All food contact surfaces must be rinsed with potable water. Do not use on glassware, utensils, or dishes.

To Use as a Restroom, Shower Room, Locker Room, Athletic Equipment Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Wipe hard, non-porous environmental surfaces with towelette.
3. Allow surfaces to remain wet for 1 minute. For certain listed viruses, allow to remain wet for 30 seconds.
4. Rinse surfaces thoroughly and let air dry.

Note: Floors will be slippery when wet or contaminated with foreign materials.

***KILLS HIV, HCV and HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (Hospitals, Nursing Homes) and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis C Virus (HCV) or Hepatitis B Virus (HBV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS), HCV OR HBV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood/body fluids must be thoroughly cleaned from surfaces/objects before application of this product.

Contact Time: Allow surface to remain wet for 30 seconds to kill HIV-1, HCV and HBV, 4 minutes for mold, and 1 minute to kill other organisms cited on label.

Disposal of Infectious Material: Blood/body fluids must be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

Animal Housing Facilities Including Veterinary Clinics, Equine Farms, Swine and Poultry Houses:

For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities (including barns, pens, stalls, chutes, and barns) occupied or traversed by animals.
3. Empty or cover all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water.
5. Wipe hard, non-porous environmental surfaces with towelettes. To disinfect, all surfaces must remain wet for 1 minute. For certain listed viruses, allow to remain wet for 30 seconds.
6. Allow to air dry or rinse with potable water if necessary.
7. Ventilate buildings and other closed spaces.
8. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
9. Thoroughly scrub treated feed racks, troughs, mangers, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

To Kill Fungi: Pre-clean heavily soiled areas. Wipe hard, non-porous environmental surfaces with towelette. Allow to remain wet for 1 minute. Use additional towelettes if needed to ensure the correct wet contact time. Wipe surfaces or let air dry.

To Kill Mold and Mildew: Pre-clean surfaces. Wipe hard, nonporous surfaces with towelettes until completely wet. Allow to remain wet for 4 minutes. Let air dry.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Store in original container in areas inaccessible to small children in a cool, dry area away from direct sunlight and heat. Keep securely closed.

Towelette Storage/Disposal: Dispose of wipe in trash after use. Do not flush. Tightly close lid between uses to retain moisture.

Container Disposal: Non-refillable container. Do not reuse or refill this container. Offer empty container for recycling. If recycling is not available, discard container in trash.

Questions? Comments
1-800-558-2332

See SDS # MS0800916

EPA Reg. No. 70627-77

EPA Est. No. 74559-CAN-1 [VX]

Lot code letters indicate establishment number.

EMERGENCY PHONE : 1-800-851-7145



Manufactured for:
©2016 Diversey, Inc.,
PO Box 19747,
Charlotte, NC 28219-0747



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO.

8017.176

300153653 (16/126)

Oxivir® 1 Wipes

When used as directed, this product is highly effective against a wide-variety (broad-spectrum) of pathogenic microorganisms including bacteria, antibiotic-resistant bacteria, *viruses, and fungi. Using approved AOAC test methods, in the presence of 5% serum load and 1 minute contact time, unless otherwise noted, this product kills the following on hard nonporous inanimate surfaces:

Bacteria:

Pseudomonas aeruginosa, (ATCC 15442)
Staphylococcus aureus, (ATCC 6538)
Salmonella enterica, (ATCC 10708)
 formerly known as *Salmonella choleraesuis*
Bordetella bronchiseptica, (ATCC 10580)
Burkholderia cepacia, ((ATCC 25416)
Bordetella pertussis, (ATCC 12743)

Enterobacter aerogenes, (ATCC 13048)
Enterococcus faecium, (ATCC 51559)
Escherichia coli, (ATCC 11229)
Escherichia coli O157:H7, (ATCC 35150)
Klebsiella pneumoniae, (ATCC 4352)
Legionella pneumophila, (ATCC 33153)

Pasteurella multocida, (ATCC 12947)
Serratia marcescens, (ATCC 14756)
Staphylococcus pseudintermedius, (ATCC 49444)
Shigella dysenteriae, (ATCC 11835)
Stenotrophomonas (Pseudomonas) maltophilia, (ATCC 25596)
Streptococcus pyogenes, (ATCC19615)

Antibiotic-Resistant Bacteria:

Acinetobacter baumannii,

Multi-Drug Resistant (MDR) (ATCC 19606)
 Ampicillin, Cefazolin, Cefepime, Gentamicin,
 Ertapenem, and Trimethoprim/Sulfa
Enterobacter cloacae, (New Delhi metallo-β-lactamase
 1 positive (producing) (NDM 1)) (CDC 1000654)
Enterococcus faecalis, (ATCC 51575) (Vancomycin Resistant (VRE))
Escherichia coli, Extended Spectrum Beta-lactamase
 producing (positive) (ESBL) (ATCC BAA-196)
Escherichia coli, (New Delhi metallo-β-lactamase 1 (NDM-1) producing
 (positive)) (CDC 1001728)

Klebsiella pneumoniae, (Carbapenem Resistant (CRE)) (ATCC BAA 1705)
Klebsiella pneumoniae, (Extended Spectrum beta-lactamase (ESBL) producing
 (positive)) (ATCC 700603)
Klebsiella pneumoniae, (New Delhi metallo-β-lactamase 1 (NDM-1)
 producing (positive)) (CDC 1000527)
Pseudomonas aeruginosa, (Metallo-beta lactamase (MBL)
 positive (CDC 2012059))
Staphylococcus aureus, Community Associated Methicillin Resistant (CA-MRSA)
 (NARSA NRS 123) (Genotype USA400)
Staphylococcus aureus, (Methicillin Resistant (MRSA)) (ATCC 33591)

Staphylococcus aureus, (ATCC 33592) (Methicillin Resistant (MRSA))
Staphylococcus aureus, (Vancomycin Intermediate Resistant (VISA)) [HIP 5836]
Staphylococcus aureus, (Vancomycin Resistant (VRSA))[VRS1]
Staphylococcus epidermidis, (Methicillin Resistant (MRSE)) (ATCC 51625)
Streptococcus pneumoniae, (Penicillin Resistant (PRSP)) (ATCC 700677)

***Virucidal** in the presence of 5% serum load with a 30 second contact time, unless otherwise noted.

*Viruses:

*Adenovirus Type 8, (ATCC VR-1368)
 *Cytomegalovirus, (ATCC VR-538)
 *Hepatitis B Virus (HBV), Duck Hepatitis B Virus as surrogate
 (Acquired from Hepadnavirus Testing)
 *Hepatitis C Virus (HCV), Bovine Viral Diarrhea
 Virus as surrogate (Acquired from National Veterinary
 Services Laboratories)

*Herpes Simplex Virus, Type 1 (HSV-1) (ATCC VR-733)
 *Herpes Simplex Virus, Type 2 (HSV-2), (ATCC VR-734)
 *HIV-1 (AIDS Virus), Strain HTLV-III_B (HIV-1),
 (Acquired from Adv. Biotechnologies, Inc.)
 *Human Coronavirus, (ATCC VR-740)
 *Influenza A Virus (Hong Kong Strain), (ATCC VR-544)
 *Influenza B Virus, (ATCC VR-823)

*Norovirus, (Feline Calicivirus, ATCC VR-782 as the surrogate)
 *Pandemic Influenza A Virus H1N1
 *Parainfluenza Virus Type 3, (ATCC VR-93)
 *Respiratory Syncytial Virus, (RSV) (ATCC VR-26)
 *Rhinovirus Type 37, Strain 151-1, (ATCC VR-1147)
 *Rotavirus, (WA strain) (Acquired from University of Ottawa)
 *Vaccinia Virus, (ATCC VR-119)

*Animal Viruses:

*Avian Influenza A Virus (H7N9)
 *Avian Influenza A Virus (H3N2), (ATCC VR-2072)

*Bovine Viral Diarrhea Virus (BVDV)
 *Duck Hepatitis B Virus
 *Equine Influenza A Virus (H3N8) (EIV)

*Feline Calicivirus, Strain F9 (ATCC VR-782)
 *Murine Norovirus

*Animal Viruses with a 1 minute contact time:

*Feline Infectious Peritonitis Virus, (FIPV) (ATCC VR-2022)
 *Murine Hepatitis Virus, (ATCC VR-765)

*Feline Herpes Virus, (Feline Rhinotracheitis) (ATCC VR-636)

*Minute Virus of Mouse, (Murine Parvovirus) (ATCC VR-1346)

TUBERCULOCIDAL against *Mycobacterium bovis* (Tb) (BCG) (OT 451C150) in the presence of a 5% serum load and 1 minute contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

FUNGICIDAL: This product is fungicidal against the pathogenic fungi *Trichophyton mentagrophytes* (ATCC 9533) (the fungus which causes Athlete's Foot) in the presence of a 5% soil and 30 second contact time when used as directed on hard surfaces found in bathrooms, shower stalls, locker rooms, or other clean, non-porous, hard surfaces commonly contacted by bare feet

FUNGICIDAL against *Candida albicans* (ATCC 10231) and *Microsporum canis* (ATCC 10214) in the presence of 5% soil with a 1 minute contact time.

Mold/Mildew (Mildewcidal Activity): Kills the growth of mold and mildew with a 4 minute contact time: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) (ATCC 16404) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Mold/Mildew (Mildewstatic Activity): Controls and prevents the growth of mold and mildew: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Non-Food Contact Surface Sanitizer for hard non-porous surfaces in the presence of a 5% serum load and 10 second contact time on hard, non-porous environmental surfaces.

Acinetobacter baumannii, (ATCC 19606)
Campylobacter jejuni, (ATCC 29428)
Enterobacter aerogenes, (ATCC 13048)
Enterococcus faecalis, (Vancomycin Resistant (VRE)) (ATCC 51575)

Escherichia coli O157:H7, (ATCC 35150)
Klebsiella pneumoniae, (ATCC 4352)
Listeria monocytogenes, (ATCC 19111)
Pseudomonas aeruginosa, (ATCC 15442)

Salmonella enterica, (formally known as *Salmonella choleraesuis*) (ATCC 10708)
Staphylococcus aureus, (ATCC 6538)
Staphylococcus aureus, (Methicillin Resistant (MRSA)) (ATCC 33592)

Malodor Counteractancy: eliminates odors and odor-causing bacteria on hard, nonporous surfaces in restroom areas, behind and under sinks and counters, and storage areas and other places where bacterial growth can cause malodors.

• Kills Pandemic 2009 H1N1 Influenza A virus.

See the container label for use directions and additional required information for use, storage and disposal.

EPA Reg. No. 70627-77

Manufactured for: ©2016 Diversey, Inc., PO Box 19747, Charlotte, NC 28219-0747

Sealed Air
Diversey Care

Diversey

Oxivir® 1 Wipes

*Virucide • Bactericide • Tuberculocide • Fungicide
For Institutional Use

Active Ingredient:

Hydrogen Peroxide..... 0.5%

Other Ingredients: 99.5%

Total: 100.0%



**KEEP OUT OF REACH
OF CHILDREN**

BACTERICIDAL: 1 MINUTE

VIRUCIDAL: 30 SECONDS

ANIMAL PREMISE *VIRUCIDAL: 1 MINUTE

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER: 10 SECONDS

FUNGICIDAL: 30 SECONDS



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO.

8017.176



AHP Technology and Logos are registered
trademarks of Virox Technologies Inc., used
under license.

Net Contents: **160** Wipes
6.1 in. x 6.8 in. (15.5 cm x 17.3 cm)
1.97 lb. (0.89 kg)

100850923

This product has been developed to aid in the reduction of cross-contamination between treated surfaces in: airports, athletic facilities, correctional facilities, cruise ships, day care centers, health clubs. This product can be used in: clinics, hospitals, nursing homes, operating rooms, patient rooms, rehabilitation centers, office buildings, pet shops, public restrooms, retail businesses, schools, and veterinary clinics. For daily cleaning and disinfecting of hard, nonporous inanimate surface materials: acrylic, Corian®, chrome, glass, glazed tile, laminated surfaces, linoleum, metal (except for brass or copper), painted surfaces, and plastic.

Note: This product is compatible with the listed materials. If product is intended to be used on other surfaces, it is recommended to test product by applying to a small discrete area to determine compatibility before proceeding with its use.

This product cleans, disinfects and deodorizes hard, nonporous surfaces in one step. May be used to clean and disinfect sinks, faucets, fixtures, bathrooms, restrooms, tubs, and showers.

It can be used on the following hard, nonporous surfaces: chairs, counters, faucets, sinks, urinals, exercise equipment, floors, lockers, showers, tubs, and walls. Do not use on glasses, dishes and utensils.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Not recommended for use on: brass or copper.

When used as directed, this product is highly effective against a wide-variety (broad-spectrum) of pathogenic microorganisms. Using approved AOAC test methods, in the presence of 5% serum load and 1 minute contact time, unless otherwise noted, this product kills the following on hard nonporous environmental surfaces: **Bacteria:** *Pseudomonas aeruginosa*, *Staphylococcus aureus*, **Antibiotic-Resistant Bacteria:** *Staphylococcus aureus* Methicillin Resistant (MRSA), *Staphylococcus aureus* Vancomycin Intermediate Resistant (VISA), *Staphylococcus aureus* Vancomycin Resistant (VRSA). ***Virucidal**, in the presence of 5% serum load with a 30 second contact time, unless otherwise noted. ***Viruses:** *Rhinovirus Type 37, *Norovirus (Feline Calicivirus, as the surrogate) ***Animal Virus:** *Murine Norovirus. See reference sheet for a complete list of pathogenic organisms.

Not for personal cleansing. This is not a baby wipe. Do not flush in toilet.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

To Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Wipe hard, non-porous environmental surfaces with towelette.
3. Allow surface to remain wet for 1 minute. For certain listed viruses, allow to remain wet for 30 seconds.
4. Allow to air dry or rinse with potable water if necessary.

Note: All food contact surfaces must be rinsed with potable water. Do not use on glassware, utensils, or dishes.

To Use as a Restroom, Shower Room, Locker Room, Athletic Equipment Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Wipe hard, non-porous environmental surfaces with towelette.
3. Allow surfaces to remain wet for 1 minute. For certain listed viruses, allow to remain wet for 30 seconds.
4. Rinse surfaces thoroughly and let air dry.

Note: Floors will be slippery when wet or contaminated with foreign materials.

See SDS # MS0800916

EPA Reg. No. 70627-77

EPA Est. No. 74559-CAN-1 [VX]

Lot code letters indicate establishment number.

Manufactured for:

©2016 Diversey, Inc., PO Box 19747, Charlotte, NC 28219-0747



***KILLS HIV, HCV and HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (Hospitals, Nursing Homes) and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis C Virus (HCV) or Hepatitis B Virus (HBV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS), HCV OR HBV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood/body fluids must be thoroughly cleaned from surfaces/objects before application of this product.

Contact Time: Allow surface to remain wet for 30 seconds to kill HIV-1, HCV and HBV, 4 minutes for mold, and 1 minute to kill other organisms cited on label.

Disposal of Infectious Material: Blood/body fluids must be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.



Questions? Comments
1-800-558-2332

EMERGENCY PHONE : 1-800-851-7145

Animal Housing Facilities Including Veterinary Clinics, Equine Farms, Swine and Poultry Houses:

For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities (including barns, pens, stalls, chutes, and barns) occupied or traversed by animals.
3. Empty or cover all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water.
5. Wipe hard, non-porous environmental surfaces with towelettes. To disinfect, all surfaces must remain wet for 1 minute. For certain listed viruses, allow to remain wet for 30 seconds.
6. Allow to air dry or rinse with potable water if necessary.
7. Ventilate buildings and other closed spaces.
8. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
9. Thoroughly scrub treated feed racks, troughs, mangers, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

To Kill Fungi: Pre-clean heavily soiled areas. Wipe hard, non-porous environmental surfaces with towelette. Allow to remain wet for 1 minute. Use additional towelettes if needed to ensure the correct wet contact time. Wipe surfaces or let air dry.

To Kill Mold and Mildew: Pre-clean surfaces. Wipe hard, nonporous surfaces with towelettes until completely wet. Allow to remain wet for 4 minutes. Let air dry.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Store in original container in areas inaccessible to small children in a cool, dry area away from direct sunlight and heat. Keep securely closed.

Towelette Storage/Disposal: Dispose of wipe in trash after use. Do not flush. Tightly close lid between uses to retain moisture.

Container Disposal: Non-refillable container. Do not reuse or refill this container. Offer empty container for recycling. If recycling is not available, discard container in trash.

(L)

EXP.

Oxivir® 1 Wipes

When used as directed, this product is highly effective against a wide-variety (broad-spectrum) of pathogenic microorganisms including bacteria, antibiotic-resistant bacteria, *viruses, and fungi. Using approved AOAC test methods, in the presence of 5% serum load and 1 minute contact time, unless otherwise noted, this product kills the following on hard nonporous inanimate surfaces:

Bacteria:

Pseudomonas aeruginosa, (ATCC 15442)
Staphylococcus aureus, (ATCC 6538)
Salmonella enterica, (ATCC 10708)
 formerly known as *Salmonella choleraesuis*
Bordetella bronchiseptica, (ATCC 10580)
Burkholderia cepacia, (ATCC 25416)
Bordetella pertussis, (ATCC 12743)

Enterobacter aerogenes, (ATCC 13048)
Enterococcus faecium, (ATCC 51559)
Escherichia coli, (ATCC 11229)
Escherichia coli O157:H7, (ATCC 35150)
Klebsiella pneumoniae, (ATCC 4352)
Legionella pneumophila, (ATCC 33153)

Pasteurella multocida, (ATCC 12947)
Serratia marcescens, (ATCC 14756)
Staphylococcus pseudintermedius, (ATCC 49444)
Shigella dysenteriae, (ATCC 11835)
Stenotrophomonas (Pseudomonas) maltophilia, (ATCC 25596)
Streptococcus pyogenes, (ATCC19615)

Antibiotic-Resistant Bacteria:

Acinetobacter baumannii,

Multi-Drug Resistant (MDR) (ATCC 19606)
 Ampicillin, Cefazolin, Cefepime, Gentamicin,
 Ertapenem, and Trimethoprim/Sulfa
Enterobacter cloacae, (New Delhi metallo-β-lactamase
 1 positive (producing) (NDM 1)) (CDC 1000654)

Enterococcus faecalis, (ATCC 51575) (Vancomycin Resistant (VRE))
Escherichia coli, Extended Spectrum Beta-lactamase
 producing (positive (ESBL) (ATCC BAA-196)
Escherichia coli, (New Delhi metallo-β-lactamase 1(NDM-1) producing
 (positive)) (CDC 1001728)

Klebsiella pneumoniae, (Carbapenem Resistant (CRE)) (ATCC BAA 1705)
Klebsiella pneumoniae, (Extended Spectrum beta-lactamase (ESBL) producing
 (positive)) (ATCC 700603)

Klebsiella pneumoniae, (New Delhi metallo-β-lactamase 1(NDM-1)
 producing (positive)) (CDC 1000527)

Pseudomonas aeruginosa, (Metallo-β lactamase (MBL)
 positive (CDC 2012059))

Staphylococcus aureus, Community Associated Methicillin Resistant (CA-MRSA)
 (NARSA NRS 123) (Genotype USA400)
Staphylococcus aureus, (Methicillin Resistant (MRSA)) (ATCC 33591)

Staphylococcus aureus, (ATCC 33592) (Methicillin Resistant (MRSA))
Staphylococcus aureus, (Vancomycin Intermediate Resistant (VISA)) [HIP 5836]
Staphylococcus aureus, (Vancomycin Resistant (VISA))[VRS1]
Staphylococcus epidermidis, (Methicillin Resistant (MRSE)) (ATCC 51625)
Streptococcus pneumoniae, (Penicillin Resistant (PRSP)) (ATCC 700677)

***Virucidal** in the presence of 5% serum load with a 30 second contact time, unless otherwise noted.

*Viruses:

*Adenovirus Type 8, (ATCC VR-1368)
 *Cytomegalovirus, (ATCC VR-538)
 *Hepatitis B Virus (HBV), Duck Hepatitis B Virus as surrogate
 (Acquired from Hepadnavirus Testing)
 *Hepatitis C Virus (HCV), Bovine Viral Diarrhea
 Virus as surrogate (Acquired from National Veterinary
 Services Laboratories)

*Herpes Simplex Virus, Type 1 (HSV-1) (ATCC VR-733)
 *Herpes Simplex Virus, Type 2 (HSV-2), (ATCC VR-734)
 *HIV-1 (AIDS Virus), Strain HTLV-III_B (HIV-1),
 (Acquired from Adv. Biotechnologies, Inc.)
 *Human Coronavirus, (ATCC VR-740)
 *Influenza A Virus (Hong Kong Strain), (ATCC VR-544)
 *Influenza B Virus, (ATCC VR-823)

*Norovirus, (Feline Calicivirus, ATCC VR-782 as the surrogate)
 *Pandemic Influenza A Virus H1N1
 *Parainfluenza Virus Type 3, (ATCC VR-93)
 *Respiratory Syncytial Virus, (RSV) (ATCC VR-26)
 *Rhinovirus Type 37, Strain 151-1, (ATCC VR-1147)
 *Rotavirus, (WA strain) (Acquired from University of Ottawa)
 *Vaccinia Virus, (ATCC VR-119)

*Animal Viruses:

*Avian Influenza A Virus (H7N9)
 *Avian Influenza A Virus (H3N2), (ATCC VR-2072)

*Bovine Viral Diarrhea Virus (BVDV)
 *Duck Hepatitis B Virus
 *Equine Influenza A Virus (H3N8) (EIV)

*Feline Calicivirus, Strain F9 (ATCC VR-782)
 *Murine Norovirus

*Animal Viruses with a 1 minute contact time:

*Feline Infectious Peritonitis Virus, (FIPV) (ATCC VR-2022)
 *Murine Hepatitis Virus, (ATCC VR-765)

*Feline Herpes Virus, (Feline Rhinotracheitis) (ATCC VR-636)

*Minute Virus of Mouse, (Murine Parvovirus) (ATCC VR-1346)

TUBERCULOCIDAL against *Mycobacterium bovis* (Tb) (BCG) (OT 451C150) in the presence of a 5% serum load and 1 minute contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

FUNGICIDAL: This product is fungicidal against the pathogenic fungi *Trichophyton mentagrophytes* (ATCC 9533) (the fungus which causes Athlete's Foot) in the presence of a 5% soil and 30 second contact time when used as directed on hard surfaces found in bathrooms, shower stalls, locker rooms, or other clean, non-porous, hard surfaces commonly contacted by bare feet

FUNGICIDAL against *Candida albicans* (ATCC 10231) and *Microsporium canis* (ATCC 10214) in the presence of 5% soil with a 1 minute contact time.

Mold/Mildew (Mildewcidal Activity): Kills the growth of mold and mildew with a 4 minute contact time: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) (ATCC 16404) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Mold/Mildew (Mildewstatic Activity): Controls and prevents the growth of mold and mildew: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Non-Food Contact Surface Sanitizer for hard non-porous surfaces in the presence of a 5% serum load and 10 second contact time on hard, non-porous environmental surfaces.

Acinetobacter baumannii, (ATCC 19606)
Campylobacter jejuni, (ATCC 29428)
Enterobacter aerogenes, (ATCC 13048)
Enterococcus faecalis, (Vancomycin Resistant (VRE)) (ATCC 51575)

Escherichia coli O157:H7, (ATCC 35150)
Klebsiella pneumoniae, (ATCC 4352)
Listeria monocytogenes, (ATCC 19111)
Pseudomonas aeruginosa, (ATCC 15442)

Salmonella enterica, (formally known as *Salmonella choleraesuis*) (ATCC 10708)
Staphylococcus aureus, (ATCC 6538)
Staphylococcus aureus, (Methicillin Resistant (MRSA)) (ATCC 33592)

Malodor Counteractancy: eliminates odors and odor-causing bacteria on hard, nonporous surfaces in restroom areas, behind and under sinks and counters, and storage areas and other places where bacterial growth can cause malodors.

• Kills Pandemic 2009 H1N1 Influenza A virus.

See the container label for use directions and additional required information for use, storage and disposal.

EPA Reg. No. 70627-77

Manufactured for: ©2016 Diversey, Inc., PO Box 19747, Charlotte, NC 28219-0747

Oxivir® 1 Wipes

***Virucide • Bactericide • Tuberculocide • Fungicide**

For Institutional Use

Active Ingredient:

Hydrogen Peroxide..... 0.5%

Other Ingredients: 99.5%

Total: 100.0%

KEEP OUT OF REACH OF CHILDREN

BACTERICIDAL: 1 MINUTE

VIRUCIDAL: 30 SECONDS

ANIMAL PREMISE *VIRUCIDAL: 1 MINUTE

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER: 10 SECONDS

FUNGICIDAL: 30 SECONDS

Net Contents: 160 Wipes
11 in. x 12 in. (28 cm x 30 cm)
5.18 lb. (2.35 kg)



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO.
8017.176



AHP Technology and Logos are registered trademarks of Virox Technologies Inc., used under license.

This product has been developed to aid in the reduction of cross-contamination between treated surfaces in: airports, athletic facilities, correctional facilities, cruise ships, day care centers, health clubs. This product can be used in: clinics, hospitals, nursing homes, operating rooms, patient rooms, rehabilitation centers, office buildings, pet shops, public restrooms, retail businesses, schools, and veterinary clinics.

For daily cleaning and disinfecting of hard, nonporous inanimate surface materials: acrylic, Corian®, chrome, glass, glazed tile, laminated surfaces, linoleum, metal (except brass or copper), painted surfaces, and plastic.

Note: This product is compatible with the listed materials. If product is intended to be used on other surfaces, it is recommended to test product by applying to a small discrete area to determine compatibility before proceeding with its use.

This product cleans, disinfects and deodorizes hard, nonporous surfaces in one step. May be used to clean and disinfect sinks, faucets, fixtures, bathrooms, restrooms, tubs, and showers.

It can be used on the following hard, nonporous surfaces: chairs, counters, faucets, sinks, urinals, exercise equipment, floors, lockers, showers, tubs, and walls. Do not use on glasses, dishes and utensils.

100850924



DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not recommended for use on: brass or copper.

When used as directed, this product is highly effective against a broad-spectrum of pathogenic microorganisms. Using approved AOAC test methods, in the presence of 5% serum load and 1 minute contact time, unless otherwise noted, this product kills the following on hard nonporous environmental surfaces: **Bacteria:** *Pseudomonas aeruginosa*, *Staphylococcus aureus*, **Antibiotic-Resistant Bacteria** *Staphylococcus aureus* Methicillin Resistant (MRSA), *Staphylococcus aureus* Vancomycin Intermediate Resistant (VISA), *Staphylococcus aureus* Vancomycin Resistant (VRSA). ***Virucidal**, in the presence of 5% serum load with a 30 second contact time, unless otherwise noted. ***Viruses** *Rhinovirus Type 37, *Norovirus (Feline Calicivirus, as the surrogate) ***Animal Virus** *Murine Norovirus.

See reference sheet for a complete list of pathogenic organisms.

Not for personal cleansing. This is not a baby wipe. Do not flush in toilet. This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.



To Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Wipe hard, non-porous environmental surfaces with towelette.
3. Allow surface to remain wet for 1 minute. For certain listed viruses, allow to remain wet for 30 seconds.
4. Allow to air dry or rinse with potable water if necessary.

Note: All food contact surfaces must be rinsed with potable water. Do not use on glassware, utensils, or dishes.

To Use as a Restroom, Shower Room, Locker Room, Athletic Equipment Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Wipe hard, non-porous environmental surfaces with towelette.
3. Allow surfaces to remain wet for 1 minute. For certain listed viruses, allow to remain wet for 30 seconds.
4. Rinse surfaces thoroughly and let air dry.

Note: Floors will be slippery when wet or contaminated with foreign materials.

***KILLS HIV, HCV and HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (Hospitals, Nursing Homes) and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis C Virus (HCV) or Hepatitis B Virus (HBV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS), HCV OR HBV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood/body fluids must be thoroughly cleaned from surfaces/objects before application of this product.

Contact Time: Allow surface to remain wet for 30 seconds to kill HIV-1, HCV and HBV, 4 minutes for mold, and 1 minute to kill other organisms cited on label.

Disposal of Infectious Material: Blood/body fluids must be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

Animal Housing Facilities Including Veterinary Clinics, Equine Farms, Swine and Poultry Houses:

For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities (including barns, pens, stalls, chutes, and barns) occupied or traversed by animals.
3. Empty or cover all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water.
5. Wipe hard, non-porous environmental surfaces with towelettes. To disinfect, all surfaces must remain wet for 1 minute. For certain listed viruses, allow to remain wet for 30 seconds.
6. Allow to air dry or rinse with potable water if necessary.
7. Ventilate buildings and other closed spaces.
8. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
9. Thoroughly scrub treated feed racks, troughs, mangers, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

To Kill Fungi: Pre-clean heavily soiled areas. Wipe hard, non-porous environmental surfaces with towelette. Allow to remain wet for 1 minute. Use additional towelettes if needed to ensure the correct wet contact time. Wipe surfaces or let air dry.

To Kill Mold and Mildew: Pre-clean surfaces. Wipe hard, nonporous surfaces with towelettes until completely wet. Allow to remain wet for 4 minutes. Let air dry.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Store in original container in areas inaccessible to small children in a cool, dry area away from direct sunlight and heat. Keep securely closed.

Towelette Storage/Disposal: Dispose of wipe in trash after use. Do not flush. Tightly close lid between uses to retain moisture.

Container Disposal: Refillable container. Refill this container with this brand name pesticide only. Do not reuse or refill this container for any other purpose. After final use, offer empty container for recycling. If recycling is not available, discard container in trash.



See SDS # MS0800916

EPA Reg. No. 70627-77
EPA Est. No. 74559-CAN-1 [VX]
Lot code letters indicate establishment number.

EMERGENCY PHONE : 1-800-851-7145

Manufactured for: ©2016 Diversey, Inc., PO Box 19747, Charlotte, NC 28219-0747

Oxivir® 1 Wipes

When used as directed, this product is highly effective against a wide-variety (broad-spectrum) of pathogenic microorganisms including bacteria, antibiotic-resistant bacteria, *viruses, and fungi. Using approved AOAC test methods, in the presence of 5% serum load and 1 minute contact time, unless otherwise noted, this product kills the following on hard nonporous inanimate surfaces:

Bacteria:

Pseudomonas aeruginosa, (ATCC 15442)
Staphylococcus aureus, (ATCC 6538)
Salmonella enterica, (ATCC 10708)
 formerly known as *Salmonella choleraesuis*
Bordetella bronchiseptica, (ATCC 10580)
Burkholderia cepacia, (ATCC 25416)
Bordetella pertussis, (ATCC 12743)

Enterobacter aerogenes, (ATCC 13048)
Enterococcus faecium, (ATCC 51559)
Escherichia coli, (ATCC 11229)
Escherichia coli O157:H7, (ATCC 35150)
Klebsiella pneumoniae, (ATCC 4352)
Legionella pneumophila, (ATCC 33153)

Pasteurella multocida, (ATCC 12947)
Serratia marcescens, (ATCC 14756)
Staphylococcus pseudintermedius, (ATCC 49444)
Shigella dysenteriae, (ATCC 11835)
Stenotrophomonas (Pseudomonas) maltophilia, (ATCC 25596)
Streptococcus pyogenes, (ATCC19615)

Antibiotic-Resistant Bacteria:

Acinetobacter baumannii,

Multi-Drug Resistant (MDR) (ATCC 19606)
 Ampicillin, Cefazolin, Cefepime, Gentamicin,
 Ertapenem, and Trimethoprim/Sulfa

Enterobacter cloacae, (New Delhi metallo-β-lactamase
 1 positive (producing) (NDM 1)) (CDC 1000654)

Enterococcus faecalis, (ATCC 51575) (Vancomycin Resistant (VRE))

Escherichia coli, Extended Spectrum Beta-lactamase
 producing (positive) (ESBL) (ATCC BAA-196)

Escherichia coli, (New Delhi metallo-β-lactamase 1(NDM-1) producing
 (positive)) (CDC 1001728)

Klebsiella pneumoniae, (Carbapenem Resistant (CRE)) (ATCC BAA 1705)
Klebsiella pneumoniae, (Extended Spectrum beta-lactamase (ESBL) producing
 (positive)) (ATCC 700603)

Klebsiella pneumoniae, (New Delhi metallo-β-lactamase 1(NDM-1)
 producing (positive)) (CDC 1000527)

Pseudomonas aeruginosa, (Metallo-beta lactamase (MBL)
 positive (CDC 2012059))

Staphylococcus aureus, Community Associated Methicillin Resistant (CA-MRSA)

(NARSA NRS 123) (Genotype USA400)

Staphylococcus aureus, (Methicillin Resistant (MRSA)) (ATCC 33591)

Staphylococcus aureus, (ATCC 33592) (Methicillin Resistant (MRSA))

Staphylococcus aureus, (Vancomycin Intermediate Resistant (VISA)) [HIP 5836]

Staphylococcus aureus, (Vancomycin Resistant (VISA))[VRS1]

Staphylococcus epidermidis, (Methicillin Resistant (MRSE)) (ATCC 51625)

Streptococcus pneumoniae, (Penicillin Resistant (PRSP)) (ATCC 700677)

***Virucidal** in the presence of 5% serum load with a 30 second contact time, unless otherwise noted.

*Viruses:

*Adenovirus Type 8, (ATCC VR-1368)

*Cytomegalovirus, (ATCC VR-538)

*Hepatitis B Virus (HBV), Duck Hepatitis B Virus as surrogate
 (Acquired from Hepadnavirus Testing)

*Hepatitis C Virus (HCV), Bovine Viral Diarrhea

Virus as surrogate (Acquired from National Veterinary
 Services Laboratories)

*Herpes Simplex Virus, Type 1 (HSV-1) (ATCC VR-733)

*Herpes Simplex Virus, Type 2 (HSV-2), (ATCC VR-734)

*HIV-1 (AIDS Virus), Strain HTLV-III_B (HIV-1),

(Acquired from Adv. Biotechnologies, Inc.)

*Human Coronavirus, (ATCC VR-740)

*Influenza A Virus (Hong Kong Strain), (ATCC VR-544)

*Influenza B Virus, (ATCC VR-823)

*Norovirus, (Feline Calicivirus, ATCC VR-782 as the surrogate)

*Pandemic Influenza A Virus H1N1

*Parainfluenza Virus Type 3, (ATCC VR-93)

*Respiratory Syncytial Virus, (RSV) (ATCC VR-26)

*Rhinovirus Type 37, Strain 151-1, (ATCC VR-1147)

*Rotavirus, (WA strain) (Acquired from University of Ottawa)

*Vaccinia Virus, (ATCC VR-119)

*Animal Viruses:

*Avian Influenza A Virus (H7N9)

*Avian Influenza A Virus (H3N2), (ATCC VR-2072)

*Bovine Viral Diarrhea Virus (BVDV)

*Duck Hepatitis B Virus

*Equine Influenza A Virus (H3N8) (EIV)

*Feline Calicivirus, Strain F9 (ATCC VR-782)

*Murine Norovirus

*Animal Viruses with a 1 minute contact time:

*Feline Infectious Peritonitis Virus, (FIPV) (ATCC VR-2022)

*Murine Hepatitis Virus, (ATCC VR-765)

*Feline Herpes Virus, (Feline Rhinotracheitis) (ATCC VR-636)

*Minute Virus of Mouse, (Murine Parvovirus) (ATCC VR-1346)

TUBERCULOCIDAL against *Mycobacterium bovis* (Tb) (BCG) (OT 451C150) in the presence of a 5% serum load and 1 minute contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

FUNGICIDAL: This product is fungicidal against the pathogenic fungi *Trichophyton mentagrophytes* (ATCC 9533) (the fungus which causes Athlete's Foot) in the presence of a 5% soil and 30 second contact time when used as directed on hard surfaces found in bathrooms, shower stalls, locker rooms, or other clean, non-porous, hard surfaces commonly contacted by bare feet

FUNGICIDAL against *Candida albicans* (ATCC 10231) and *Microsporium canis* (ATCC 10214) in the presence of 5% soil with a 1 minute contact time.

Mold/Mildew (Mildewcidal Activity): Kills the growth of mold and mildew with a 4 minute contact time: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) (ATCC 16404) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Mold/Mildew (Mildewstatic Activity): Controls and prevents the growth of mold and mildew: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Non-Food Contact Surface Sanitizer for hard non-porous surfaces in the presence of a 5% serum load and 10 second contact time on hard, non-porous environmental surfaces.

Acinetobacter baumannii, (ATCC 19606)

Campylobacter jejuni, (ATCC 29428)

Enterobacter aerogenes, (ATCC 13048)

Enterococcus faecalis, (Vancomycin Resistant (VRE)) (ATCC 51575)

Escherichia coli O157:H7, (ATCC 35150)

Klebsiella pneumoniae, (ATCC 4352)

Listeria monocytogenes, (ATCC 19111)

Pseudomonas aeruginosa, (ATCC 15442)

Salmonella enterica, (formally known as *Salmonella choleraesuis*) (ATCC 10708)

Staphylococcus aureus, (ATCC 6538)

Staphylococcus aureus, (Methicillin Resistant (MRSA)) (ATCC 33592)

Malodor Counteractancy: eliminates odors and odor-causing bacteria on hard, nonporous surfaces in restroom areas, behind and under sinks and counters, and storage areas and other places where bacterial growth can cause malodors.

• Kills Pandemic 2009 H1N1 Influenza A virus.

See the container label for use directions and additional required information for use, storage and disposal.

EPA Reg. No. 70627-77

Manufactured for: ©2016 Diversey, Inc., PO Box 19747, Charlotte, NC 28219-0747

Oxivir® 1 Wipes

*Virucide • Bactericide • Tuberculocide • Fungicide
For Institutional Use

Active Ingredient:

Hydrogen Peroxide.....0.5%

Other Ingredients:.....99.5%

Total:.....100.0%

KEEP OUT OF REACH OF CHILDREN

BACTERICIDAL: 1 MINUTE

VIRUCIDAL: 30 SECONDS

ANIMAL PREMISE *VIRUCIDAL: 1 MINUTE

BROAD-SPECTRUM NON-FOOD CONTACT SANITIZER: 10 SECONDS

FUNGICIDAL: 30 SECONDS



AHP Technology and Logos are registered trademarks of Virox Technologies Inc., used under license.

Net Contents: 160 Wipes
11 in. x 12 in. (28 cm x 30 cm)
5.18 lb. (2.35 kg)

100850925

Sealed Air
Diversey Care

Diversey®



DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not recommended for use on: brass or copper.

When used as directed, this product is highly effective against a wide variety (broad-spectrum) of pathogenic Microorganisms. Using approved AOAC test methods, in the presence of 5% serum load and 1 minute contact time, unless otherwise noted, this product kills the following on hard nonporous environmental surfaces: **Bacteria:** *Pseudomonas aeruginosa*, *Staphylococcus aureus*,

Antibiotic-Resistant Bacteria *Staphylococcus aureus* Methicillin Resistant (MRSA), *Staphylococcus aureus* Vancomycin Intermediate Resistant (VISA), *Staphylococcus aureus* Vancomycin Resistant (VISA). ***Virucidal**, in the presence of 5% serum load with a 30 second contact time, unless otherwise noted. ***Viruses:** *Rhinovirus Type 37, *Norovirus (Feline Calicivirus, as the surrogate) ***Animal Viruses:** *Murine Norovirus.

See reference sheet for a complete list of pathogenic organisms.

Not for personal cleansing. This is not a baby wipe. Do not flush in toilet.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

To Use as a One-Step Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Wipe hard, non-porous environmental surfaces with towelette.
3. Allow surface to remain wet for 1 minute. For certain listed viruses, allow to remain wet for 30 seconds.
4. Allow to air dry or rinse with potable water if necessary.

Note: All food contact surfaces must be rinsed with potable water. Do not use on glassware, utensils, or dishes.

To Use as a Restroom, Shower Room, Locker Room, Athletic Equipment Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Wipe hard, non-porous environmental surfaces with towelette.
3. Allow surfaces to remain wet for 1 minute. For certain listed viruses, allow to remain wet for 30 seconds.
4. Rinse surfaces thoroughly and let air dry.

Note: Floors will be slippery when wet or contaminated with foreign materials.

*KILLS HIV, HCV and HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS

PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings (Hospitals, Nursing Homes) and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis C Virus (HCV) or Hepatitis B Virus (HBV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS), HCV OR HBV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood/body fluids must be thoroughly cleaned from surfaces/objects before application of this product.

Contact Time: Allow surface to remain wet for 30 seconds to kill HIV-1, HCV and HBV, 4 minutes for mold, and 1 minute to kill other organisms cited on label.

Disposal of Infectious Material: Blood/body fluids must be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

Animal Housing Facilities Including Veterinary Clinics, Equine Farms, Swine and Poultry Houses:

For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities (including barns, pens, stalls, chutes, and barns) occupied or traversed by animals.
3. Empty or cover all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water.
5. Wipe hard, non-porous environmental surfaces with towelettes. To disinfect, all surfaces must remain wet for 1 minute. For certain listed viruses, allow to remain wet for 30 seconds.
6. Allow to air dry or rinse with potable water if necessary.
7. Ventilate buildings and other closed spaces.
8. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
9. Thoroughly scrub treated feed racks, troughs, mangers, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

To Kill Fungi: Pre-clean heavily soiled areas. Wipe hard, non-porous environmental surfaces with towelette. Allow to remain wet for 1 minute. Use additional towelettes if needed to ensure the correct wet contact time. Wipe surfaces or let air dry.

To Kill Mold and Mildew: Pre-clean surfaces. Wipe hard, nonporous surfaces with towelettes until completely wet. Allow to remain wet for 4 minutes. Let air dry.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Store in original container in areas inaccessible to small children in a cool, dry area away from direct sunlight and heat. Keep securely closed.

Towelette Storage/Disposal: Dispose of wipe in trash after use. Do not flush. Tightly close lid between uses to retain moisture.

Container Disposal: Non-refillable container. Do not reuse or refill this container. Discard container in trash.

EMERGENCY PHONE : 1-800-851-7145

See SDS # MS0800916

Questions? Comments
1-800-558-2332

EPA Reg. No. 70627-77

EPA Est. No. 74559-CAN-1 [VX]

Lot code letters indicate establishment number.



Manufactured for:

©2016 Diversey, Inc., PO Box 19747,
Charlotte, NC 28219-0747

300153659 (16/126)



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO.

8017.176

Oxivir® 1 Wipes

When used as directed, this product is highly effective against a wide-variety (broad-spectrum) of pathogenic microorganisms including bacteria, antibiotic-resistant bacteria, *viruses, and fungi. Using approved AOAC test methods, in the presence of 5% serum load and 1 minute contact time, unless otherwise noted, this product kills the following on hard nonporous inanimate surfaces:

Bacteria:

Pseudomonas aeruginosa, (ATCC 15442)
Staphylococcus aureus, (ATCC 6538)
Salmonella enterica, (ATCC 10708)
 formerly known as *Salmonella choleraesuis*
Bordetella bronchiseptica, (ATCC 10580)
Burkholderia cepacia, (ATCC 25416)
Bordetella pertussis, (ATCC 12743)

Enterobacter aerogenes, (ATCC 13048)
Enterococcus faecium, (ATCC 51559)
Escherichia coli, (ATCC 11229)
Escherichia coli O157:H7, (ATCC 35150)
Klebsiella pneumoniae, (ATCC 4352)
Legionella pneumophila, (ATCC 33153)

Pasteurella multocida, (ATCC 12947)
Serratia marcescens, (ATCC 14756)
Staphylococcus pseudintermedius, (ATCC 49444)
Shigella dysenteriae, (ATCC 11835)
Stenotrophomonas (Pseudomonas) maltophilia, (ATCC 25596)
Streptococcus pyogenes, (ATCC19615)

Antibiotic-Resistant Bacteria:

Acinetobacter baumannii.

Multi-Drug Resistant (MDR) (ATCC 19606)
 Ampicillin, Cefazolin, Cefepime, Gentamicin,
 Ertapenem, and Trimethoprim/Sulfa

Enterobacter cloacae, (New Delhi metallo- β -lactamase
 1 positive (producing) (NDM 1)) (CDC 1000654)

Enterococcus faecalis, (ATCC 51575) (Vancomycin Resistant (VRE))

Escherichia coli, Extended Spectrum Beta-lactamase
 producing (positive) (ESBL) (ATCC BAA-196)

Escherichia coli, (New Delhi metallo- β -lactamase 1(NDM-1) producing
 (positive)) (CDC 1001728)

Klebsiella pneumoniae, (Carbapenem Resistant (CRE)) (ATCC BAA 1705)
Klebsiella pneumoniae, (Extended Spectrum beta-lactamase (ESBL) producing
 (positive)) (ATCC 700603)

Klebsiella pneumoniae, (New Delhi metallo- β -lactamase 1(NDM-1)
 producing (positive)) (CDC 1000527)

Pseudomonas aeruginosa, (Metallo-beta lactamase (MBL)
 positive (CDC 2012059))

Staphylococcus aureus, Community Associated Methicillin Resistant (CA-MRSA)
 (NARSA NRS 123) (Genotype USA400)

Staphylococcus aureus, (Methicillin Resistant (MRSA)) (ATCC 33591)

Staphylococcus aureus, (ATCC 33592) (Methicillin Resistant (MRSA))

Staphylococcus aureus, (Vancomycin Intermediate Resistant (VISA)) [HIP 5836]

Staphylococcus aureus, (Vancomycin Resistant (VISA))[VRS1]

Staphylococcus epidermidis, (Methicillin Resistant (MRSE)) (ATCC 51625)

Streptococcus pneumoniae, (Penicillin Resistant (PRSP)) (ATCC 700677)

***Virucidal** in the presence of 5% serum load with a 30 second contact time, unless otherwise noted.

*Viruses:

*Adenovirus Type 8, (ATCC VR-1368)

*Cytomegalovirus, (ATCC VR-538)

*Hepatitis B Virus (HBV), Duck Hepatitis B Virus as surrogate
 (Acquired from Hepadnavirus Testing)

*Hepatitis C Virus (HCV), Bovine Viral Diarrhea

Virus as surrogate (Acquired from National Veterinary
 Services Laboratories)

*Herpes Simplex Virus, Type 1 (HSV-1) (ATCC VR-733)

*Herpes Simplex Virus, Type 2 (HSV-2), (ATCC VR-734)

*HIV-1 (AIDS Virus), Strain HTLV-III_B (HIV-1),
 (Acquired from Adv. Biotechnologies, Inc.)

*Human Coronavirus, (ATCC VR-740)

*Influenza A Virus (Hong Kong Strain), (ATCC VR-544)

*Influenza B Virus, (ATCC VR-823)

*Norovirus, (Feline Calicivirus, ATCC VR-782 as the surrogate)

*Pandemic Influenza A Virus H1N1

*Parainfluenza Virus Type 3, (ATCC VR-93)

*Respiratory Syncytial Virus, (RSV) (ATCC VR-26)

*Rhinovirus Type 37, Strain 151-1, (ATCC VR-1147)

*Rotavirus, (WA strain) (Acquired from University of Ottawa)

*Vaccinia Virus, (ATCC VR-119)

*Animal Viruses:

*Avian Influenza A Virus (H7N9)

*Avian Influenza A Virus (H3N2), (ATCC VR-2072)

*Bovine Viral Diarrhea Virus (BVDV)

*Duck Hepatitis B Virus

*Equine Influenza A Virus (H3N8) (EIV)

*Feline Calicivirus, Strain F9 (ATCC VR-782)

*Murine Norovirus

*Animal Viruses with a 1 minute contact time:

*Feline Infectious Peritonitis Virus, (FIPV) (ATCC VR-2022)

*Murine Hepatitis Virus, (ATCC VR-765)

*Feline Herpes Virus, (Feline Rhinotracheitis) (ATCC VR-636)

*Minute Virus of Mouse, (Murine Parvovirus) (ATCC VR-1346)

TUBERCULOCIDAL against *Mycobacterium bovis* (Tb) (BCG) (OT 451C150) in the presence of a 5% serum load and 1 minute contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

FUNGICIDAL: This product is fungicidal against the pathogenic fungi *Trichophyton mentagrophytes* (ATCC 9533) (the fungus which causes Athlete's Foot) in the presence of a 5% soil and 30 second contact time when used as directed on hard surfaces found in bathrooms, shower stalls, locker rooms, or other clean, non-porous, hard surfaces commonly contacted by bare feet

FUNGICIDAL against *Candida albicans* (ATCC 10231) and *Microsporium canis* (ATCC 10214) in the presence of 5% soil with a 1 minute contact time.

Mold/Mildew (Mildewcidal Activity): Kills the growth of mold and mildew with a 4 minute contact time: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) (ATCC 16404) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Mold/Mildew (Mildewstatic Activity): Controls and prevents the growth of mold and mildew: *Aspergillus brasiliensis* (formerly known as *Aspergillus niger*) and the odors caused by them when applied to hard, non-porous environmental surfaces.

Non-Food Contact Surface Sanitizer for hard non-porous surfaces in the presence of a 5% serum load and 10 second contact time on hard, non-porous environmental surfaces.

Acinetobacter baumannii, (ATCC 19606)

Campylobacter jejuni, (ATCC 29428)

Enterobacter aerogenes, (ATCC 13048)

Enterococcus faecalis, (Vancomycin Resistant (VRE)) (ATCC 51575)

Escherichia coli O157:H7, (ATCC 35150)

Klebsiella pneumoniae, (ATCC 4352)

Listeria monocytogenes, (ATCC 19111)

Pseudomonas aeruginosa, (ATCC 15442)

Salmonella enterica, (formally known as *Salmonella choleraesuis*) (ATCC 10708)

Staphylococcus aureus, (ATCC 6538)

Staphylococcus aureus, (Methicillin Resistant (MRSA)) (ATCC 33592)

Malodor Counteractancy: eliminates odors and odor-causing bacteria on hard, nonporous surfaces in restroom areas, behind and under sinks and counters, and storage areas and other places where bacterial growth can cause malodors.

• Kills Pandemic 2009 H1N1 Influenza A virus.

See the container label for use directions and additional required information for use, storage and disposal.

EPA Reg. No. 70627-77

Manufactured for: ©2016 Diversey, Inc., PO Box 19747, Charlotte, NC 28219-0747