

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Do not reuse empty container. Keep from freezing.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Nonrefillable container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer container for recycling, if available.

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin or on clothing. Wear chemical splash-proof goggles or face shield, rubber gloves and protective clothing. Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

EPA Reg. No. 70627-24
EPA Est. No. 0312-WI-3 (NW)

FOR INSTITUTIONAL USE

Questions? Comments:
1-800-558-2332



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO.

8017.33

Sealed Air
Diversey Care

Diversey™

5

Virex® II 256

One-Step Disinfectant Cleaner And Deodorant

Bactericidal • *Virucidal • Fungicidal • Mildewcidal • Deodorizer • Mildewstatic

Meets OSHA Bloodborne Pathogen Standard for *HBV & *HIV

ACTIVE INGREDIENTS:

Didecyl dimethyl ammonium chloride 8.704%
n-Alkyl (50% C14, 40% C12, 10% C16)
dimethyl benzyl ammonium chloride..... 8.190%

OTHER INGREDIENTS: 83.106%

TOTAL: 100.000%

KEEP OUT OF REACH OF CHILDREN DANGER

See additional precautionary statements on back panel.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.

**IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR
DOCTOR FOR TREATMENT ADVICE.**

1-800-851-7145

Have the product container or label with you when calling a
Poison Control Center or doctor or going in for treatment.

Note to Physician: Probable mucosal damage may contraindicate gastric lavage.

Net Contents:

3.78 L / 1 U.S. Gal.

04332

This product is a one-step disinfectant/cleaner and deodorizer designed for general cleaning, disinfecting and deodorizing of hard, non-porous environmental surfaces. It cleans quickly by removing dirt, grime, mold, mildew, body oils and other common soils found in hospitals, nursing homes, schools and colleges, medical offices, hotels, motels, public areas, restrooms and foodservice establishments.

It is designed for use on the following hard, non-porous environmental surfaces: vinyl, painted surfaces, plastic, glazed ceramic, glazed porcelain, chrome, aluminum, laminated surfaces and baked enamel surfaces associated with floors, walls, ceilings, tables, chairs, countertops, telephones, fixtures, glazed tile, toilets, urinals, sinks, shower rooms and locker rooms areas – any hard, non-porous washable surface where disinfection is required. This products non-dulling formula eliminates the time and labor normally required for rinsing. A potable water rinse is required for food contact surfaces. Do not use on glasses, dishes and utensils.

When used as directed at a 1:256 dilution (1/2 oz. per gallon of water), this product contains 660 ppm of active quaternary germicide making it highly effective against a wide variety of pathogenic microorganisms.

Using approved AOC test methods under Good Laboratory Practices, in the presence of 400 ppm hard water, 10% serum load and 10 minute contact time, this product kills the following on hard non-porous inanimate surfaces:

Bacteria – *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Salmonella enterica*, *Acinetobacter baumannii*, *Acinetobacter calcoaceticus*, *Enterococcus faecalis*, *Escherichia coli*, *Escherichia coli* O157:H7, *Klebsiella pneumoniae*, *Listeria monocytogenes*, *Proteus mirabilis*, *Proteus vulgaris*, *Salmonella enteritidis*, *Salmonella typhi*, *Serratia marcescens*, *Shigella dysenteriae*, *Staphylococcus epidermidis*, *Streptococcus agalactiae*, *Streptococcus pyogenes*

Antibiotic-Resistant Bacteria – *Enterococcus faecalis*, Resistant to Vancomycin (VRE); *Staphylococcus aureus*, Intermediate Vancomycin Resistance (VISA); *Staphylococcus aureus*, Resistant to Methicillin (MRSA) and Gentamicin (GRSA); *Staphylococcus epidermidis*, Resistant to Methicillin (MRSE); *Streptococcus pneumoniae*, Resistant to Penicillin (PRSP)

***Viruses** – *Cytomegalovirus, *Herpes Simplex virus Type 1, *Herpes Simplex virus Type 2, *Human Coronavirus, *Influenza virus Type A2 (Hong Kong), *Parinfluenza virus Type 3, *Respiratory syncytial virus, *Rotavirus, *Vaccinia virus. Kills *HIV-1 (AIDS virus) when used as directed on hard, non-porous inanimate surfaces with a 1 minute contact time. Kills *HBV & *HCV when used as directed on hard, non-porous inanimate surfaces with a 5 minute contact time.

*Kills Pandemic 2009 H1N1 Influenza A virus

Manufactured for: ©2015 Diversey, Inc., PO Box 19747, Charlotte, NC 28219-0747, U.S.A. All Rights Reserved.

SDS # MS0800549



Fungi – *Geotrichum candidum*, *Saccharomyces cerevisiae*.

Bactericidal Stability of Use-Dilution – Tests show this product, when diluted in 400 ppm hard water and in the presence of 5% serum load, remains effective against *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Salmonella enterica* for up to 1 year in storage as long as it remains sealed. If product becomes visibly dirty or contaminated, the use-dilution must be discarded and fresh product prepared. Always use clean, dry containers when diluting this product.

Using approved AOC test methods under Good Laboratory Practices, in the presence of 400 ppm hard water, 5% serum load and 10 minute contact time, this product kills the following on hard non-porous inanimate surfaces:

***Viruses** – *Adenovirus Type 2.

Fungi: *Aspergillus niger*, *Trichophyton mentagrophytes* (athlete's foot fungus).

Yeast: *Candida albicans*.

Mold/Mildew – kills the growth of mold and mildew: *Aspergillus niger* and the odors caused by them when applied to hard, non-porous environmental surfaces.

See reference sheet for additional directions for use and a complete list of organisms. This product can be applied by mop, sponge, cloth, paper towel, coarse trigger sprayer, auto-scrubber or foam gun. Change cloth, sponges or towels frequently to avoid redeposition of soil. For disinfection, all surfaces must remain wet for 10 minutes.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

To Prepare Use Solution: Add the product at 1/2 oz. per gallon of water (1:256).

For Use as a One-Step Cleaner/Disinfectant: Pre-clean heavily soiled areas. Apply Use Solution to hard, non-porous environmental surfaces. All surfaces must remain wet for 10 minutes. Wipe surfaces and let air dry.

Note – Rinsing is not necessary unless floors are to be coated with finish or restorer. All food contact surfaces such as appliances and kitchen countertops must be rinsed with potable water. Do not use on glassware, utensils, or dishes.

For Use as a Cleaner and Deodorizer: Apply Use Solution to hard, non-porous surfaces. Wipe surfaces and let air dry.

For Use as a Non-Acid Bowl Cleaner/Disinfectant in Toilet Bowls and Urinals from Use-Dilution: Pre-clean heavily soiled areas. Empty toilet bowls by forcing water through the trap. Apply Use Solution to exposed surfaces in toilet bowls and urinals. Swab entire surface area especially under the rim. Allow entire surface to remain wet for 10 minutes. Flush toilet or urinal and rinse swab applicator thoroughly.

For use as a Non-Acid Bowl Cleaner/Disinfectant in Toilet Bowls from Concentrate: Pre-clean heavily soiled areas. Add 3/8 oz. into toilet bowl for a 1:256 dilution. Swab entire surface area especially under the rim. Allow entire surface to remain wet for 10 minutes. Flush toilet and rinse swab applicator thoroughly.

For Use To Clean and Disinfect Shower Rooms, Locker Rooms and Other Large, Open Areas with Floor Drains: Pre-clean heavily soiled areas. Apply Use Solution to floors, walls and ceilings making sure not to over spray. To disinfect, all surfaces must remain wet for 10 minutes. *Special instructions for foam guns: Pour concentrate into foam gun bottle and attach bottle to spray nozzle and ensure gun is attached to hose.*

Note: See foam gun instructions for more information. Make sure setting is set for a 1:256 dilution. Once in place, squeeze the handle to dispense foam solution. To disinfect, all surfaces must remain wet for 10 minutes. Scrub using a deck brush or other coarse material as necessary. Rinse surfaces thoroughly and let air dry. NOTE: When cleaning floors position wet-floor signs around area to be cleaned. Floors will be slippery when wet or contaminated with foreign materials. Promptly clean up spills and foreign materials.

To Kill Mold and Mildew (in 5% soil load): Pre-clean heavily soiled areas. Apply Use Solution to hard, non-porous environmental surfaces. Allow surfaces to remain wet for 10 minutes. Wipe surfaces and let air dry.

To Control Mold and Mildew: Apply Use Solution to pre-cleaned hard, non-porous environmental surfaces. Allow to air dry. Repeat application weekly or when growth reappears.

To Kill Fungi: Pre-clean heavily soiled areas. Apply Use Solution to hard, non-porous environmental surfaces. Allow surface to remain wet for 10 minutes. Wipe surfaces and let air dry.

***Kills *HBV, *HCV and *HIV-1 on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids** in health care settings (Hospitals, Nursing Homes) and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of *Hepatitis B Virus (HBV), *Hepatitis C Virus (HCV) and *Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST *HBV, *HCV and *HIV-1 ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood and body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

Contact Time: Allow surface to remain wet for 1 minute to kill *HIV-1, 5 minutes to kill *HBV and *HCV, and for 10 minutes to kill all other organisms cited on the label.

Disposal of Infectious Material: Blood and other body fluids must be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

To Sanitize Non-Food Contact Surfaces:

1. Pre-clean soiled hard non-porous surfaces.
2. Apply this product until surface is thoroughly wet.
3. Let stand 1 minute, then wipe.

Note: Not for use on food contact surfaces or on food preparation areas.

Virex® II 256



This product is a one-step disinfectant cleaner and deodorant designed for general cleaning, disinfecting and deodorizing of hard, non-porous environmental surfaces. It cleans quickly by removing dirt, grime, mold, mildew, body oils and other common soils found in hospitals, nursing homes, schools and colleges, medical offices, hotels, motels, public areas, restrooms and foodservice establishments.

It is designed for use on the following hard, non-porous environmental surfaces: vinyl, painted surfaces, plastic, glazed ceramic, glazed porcelain, chrome, aluminum, laminated surfaces and baked enamel surfaces associated with floors, walls, ceilings, tables, chairs, countertops, telephones, fixtures, glazed tile, toilets, urinals, sinks, shower rooms and locker room areas - any hard, non-porous washable surface where disinfection is required. This product's non-dulling formula eliminates the time and labor normally required for rinsing. A potable water rinse is required for food contact surfaces. Do not use on glasses, dishes and utensils.

When used as directed at a 1:256 dilution (½ oz. per gallon of water), this product contains 660 ppm of active quaternary germicide making it highly effective against a wide variety of pathogenic microorganisms.

Using approved AOAC test methods under Good Laboratory Practices, in the presence of 400 ppm hard water, 10% serum load and 10 minute contact time this product kills the following on hard non-porous inanimate surfaces:

Pseudomonas aeruginosa, *Staphylococcus aureus*, *Salmonella enterica*

See reference sheet for a complete list of organisms and additional directions for use.

This product can be applied by mop, sponge, cloth, paper towel, coarse trigger sprayer, auto-scrubber or foam gun. Change cloth, sponges or towels frequently to avoid redeposition of soil. For disinfection, all surfaces must remain wet for 10 minutes.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For Use as a Non-Acid Bowl Cleaner/Disinfectant in Toilet Bowls and Urinals from Use-Dilution:

Pre-clean heavily soiled areas. Empty toilet bowls by forcing water through the trap. Apply Use Solution to exposed surfaces in toilet bowls and urinals. Swab entire surface area especially under rim. Allow entire surface to remain wet for 10 minutes. Flush toilet or urinal and rinse swab applicator thoroughly.

For use as a Non-Acid Bowl Cleaner/Disinfectant in Toilet Bowls from Concentrate:

Pre-clean heavily soiled areas. Add 3/8 oz. into toilet bowl for a 1:256 dilution. Swab entire surface area especially under the rim. Allow entire surface to remain wet for 10 minutes. Flush toilet and rinse swab applicator thoroughly.

For Use To Clean and Disinfect Shower Rooms, Locker Rooms

and Other Large, Open Areas with Floor Drains: Pre-clean heavily soiled areas. Apply Use Solution to floors, walls and ceilings making sure not to over spray. To disinfect, all surfaces must remain wet for 10 minutes. *Special instructions for foam guns: Pour concentrate into foam gun bottle and attach bottle to spray nozzle and ensure gun is attached to hose. Note: See foam gun instructions for more information. Make sure setting is set for a 1:256 dilution. Once in place, squeeze the handle to dispense foam solution. To disinfect, all surfaces must remain wet for 10 minutes.* Scrub using a deck brush or other coarse material as necessary. Rinse surfaces thoroughly and let air dry.

NOTE: When cleaning floors position wet-floor signs around area to be cleaned. Floors will be slippery when wet or contaminated with foreign materials. Promptly clean up spills and foreign materials.

To Kill Mold and Mildew (in 5% soil load):

Pre-clean heavily soiled areas. Apply Use Solution to hard, non-porous environmental surfaces. Allow surfaces to remain wet for 10 minutes. Wipe surfaces and let air dry.

To Control Mold and Mildew:

Apply Use Solution to pre-cleaned hard, non-porous environmental surfaces. Allow to air dry. Repeat application weekly or when growth reappears.

To Kill Fungi:

Pre-clean heavily soiled areas. Apply Use Solution to hard, non-porous environmental surfaces. Allow surface to remain wet for 10 minutes. Wipe surfaces and let air dry.

***Kills *HBV, *HCV and *HIV-1 on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids** in health care settings (Hospitals, Nursing Homes) and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of *Hepatitis B Virus (HBV), *Hepatitis C Virus (HCV) and *Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST *HBV, *HCV and *HIV-1 ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood and body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

Contact Time: Allow surface to remain wet for 1 minute to kill *HIV-1, 5 minutes to kill *HBV and *HCV, and for 10 minutes to kill all other organisms cited on the label.

Disposal of Infectious Material: Blood and other body fluids must be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

To Sanitize Non-Food Contact Surfaces:

1. Pre-clean soiled hard non-porous surfaces.
2. Apply this product until surface is thoroughly wet.
3. Let stand 1 minute, then wipe.

Note: Not for use on food contact surfaces or on food preparation areas.

See container label for First Aid, Precautionary Statements and complete Directions for Use.

EPA Reg. No. 70627-24