



Oxivir[®] Tb

General *Virucide, Bactericide, Tuberculocide, Fungicide, Sanitizer

ACTIVE INGREDIENT:

Hydrogen Peroxide 0.5%

OTHER INGREDIENTS: 99.5%

TOTAL: 100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

See additional precautionary statements on side panel.



Net Contents:

946 mL / 1 U.S. Qt.



*AHP is a registered trademark of Virox Technologies Inc., used under license.

4277285

For Use in Health Care, Medical, Industrial & Institutional Facilities, Food Processing Establishments, and Hospitals. This product cleans, disinfects and deodorizes hard, non-porous inanimate hospital or medical surfaces in one step with no rinsing required. Use daily on the following hard, non-porous inanimate surfaces: vinyl, painted surfaces, plastic surfaces, glazed ceramic, glazed porcelain, laminated surfaces and baked enamel surfaces associated with floors, walls, ceilings, tables, chairs, countertops, telephones, fixtures, glazed tile, toilet exteriors, urinal exteriors, sinks found in health care facilities, hospitals, patient rooms, operating rooms, shower rooms, and locker rooms. It can be used to pre-clean and disinfect hospital items: wheelchairs, patient bedrails and linings, patient bed linings, vinyl mattress covers, wash basins, bed pans, non-critical medical equipment surfaces – almost any washable food and non-food contact surface where disinfection is required. A potable water rinse is required when disinfecting food contact surfaces. Do not use on glasses, dishes and utensils.

When used as directed, this product is highly effective against a wide variety of pathogenic microorganisms including bacteria, antibiotic resistant bacteria, viruses, and fungi.

Using approved AOAC test methods under Good Laboratory Practices, 5% serum load and 1 minute contact time this product kills the following on hard-nonporous inanimate surfaces.

Bactericidal Activity: *Pseudomonas aeruginosa*, (ATCC 15442), *Staphylococcus aureus*, (ATCC 6538), *Salmonella enterica*, (ATCC 10708)

***Viruses:** *Norovirus, (Feline Calicivirus, ATCC VR-782 as the surrogate)

See reference sheet for a complete list of organisms and additional directions for use.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

For Use as a One-Step Cleaner/Disinfectant Product: Pre-clean heavily soiled areas. Apply product by coarse trigger sprayer to hard, non-porous inanimate surfaces. Spray 6 – 8 inches from surface, making sure to wet surfaces thoroughly. All surfaces must remain wet for 1 minute. Use a 5 minute contact time for Tb and a 10 minute contact time for fungi. Wipe surfaces dry, rinse, or allow to air dry.

Note: Rinsing is not necessary unless floors are to be coated with finish or restorer. All food contact surfaces must be rinsed with potable water. Do not use on glassware, utensils, or dishes.

For Use as a Cleaner and/or Deodorizer: Apply product by coarse trigger sprayer to hard, non-porous inanimate surfaces. Wipe surfaces dry, rinse or allow to air dry.

For Use as a Tuberculocide: Pre-clean heavily soiled areas. Apply product by coarse trigger sprayer to hard, non-porous inanimate surfaces. Spray 6 – 8 inches from surface, making sure to wet surfaces thoroughly. Allow the surface to remain wet for 5 minutes. Wipe surfaces dry, rinse, or allow to air dry. Change solution after each use.

To Kill Fungi: Pre-clean heavily soiled areas. Apply product by coarse trigger sprayer to hard, non-porous inanimate surfaces. Spray 6 – 8 inches from surface, making sure to wet surfaces thoroughly. Allow surface to remain wet for 10 minutes. Wipe surfaces dry, rinse or allow to air dry.

To Spot Sanitize Soft Surfaces: Remove gross filth prior to sanitization. Hold bottle upright 6 – 8" from surface. Spray until fabric is wet. DO NOT SATURATE. Allow surface to remain wet for 30 seconds. Gently blot with a clean cloth and let air dry. For difficult odors or heavy fabrics, repeat application. Repeat as needed. Note: Prior to first use on a particular surface, it is recommended to test in a small, discreet area if there is concern of incompatibility.

FOR INSTITUTIONAL USE

Manufactured for: © 2015 Diversey, Inc., PO Box 19747, Charlotte, NC 28219-0747 U.S.A. All Rights Reserved.

***KILLS HIV AND HCV AND HBV ON PRE-CLEANED INANIMATE SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (Hospitals, Nursing Homes) and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV), or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATING AGAINST HIV-1 (AIDS Virus), HBV OR HCV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, and eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood/Body Fluids must be thoroughly cleaned from surfaces/objects before application of this product.

Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1, HBV and HCV. Use a 5 minute contact time for Tb and a 10 minute contact for fungi.

Disposal of Infectious Material: Blood/Body Fluids must be autoclaved and disposed according to Federal, State and local regulations for infectious waste disposal.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Store in original container in areas inaccessible to small children. Keep securely closed.

PESTICIDE STORAGE: Store in a dry, well-ventilated area away from chemicals, direct light, heat or open flame. Do not mix with other cleaning or disinfecting products.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. For more information on the disposal of unused, unwanted product and the clean up of spills, contact the EPA Regional Office of the Manufacturer for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Wrap empty container and put in trash.

ENVIRONMENTAL HAZARDS: This product is toxic to birds, fish and aquatic invertebrates. Caution should be used when applying indoors because pets may be at risk.

IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR DOCTOR FOR TREATMENT ADVICE.

1-800-851-7145

Have the product container or label with you when calling a Poison Control Center or doctor or going in for treatment.

EPA Reg. No. 70627-56
EPA Est. No. 74559-CAN-1 [VX]; 0312-WI-3 [NW]

Lot code letters indicate establishment number.

SDS Ref. # MS0800255

Questions? Comments:

1-800-558-2332

Made In Canada

6003368 (15/203)

Oxivir® Tb



Using approved AOAC test methods under Good Laboratory Practices, 5% serum load and 1 minute contact time this product kills the following on hard-nonporous inanimate surfaces.

Bactericidal Activity:

Staphylococcus aureus, (ATCC 6538)
Salmonella enterica, (ATCC 10708)
formerly known as *Salmonella choleraesuis*
Pseudomonas aeruginosa, (ATCC 15442)

Acinetobacter baumannii, (ATCC 19606)
Escherichia coli O157:H7, (ATCC 35150)
Escherichia coli with extended beta-lactamase
resistance (ESBL), (ATCC BAA-196)

Klebsiella pneumoniae, (ATCC 4352)
Shigella dysenteriae, (ATCC 11835)

Antibiotic-Resistant Bacteria Activity:

Enterococcus faecalis, (ATCC 51575)
(Resistant to Vancomycin [VRE])

Staphylococcus aureus (CA-MRSA),
(NARSA NRS 384)(Genotype USA300)
Community Associated Methicillin Resistant

Staphylococcus aureus (CA-MRSA),
(NARSA NRS 123)(Genotype USA400)
Community Associated Methicillin Resistant

Staphylococcus aureus, (ATCC 33592)
(Resistant to Methicillin [MRSA])

*** Virucidal Activity:**

*Adenovirus type 8, (ATCC VR-1368)

*Herpes Simplex Virus, Type 2 (HSV-2),
(ATCC VR-734)

*Norovirus, (Feline Calicivirus, ATCC VR-782
as the surrogate)

*Hepatitis B Virus (HBV), Duck Hepatitis
B Virus (DHBV) as surrogate
(Acquired from Hepadnavirus Testing)

*HIV-1 (AIDS Virus), Strain HTLV-III_B (HIV-1),
(Acquired from Adv. Biotechnologies, Inc.)

*Poliovirus Type 1, Strain Brunhilde,
(ATCC VR-1000)

*Hepatitis C Virus (HCV), Bovine Viral Diarrhea
Virus (BVDV) as surrogate (Acquired from
National Veterinary Services Laboratories)

*Human Coronavirus, (ATCC VR-740)

*Rhinovirus Type 37, Strain 151-1,
(ATCC VR-1147)

*Herpes Simplex Virus, Type 1 (HSV-1),
(ATCC VR-733)

*Influenza A/Hong Kong, (ATCC VR-544)

*Rotavirus WA, (Acquired from University of
Ottawa)

***Veterinary Viruses:**

*Avian Influenza A (H3N2), (ATCC VR-2072)

*Feline Calicivirus, Strain F9, (ATCC VR-782)

TUBERCULOCIDAL in the presence of a 5% serum load and 5 minute contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

Mycobacterium bovis (Tb) (OT 451C150)

FUNGICIDAL: This product is fungicidal against the pathogenic fungi *Trichophyton mentagrophytes* (ATCC 9533) (the fungus which causes Athlete's Foot) in the presence of a 5% serum load and 10 minute contact time when used as directed on hard surfaces found in bathrooms, shower stalls, locker rooms, or other clean, non-porous, hard surfaces commonly contacted by bare feet.

Soft Surface Spot Sanitizer in the presence of 5% soil with a 30 second contact time on cotton or polyester fabrics against the following bacteria:

Staphylococcus aureus, (ATCC 6538)

Acinetobacter baumannii, (ATCC 19606)

Klebsiella pneumoniae, *Enterobacter aerogenes*, (ATCC 13048)

(Multi-Drug Resistant, Resistant to Cefazolin, Gentamicin, Trimethoprim/Sulfa, and Ertapenem)
Enterococcus faecalis, (ATCC 51575) (ATCC BAA-1705) (Carbapenem-Resistant)
(Resistant to Vancomycin [VRE])

Staphylococcus aureus, (ATCC 33592) (Resistant to Methicillin [MRSA])

Non-Food Contact Surface Sanitizer for hard non-porous surfaces in the presence of a 5% serum load and 30 second contact time on hard, non-porous environmental surfaces.

Enterococcus faecalis, (ATCC 51575)

Pseudomonas aeruginosa, (ATCC 15442)

Staphylococcus aureus, (ATCC 6538)

(Resistant to Vancomycin [VRE])

Escherichia coli O157:H7, (ATCC 35150)

Salmonella enterica, (ATCC 10708)

Staphylococcus aureus, (ATCC 33592)
(Resistant to Methicillin [MRSA])

Klebsiella pneumoniae, (ATCC 4352)

Malodor Counteractancy – eliminates odors and odor-causing bacteria on hard, nonporous surfaces in restroom areas, behind and under sinks and counters, storage areas, and other places where bacterial growth can cause malodors.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For use as a Non-Food Contact Sanitizer: Pre-clean heavily soiled areas. Apply Solution with spray, cloth, disposable wipe or mop to hard, non-porous environmental surfaces. Allow the surface to remain wet for 30 seconds. Wipe surfaces dry, rinse or allow to air dry. Change solution after each use application.

Animal Housing Facilities Including Veterinary Clinics, Farms, Livestock, Swine and Poultry Houses: For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities (including pens, chutes, and barns) occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding or watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water. Saturate surfaces by spraying with product for a period of 1 minute. Use a 5 minute contact time for TB and a 10 minute contact time for fungi.
5. Spray handling and restraining equipment such as leashes, muzzles, halters or ropes, as well as forks, shovels, and scrapers used for removing litter and manure.
6. Ventilate buildings and other closed spaces.
7. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub treated feed racks, troughs, mangers, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

Read the label affixed to the container for this product before applying.

Use of this product according to this label is subject to the use precautions and limitations imposed by the label affixed to the container for this product.

EPA Reg. No. 70627-56

SDS # MS0800255

Manufactured for: ©2015 Diversey, Inc., PO Box 19747, Charlotte, NC 28219-0747 U.S.A. All Rights Reserved