



LICENSED

PERIOD 2012-2014 LIC. NO.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2015-2017 LIC. NO.

8053.8



Powered by **OdoBan**®

Germ Control

Effective against Canine Parvovirus*



Active Ingredients:

Alkyl (C ₁₄ 60%, C ₁₆ 30%, C ₁₂ 5%, C ₁₈ 5%)	
dimethyl benzyl ammonium chloride	4.5%
Alkyl (C ₁₂ 68%, C ₁₄ 32%)	
dimethyl ethylbenzyl ammonium chloride	4.5%
Inert Ingredients	91.0%
Total	100.0%

EPA Reg. No. 10324-105-66243
EPA Est. No. 66243-GA-001

Net Contents:

1 Gallon (3.79 L)

KEEP OUT OF REACH OF CHILDREN
DANGER. See side panel for additional precautionary statements and first aid statements.

Cleans by removing dirt, grime, mold, mildew, blood, urine, fecal matter and other common soils found in animal housing facilities, livestock, swine poultry facilities, grooming facilities, farms, kennels, pet stores, veterinary clinics, laboratories or other small animal facilities. It also eliminates odors leaving surfaces smelling clean and fresh.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Apply this product with a cloth, mop or mechanical spray device. When applied with a mechanical spray device, surfaces must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. For sprayer applications, use a coarse pump or trigger sprayer. Spray 6-8 inches from surface. Rub with brush, sponge or cloth. Do not breathe spray. **NOTE:** With spray applications, cover or remove all food products. Prepare fresh solution at least daily or when use solution becomes visibly dirty. For heavily soiled areas, a preliminary cleaning is required.

DISINFECTION PERFORMANCE IN NON-MEDICAL FACILITIES:

At 1 ounce of this product to one gallon of water (703 ppm active quat) use level, this product is bactericidal according to the AOAC Use Dilution Test method on hard inanimate surfaces modified in the presence of 5% organic serum with a 10-minute contact time against: *Bordetella bronchiseptica*, *Campylobacter jejuni*, *Corynebacterium ammoniagenes*, *Enterobacter aerogenes*, *Enterococcus faecalis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Listeria monocytogenes*, *Salmonella enterica*, *Salmonella schottmuelleri*, *Salmonella typhi*, *Shigella dysenteriae*, *Staphylococcus aureus*, Community associated Methicillin resistant *Staphylococcus aureus* (CA MRSA), *Staphylococcus aureus* Methicillin resistant, *Streptococcus salivarius*

***ANIMAL VIRUCIDAL PERFORMANCE:** At 1 ounce per gallon of water use level, this product was evaluated in the presence of 5% serum with a 10-minute contact time and found to be effective against the following viruses on hard, non-porous, environmental surfaces: Avian Influenza A H5N1 virus, Avian influenza A H9N2/Turkey/Wisconsin virus, Avian Reovirus, Bovine Viral Diarrhea virus, Canine Distemper virus, Canine Coronavirus, Equine Arteritis Virus, Infectious Avian Laryngotracheitis virus, Infectious Bovine Rhinotracheitis virus (IBR), Infectious Bronchitis Virus, Newcastle disease virus, Porcine Respiratory & Reproductive Virus (PRRSV), Porcine Rotavirus, Pseudorabies virus, Transmissible Gastroenteritis virus (TGE), Vaccinia virus

At 1.25 ounces per gallon of water use level, this product is found to be effective against the following viruses on hard, non-porous environmental surfaces in the presence of 5% serum with a 10-minute contact time: **Canine Adenovirus, Canine**

Parainfluenza Virus, Feline Calicivirus, Feline Infectious Peritonitis virus, Feline leukemia virus, Feline Panleukopenia virus, Feline Picornavirus, Feline Rhinotracheitis virus, Infectious Canine Hepatitis Virus, Murine Parainfluenza virus type 1, Rabies virus

At 4 ounces per gallon of water use level, this product was evaluated in the presence of 5% serum with a 10-minute contact time and found to be effective against the following viruses on hard, non-porous environmental surfaces: **Canine Parvovirus (CPV), Minute virus of Mice Parvovirus, Porcine Parvovirus**

***FOR DISINFECTING HARD, NON-POROUS SURFACES AGAINST AVIAN INFLUENZA A VIRUS:** For use in animal housing facilities, follow the animal premise directions.

***FOR CANINE PARVOVIRUS ACTIVITY:** This product is effective against Canine Parvovirus at 4 oz. per gallon. Follow "Veterinary Clinics/Animal Life Science Laboratory/Animal Care Facilities/Animal Research Centers/Animal Quarantine Areas/Zoos/Pet Shop/Kennels/Animal Breeding Facilities/Breeding and Grooming Establishments/Tack Shops Disinfection Directions".

VETERINARY CLINICS/ANIMAL LIFE SCIENCE LABORATORY/ANIMAL CARE FACILITIES/ANIMAL RESEARCH CENTERS/ANIMAL QUARANTINE AREAS/ZOOS/PET SHOP/KENNELS/ANIMAL BREEDING FACILITIES/ANIMAL HUSBANDRY/BREEDING AND GROOMING ESTABLISHMENTS/TACK SHOPS

DISINFECTION DIRECTIONS: For cleaning and disinfecting the following hard non-porous surfaces: equipment, utensils, instruments, cages, kennels, stables, stalls and catteries. Remove all animals and feeds from premises, animal transportation vehicles, crates etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of the disinfecting and virucidal solution (1 oz. per gallon of water) for a period of 10 minutes. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals as well as forks, shovels, scrapers used in removing litter and manure. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, automatic feeders, waterers and other equipment which dispenses food or water with soap or detergent, and rinse with potable water before reuse.

***SPECIAL INSTRUCTIONS FOR INACTIVATING AVIAN INFLUENZA A VIRUS:** Remove all poultry and feeds from premises, trucks, coops, and crates. Remove all litter and droppings from floors, walls, and surfaces of facilities occupied or traversed by poultry. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with the disinfecting solution (1 oz per gallon of water) for a period of 10

minutes. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels, scrapers used in removing litter and manure. Ventilate buildings, coops, and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set, or dried. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

FUNGICIDAL PERFORMANCE: At 1 ounce per gallon use-level, this product is effective against Trichophyton mentagrophytes-athlete's foot fungus/a cause of Ringworm and Aspergillus niger on inanimate surfaces in the presence of 5% blood serum in locker rooms, dressing rooms, shower and bath areas and exercise facilities. Saturate surfaces with use solution for a period of 10 minutes. Allow solution to dry on surface and repeat treatment every seven days or when new growth appears.

MOLD AND MILDEW CONTROL: At 1 ounce per gallon (or equivalent use dilution) (703 ppm active quat) this product will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard non-porous surfaces. Saturate surfaces with use solution for a period of 10 minutes. Allow solution to dry on surface and repeat treatment every seven days or when new growth appears.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. PESTICIDE STORAGE: Open dumping is prohibited. Store only in original container. Do not reuse empty container. If a leaky container must be contained within another, mark the outer container to identify the contents. Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep this product under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use. **PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Non-refillable container. Do not reuse this container to hold materials other than pesticides or rinsate. Offer for recycling if available or puncture and dispose in a sanitary landfill, or by other procedures approved by state and local authorities in accordance with local, state and federal regulations. **RESIDUE REMOVAL INSTRUCTIONS:** Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Fill the container ¼ full with water and recap. Shake for 10 seconds. Follow Pesticide Disposal instructions for rinsate disposal. Drain for 10 seconds after the flow begins to drip. Repeat procedure two more times.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER. KEEP OUT OF REACH OF CHILDREN. Corrosive. Causes irreversible eye damage and skin burns. Harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves and protective clothing when handling. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

PHYSICAL OR CHEMICAL HAZARDS: Do not use or store near heat or open flame. Do not mix with oxidizers, anionic soaps and detergents.

FIRST AID: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.



Pets Rule Division • PO Box 7444
Warner Robins, GA 31095
odobanpetsrule.com

13771-G 300212 120504

