




LICENSED

PERIOD 2015-2017 LIC. NO.

8054.10



Andersons™
GOLF PRODUCTS

LICENSED
HAWAII
DEPT. OF AGRICULTURE
Period ~~2003-2005~~ No. 8054.10

Regis



LICENSED

PERIOD 2006-2008 LIC. NO.



LICENSED

PERIOD 2012-2014 LIC. NO.



LICENSED

PERIOD 2009-2011 LIC. NO.

Goosegrass/ Crabgrass Control

- **Prevents goosegrass and crabgrass in bermudagrass, zoysiagrass, tall fescue, bentgrass, perennial bluegrass, and perennial ryegrass**
- **For use on bentgrass fairways (in listed states) and tees**
- **For use on bentgrass and bermudagrass greens**

KEEP OUT OF REACH OF CHILDREN
WARNING

See back panel for additional precautionary statements

ACTIVE INGREDIENTS:

Bensulide†	5.25%
Oxadiazon††	1.31%
OTHER INGREDIENTS:	<u>93.44%</u>
Total	100.00%

†CAS# 741-58-2
††CAS# 19666-30-6

EPA Reg. No. 9198-176

EPA Est. Nos. 538-OH-1S, 9198-OH-1M
Underlined letter used is first letter of run code on end of bag.

SGN 80

Treats Up To 22,000 Square Feet

Recommended For Use By Professional Turfgrass Managers

Net Wt 28.87 Lb (13.1 kg)



Goosegrass/Crabgrass Control

AGC8656-1

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: **WARNING:** causes substantial but temporary eye injury. Do not get in eyes, or on clothing. Wear goggles or face shield. Harmful if swallowed. Wash thoroughly with soap and water after reuse.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance selection chart.

Applicators using a push type spreader and all loaders must wear:

- Coveralls over long-sleeved shirt and long pants,
- Chemical resistant gloves such as barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils, Viton \geq 14 mils,
- Chemical resistant footwear plus socks, and
- A NIOSH-approved dust mist filtering respirator with MSA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator with any N₂, R, P, or HE filter.

All other applicators and handlers must wear:

- Long-sleeved shirt and long pants,
- Chemical resistant gloves such as barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils, Viton \geq 14 mils, and
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides CFR (d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

If swallowed: Call a poison control center immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

For best results apply to DRY FOLIAGE and ensure that the product granules reach the soil surface by removing debris and mowing if necessary prior to treatment.

Apply uniformly at the NORMAL RATE (1X) in the spring (South: January to early April; North: March to early May) prior to crabgrass or goosegrass germination.

ADDITIONAL SEEDING, OVERSEEDING, SPRIGGING, AND STOLONIZING INFORMATION
Do not use on newly seeded bermudagrass until after the second mowing.

Aerification, coring or other practices which break the herbicidal barrier will increase germination if performed just before seeding.

Residual effectiveness requires a delay of 5 growing months after the last treatment before overseeding. On newly sprigged bermudagrass wait until the grass has filled in and the stolons have rooted down before application.

Where large areas of bermudagrass have been severely thinned (25% or more), withhold application until desired fill-in and rooting of the stolons are achieved.

SPREADER APPLICATION RECOMMENDATIONS

Correct spreader calibration is important to minimize undesirable results.

For best results on bentgrass greens, application using a properly calibrated drop spreader is recommended.

When applying the product at the NORMAL RATE (1X), a HALF WIDTH application can be made to ensure a more uniform coverage and particle distribution.

Heavy Weed Pressure

Where heavy weed pressure is expected, apply a NORMAL RATE (1X) followed by a repeat application at the HALF RATE (1/2 X), 6 to 8 weeks later. DO NOT APPLY THIS REPEAT APPLICATION TO BENTGRASS GREENS OR FAIRWAYS.

Reduce Yellowing or Injury

To reduce the potential for yellowing or injury on established bentgrass greens, bermudagrass greens, and perennial ryegrass winter overseedings, apply an initial treatment to dry foliage at the HALF RATE (1/2 X). Apply a repeat treatment 10-14 days later at the HALF RATE (1/2 X).

Where herbicidal effects at the NORMAL RATE (1X) can be tolerated, this product may be applied at the HALF WIDTH to ensure proper particle coverage.

Poa trivialis

When using this product on established winter overseedings of Poa trivialis or blends containing Poa trivialis, apply an initial treatment to dry foliage at the HALF RATE (1/2 X). Repeat

Do not give anything by mouth to an unconscious person.

If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling poison control center or doctor, or going for treatment.

Environmental Hazards:

This pesticide is toxic to fish and aquatic vertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Do not apply when weather conditions favor drift from the treated area. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This product may impair reproduction in birds if used during the breeding season.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Controls crabgrass and goosegrass as they sprout.

This product may only be applied by tractor-drawn spreader or by push type spreader.

For use only on golf courses (greens, tees and bentgrass fairways only) and on residential lawns.

Applications on bentgrass fairways is limited to the following states: OH, PA, NY, MI, CT, MA, IN, IL, NJ, WV, MN, WI, VT, NH, RI, DE, MD, VA. Applications on bentgrass fairways may only be made in the fall and are limited to one application per year.

Applications to greens and tees are limited to two per year.

PRECAUTIONS

IMPORTANT: THIS PRODUCT WILL NOT WORK UNLESS IT IS WATERED IN FOR 10-15 MINUTES. FOR SAFETY REASONS, WATER THIS PRODUCT IN AS SOON AS POSSIBLE FOLLOWING APPLICATION FOR 10-15 MINUTES AND DO NOT ALLOW CHILDREN OR PETS ON TREATED AREAS UNTIL DRY FOLLOWING THE WATERING.

WATERING-IN MUST BE PERFORMED BY THE COMMERCIAL APPLICATOR OR THE

Poa trivialis

When using this product on established winter overseedings of Poa trivialis or blends containing Poa trivialis, apply an initial treatment to dry foliage at the HALF RATE (1/2 X). Repeat treatment again 10 - 14 days later at the HALF RATE (1/2 X).

STORAGE AND DISPOSAL

STORAGE: Store in a cool, clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION. (RVnot030102)

**Visit us at our website at:
andersonsgolfproducts.com**

Manufactured for:
The Andersons Lawn Fertilizer Division, Inc.
P. O. Box 119

PRECAUTIONS

IMPORTANT: THIS PRODUCT WILL NOT WORK UNLESS IT IS WATERED IN FOR 10-15 MINUTES. FOR SAFETY REASONS, WATER THIS PRODUCT IN AS SOON AS POSSIBLE FOLLOWING APPLICATION FOR 10-15 MINUTES AND DO NOT ALLOW CHILDREN OR PETS ON TREATED AREAS UNTIL DRY FOLLOWING THE WATERING.

WATERING-IN MUST BE PERFORMED BY THE COMMERCIAL APPLICATOR OR THE COMMERCIAL APPLICATOR MUST PROVIDE THE ABOVE WATERING-IN INSTRUCTIONS TO THE RESIDENT IN WRITING USING BOLD FACE TYPE AND/OR RED PRINT OR OTHER METHODS TO ENSURE WARNING IS PROMINENTLY DISPLAYED ON THE MATERIAL PROVIDED TO THE RESIDENT.

Do not apply this product in a way that will contact workers or other persons.


Do not use on parks, recreation areas, or other public sites.

Visit us at our website at:
andersonsgolfproducts.com

Manufactured for:
 The Andersons Lawn Fertilizer Division, Inc.
 P. O. Box 119
 Maumee, Ohio 43537

AGC8656-1

©2002 T.A.I.
 Andersons Golf Products is a trademark of
 The Andersons AgServices, Inc.



ESTABLISHED COOL AND WARM-SEASON GRASSES
 This product may be used on bentgrass and bermudagrass putting greens only under conditions of very heavy goosegrass infestations and where the herbicide side effects are tolerable.

This product may cause yellowing and/or thinning for up to 4 weeks on bentgrass and bermudagrass greens.

Do not use this product when the temperature is consistently below 50°F or above 80°F.

Not for use where Poa annua, annual ryegrass, or fine fescue are the desired turf or on dichondra or similar groundcovers.

Complete any raking or aerifying of the turf before treatment so that the protective chemical barrier is not destroyed.

SUGGESTED SPREADER SETTINGS

SPREADER	GROUND SPEED	NORMAL RATE (1X)				HALF RATE (1/2 X)	
		SETTING	FULL WIDTH OF COVERAGE	SETTING	HALF WIDTH OF COVERAGE	SETTING	FULL WIDTH OF COVERAGE
Andersons SS-2 Drop	3 mph	3	overlap wheels	-	-	2 1/4	overlap wheels
Andersons Rotarys: AccuPro/SR-2000 (Cone fit)	3 mph	H	9 feet	G	4.5 feet	G	9 feet
Scotts Rotarys: R-8, R-8A (Cone fit) R-7X	3 mph 3 mph	H G 1/2	9 feet 9 feet	F 1/2 F 1/4	4.5 feet 4.5 feet	F 1/2 F 1/4	9 feet 9 feet
Vicon (Q3 Series-Standard Spout)	4.5 mph	14	18 feet	9	9 feet	9	18 feet
Teley Models	4.5 mph	3 1/4 III	18 feet	2 1/2 III	9 feet	2 1/2 III	18 feet
APPLICATION INFORMATION							
lb. active ingredient/ acre (Bifenox, Oxadiazon)		5.0, 1.5		5.0, 1.5		3.0, 0.75	
lb. product/1,000 sq. ft.		2.52		2.52		1.31	
Number of Bags/Acre		4		4		2	

Attention: This product contains a chemical known to the state of California to cause cancer, birth defects or reproductive harm.