



0.48% Barricade® on DG Pro™

For preemergence control of grass and broadleaf weeds in established turfgrasses (excluding golf course putting greens), lawns, sod nurseries; landscape ornamentals; established perennial and wildflower plantings; non-crop areas on or surrounding managed rights-of-way for transportation systems and utilities (including roadways, roadsides, railways and equipment yards); facilities including substations, tankfarms, pump- ing stations, parking and storage areas, and ungrazed fencerows.

ACTIVE INGREDIENT:
 Prodiamine™ 0.48%

OTHER INGREDIENTS: 99.52%

Total 100.00%

*(CAS No. 29091-21-2)
 EPA Reg. No. 9198-214

EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A

Barricade is a registered trademark of Syngenta Crop Protection, Inc.
 The Andersons logo is a registered trademark of The Andersons Agriservices, Inc.: DGPro and Professional Turf are trademarks of The Andersons Agriservices, Inc.

KEEP OUT OF REACH OF CHILDREN

CAUTION PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE)

- WPS USES:**
 Applicators and other handlers:
- Long sleeved shirt and long pants
 - Waterproof gloves
 - Shoes plus socks

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash all PPE separately from other laundry.

NON-WPS USES:

Mixers and loaders who handles this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) – in general, only agricultural-plant uses are covered by the WPS – must wear:

- Chemical-resistant gloves

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash all PPE separately from other laundry.

Engineering Controls:

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

- Users should:
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - After handling this product, immediately wash the outside of gloves before removing them, then remove gloves and all other PPE. Immediately wash thoroughly and change into clean clothing.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

Environmental Hazards

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations substantially above the level of water solubility, it may be toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

If material is spilled, contain and sweep up material of spill for disposal or reuse per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under or cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Workers Protection Standard 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to use of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The Worker Protection Standard applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dusts have settled.

GENERAL INFORMATION

This product is a preemergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses (excluding golf course putting greens), lawns, sod nurseries, landscape ornamentals, established perennial & wildflower plantings, non-crop areas on or surrounding managed rights-of-way for transportation systems and utilities (including roadways, roadsides, railways & equipment yards), facilities including substations, tankfarms, pumping stations, parking & storage areas, ungrazed fencerows.

This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Do not graze or feed livestock forage cut from areas treated with this product. Do not apply this product through any type of irrigation system unless otherwise instructed by this label. Do not apply aerially. Do not apply to golf course putting greens.


APPLICATION DIRECTIONS

Apply uniformly through suitable, calibrated application equipment.

ESTABLISHED TURF

This product is a preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the Maximum Annual Rates section of this label.

Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.



- Approx. Date:**
- 1 - After May 30
 - 2 - After May 10
 - 3 - After April 20
 - 4 - After March 20
 - 5 - Jan. 1 to March 20

Rates of Application

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

Maximum Annual Rates

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Application Rate of this product per Calendar Year by Turf Grass Species ¹			
Turf Species	Lbs. AI/A	Lbs. product /A	Lbs. product/ 1,000 sq. ft.
Bermudagrass ² , Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including turf-type), Zoysia	1.5	312	7.1
Buffalograss, Kentucky Bluegrass, Perennial Ryegrass	1.0	208	4.7
Creeping Red Fescue	0.75	156	3.5
Creeping Bentgrass	0.65	135	3.1

¹These are the maximum rates per calendar year by species limitations.

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 108 lb./acre or 2.4 LB./1,000 sq. ft. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

Use higher rates of this product to achieve higher levels weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates table.

Weeds Controlled

When used at the recommended rates this product will control:

Barnyardgrass	Goosegrass ¹	Pigweed
Bluegrass, Annual	Henbit	Purslane, Common
(<i>Poa annua</i>)	Itchgrass	Pusley, Florida
Carpetweed	Johnsongrass	Rescuegrass ³
Chickweed, Common	(from seed)	Shepherd's Purse ²
Chickweed, Mouseear	Junglerice	Signalgrass, Broadleaf
(from seed)	Knotweed	Speedwell, Persian
Crabgrass	Kochia	Sprangletop
(Large, Smooth)	Lambsquarter, Common	Spurge, Prostrate
Crowfootgrass	Lovegrass	Witchgrass
Cupgrass, Woolly	Oxalis	Woodsorrel, Yellow
Foxtails, Annual	Panicum	(from seed)
	(Texas, Fall, Browntop)	

¹In many areas a single application of 3.1 to 7.1 lbs./1,000 sq.ft. (135-315 lbs./A) of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 3.1 to 4.7 lbs./1,000 sq. ft. (135-208 lbs./A) followed after 60 to 90 days by a application at doses that would not exceed those given in the Maximum Application Rates Table. Do not exceed the maximum rate for turf grass species listed in the Maximum Application Rates Table above.

²Applications for this weed should be made in late summer, fall, or winter prior to germination.

³Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

When to Apply After Overseeding Turf

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

When to Overseed After Application - All States*

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

The Andersons Professional Turf Products 0.48% Barricade on DG Pro			Interval Before Overseeding		
Lbs/1,000 sq. ft.	Lbs./A	AI/A	North	Transition	South
2.3	104	0.50	4 mo	4 mo	4 mo
3.1	135	0.66	5 mo	4 mo	4 mo
3.5	156	0.74	6 mo	5 mo	5 mo
3.8	166	0.80	---	6 mo	6 mo
4.7	208	1.00	---	7 mo	7 mo
5.4	237	1.14	---	---	9 mo
6.1	270	1.30	---	---	10 mo
7.1	312	1.50	---	---	12 mo

*Note: In AZ, CA, NV and TX, the overseeding interval can be shorter in established bermudagrass that has been overseeded with perennial ryegrass. See the next section "*Poa annua* Control in Established Bermudagrass Overseeded with Perennial Ryegrass (AZ, CA, NV and TX only)".

Poa annua Control in Established Bermudagrass Overseeded with Perennial Ryegrass (AZ, CA, NV and TX only)

Use on golf courses (excluding golf course putting greens) and lawns when overseeding with perennial ryegrass (minimum seed rate of 350 lbs./A).

Amount to Apply
 Lbs./Acre Lbs/1,000
 78 – 135 1.8 – 3.1

When to Apply
 6 to 8 weeks before ryegrass overseeding

Second application 4-8 weeks after overseeding date or when perennial ryegrass roots are in the second inch of soil

How Much and When to Apply

Expected Control
 One application for 70% or greater control of *Poa annua*

Second application should enhance the residual under normal weather conditions

Use Precautions

1. Some seed mortality and temporary reduction in root growth of new seedlings may occur.
2. To reduce the potential for seeding mortality, maintain a moist seedbed with light, frequent irrigation.
3. Make no more than 2 applications per year for this use, and do not exceed 4 lbs./1,000 sq. ft. (176 lbs./A) for these two applications for *Poa annua* control in ryegrass overseeded turf.
4. Do not make a second application if any injury to the ryegrass is observed after the first application.
5. Do not make a second application unless the product was first applied before overseeding.



LICENSED

PERIOD 2007-2009 LIC. NO.

Use Precautions

The following precautions apply to the use of this product in turf grasses and lawns:

Application of this product may thin emerged annual bluegrass and newly over seeded grasses. Do not apply to over seeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil. Do not apply to newly set sod until the following year. Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control. Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bluegrass (*Poa annua*) are desirable species.

SPREADER SETTINGS

Recommended Spreader Settings					
These suggested spreader settings are not intended to replace calibration. Calibrate your spreader before applying product.					
50 lbs. treats 16,100 sq. ft. (0.37 acre) at the Low Rate (3.1 lbs./1,000 Sq. Ft.)					
50 lbs. treats 14,200 sq. ft. (0.33 acre) at the Medium Rate (3.5 lbs./1,000 Sq. Ft.)					
50 lbs. treats 10,600 sq. ft. (0.24 acre) at the High Rate (4.7 lbs./1,000 Sq. Ft.)					
SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS		
			LOW RATE	MEDIUM RATE	HIGH RATE
Andersons AccuPro 1000, 2000, SR-2000 (cone 6)	3 mph	11 feet	K 1/2	L 1/2	M 1/2
Andersons SS-2 Drop	3 mph	overlap wheels	4 3/4	5	5 1/2
Scotts Rotaries: R-8, R-8A (cone 6)	3 mph	11 feet	J 1/2	K	L 3/4
LESCO	3 mph	5 feet	D 1/2	E	E 3/4
Lely Models WTR, WFT, HR, 1250 (PTO at 450 rpm)	4.5 mph	34 feet	5 IIC	5 1/4 IIC	6 1/4 IIC
Prizelawn CB85 B	3 mph	5 feet	3 1/2	3 3/4	4 1/4
Vicon (03 series)	4.5 mph	20 feet	17	18	22

LANDSCAPE ORNAMENTALS

This product may be applied for residual preemergence weed control in ornamentals (including established perennial and wildflower plantings).

Use Rates

Apply this product at 3.1 to 7.1 lbs./1,000 sq. ft. (135 to 312 lbs./A) in fall and/or spring. Use higher rate of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 7.1 lbs./1,000 sq. ft. (312 lbs./acre).

Application Timing and Information

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves and mulch. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

Apply this product to dry foliage only. Overhead irrigation or rainfall soon after application will wash granules off plant foliage and activate this product in the soil.

Tolerant Ornamental Species

This product will not harm most trees, shrubs, vines, and flowers. The species listed on back panel are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface. This product may be applied over the top of these species.

When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

LICENSED

PERIOD 2013-2015 LIC. NO.

8054.24 Disc.

LICENSED

PERIOD 2010-2012 LIC. NO.

G4 (BLKP1929B) Front

APTDG48BR5.1

Fir species** (Balsam, Fraser, Noble, etc.)

Abelia: Sherwood
Japanese Maple
Norway Maple
Yarrow:King Edward
Kiwi*
Lily-of-the-Nile (African Lily)
Five-Leaf or Chocolate Vine
Lady's Leek, Nodding Onion
Japanese Anemone
Aquilegia: Red and gold
Vine Hill Manzanita
Cape Weed
Wormwood: Silver Mound, Castle
Japanese Aucuba
Barberry
Aster: Bonny Blue, Purple Dome
Lady Fern; Fern Lady
Fibrous Begonia; Hardy Grandis
Barberry
Wintergreen Barberry
Mentor Barberry
Japanese Barberry
Warty Barberry
Snowbank
Bougainvillea
Butterfly-Bush (Dwarf Blue); Royal Red
Japanese Boxwood
Crimson Bottlebrush
Weeping Bottlebrush
Scotch Heather
Tussock Bellflower: (White Clips)
Trumpet Creeper, Trumpet Flower; Madame Galen
Hottentot Fig (Ice Plant)
Feathery Cassia
Wild Lilac
False Cypress
Cleyera
Citrus species*
Coreopsis (Calliopsis): Early Sunrise
Moonbean
Flowering Dogwood
American Dogwood
Pampas Grass
Cranberry Cotoneaster
Cotoneaster
Bearberry Cotoneaster
Rockspray Cotoneaster
Hawthorn
Lucifer
Italian Cypress
White Trailing Ice Plant
Cooperi Pink
Larkspur; Blue Elf
Dianthus, Maiden Pinks 'Zing'
Cheddar Pink
Hop Bush
Coneflower, Purple; Magnus
Silverberry
Wintercreeper
Japanese Spindle Tree (Evergreen Euonymus)
Spreading Euonymus
Japanese Aralia
Border Forsythia
Weeping Forsythia
Greenstem Forsythia
Galliaria, Blanket Flower: 'Goblin'
Gardenia, Cape-Jasmine; Radicans
Gentian
Cranesbill
Gladiolus species**
Baby's Breath
English Ivy
Sunrose
Daylily: Aztec Gold, Stella De Oro, Tender Love
Coral Bell; Bridget Bloom
Rose of Sharon**
Chinese Hibiscus
Mallow; Disco Belle White
Hosta, Plantain Lily (Fragrant)
Hosta, 'Searsucker'
Bigleaf Hydrangea
Chinese Holly
Japanese Holly; Helleri
American Holly
Holly
Yaupon Holly; Schillings
Sword-Leaved Iris; Jodlesong
Siberian Iris; Cabernet
Iris species**
Winter Jasmine
Chinese Juniper; Nick's Compact, Parsonii
Shore Juniper; Blue Pacific
Parsoni
Creeping Juniper
Walnut*
Shrimp Plant
Crape Myrtle
Crape Myrtle; Tuscarora
Weeping Lantana
Lavender; Munstead
Edleweiss
Chinese Privet; Variiegata
Amur Privet
Japanese Privet
Glossy Privet (Wax-Leaf)

Lily: Jazz
Big Blue Lillyturf; Turf, Silver Mound, Evergreen
Giant
Liriope, Variegated
Liriope, Creeping
Cardinal Flower, Indian Pink
Japanese Honeysuckle
Tatarian Honeysuckle
Burgundy
Loosestrife; Modern Pink
Magnolia species**
Ice Plant
Crabapple*
Yaku Jima, Silberfeder
Heavenly Bamboo
Narcissus species**
Oleander
Evening Primrose
Olive*
Mondo Grass
Osmanthus (False Holly); Gulf Tide
Trailing African Daisy
Sourwood
Tree Peony
Fountain Grass (Dwarf)
Avocado*
Frasier's Photinia (Redtip)
Dragonhead; False; Vivid
Spruce species** (Colorado Blue, Norway, etc.)
Lily-of-the-Valley Shrub
Calabrian Pine
Canary Island Pine
Slash Pine
Aleppo Pine
Austrian Black Pine
Longleaf Pine
Monterey Pine
Japanese Black Pine
Eastern White Pine
Scotch Pine
Loblolly Pine
Virginia Pine
Pistachio*
Queensland Pittosporum
Japanese Pittosporum
Japanese Yew
English Laurel
Almond, Apricot, Nectarine,
Peach, Plum and Prune*
Douglas Fir**
Firethorn Scarlet
Firethorn
Firethorn
Bradford Pear spp.
Oak, Shumard's Red
Oak species
Indian Hawthorn
Yedda Hawthorn
Rhododendron (including Azalea)
'Coral Bells'
'Delaware Valley White'
'Flame Creeper'
'Formosa'
'Girard Crimson'
'George I. Tabor'
'Hino-crimson'
'PJM'
'Roseum Elegans'
Wakeiebisu
White Gumbo
Lady Bank's Rose
Rosemary
Black-Eyed Susan; Goldstrum
Leatherleaf Fern
Santolina
Saxifrage; Purple Dome
Pincushion Flower
Stonecrop; Lidakense, Dragon's Blood
Spirea; Anthony Waterer
Australian Brushcherry
Japanese Yew
Yew
Germander
Meadow Rue
American Arborvitae
Star Jasmine
Canada Hemlock
Tulip species
Veronica Speedwell; Sunny Border
Japanese Viburnum
Sweet Viburnum
Japanese Snowball
Canary Island Viburnum
Arrowood Viburnum
Laurustinus
Cranberry Bush
Leatherleaf Viburnum
Greater Periwinkle, Vinca
Common Periwinkle, Vinca
Grape*
Old Fashioned Weigela
Spanish Bayonet, Yucca
Adam's Needle, Yucca

*Do not use on food producing trees, vines, or plants.

**Not for use on container grown plants.

VEGETATION MANAGEMENT

This product may be applied to soil surfaces for preemergence control of many grass and broadleaf weeds in non-cropland areas on or surrounding:

- Non-crop areas on or surrounding managed rights-of-way for transportation systems and utilities (including roadways, roadsides, railways & equipment yards)
- Facilities including substations, tankfarms, pumping stations, parking and storage areas and ungrazed fencerows

Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Apply this product at 3.1 to 7.1 lbs./1,000 (135 to 312 lbs./A) in fall and/or spring. Use higher rate of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 7.1 lbs./1,000 (312 lbs./A).

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in original container away from fertilizer, feed, or food stuffs and separated from other pesticides.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty container into application equipment. Then dispose of empty bag or box in a sanitary landfill or at an incineration facility; or, if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

**Manufactured by:
The Andersons
Lawn Fertilizer Division, Inc.
P. O. Box 119, Maumee, OH 43537**

**Net. Wt. 50 Lbs. (22.68 kg)
Treats Up To 16,100 Sq. Ft.**

G4 (BLKP1929B) Back

APTDG48BR5.1