



STATE OF HAWAII  
Department of Agriculture  
**ACCEPTED**  
LICENSE NO. **8054.49**

*Fortify*

# Phosphorus Free Winterizer

Plus Weed Control

**18-0-12**

APPLY IN THE FALL

*Fortify*

Phosphorus Free  
**Winterizer**  
Plus Weed Control

**18-0-12**

**18-0-12**

Phosphorus Free  
**Winterizer**  
Plus Weed Control

*Fortify*  
APPLY IN THE FALL


**APPLY IN THE FALL**



- Contains Dissolve® for control of over **250** broadleaf, creeping & woody weeds
- Extended greening from controlled release nitrogen

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**  
See back panel for First Aid and Additional Precautionary Statements  
EPA Reg. No. 228-281-9198  
EPA Est. Nos. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A  
(Underlined letter used is first letter of run code on end of bag.)

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300.  
For Medical Emergencies Only, Call (877) 325-1840




Net Weight  
**18 lbs.**  
(8.16 kg)  
Covers 5,000 Square Feet

<b>ACTIVE INGREDIENTS:</b>	
Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	0.580%
Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**†	0.145%
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy)propionic Acid***†	0.143%
<b>OTHER INGREDIENTS:</b>	<b>99.152%</b>
<b>TOTAL</b>	<b>100.000%</b>
<b>ISOMER SPECIFIC METHOD, EQUIVALENT TO:</b>	
* 2,4-Dichlorophenoxyacetic Acid	0.46%
** (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	0.12%
*** (+)-R-2-(2,4-Dichlorophenoxy)propionic Acid	0.12%
†CONTAINS THE SINGLE ISOMER FORMS OF MECOPROP-p AND DICHLORPROP-p	

*Fortify*  
APPLY IN THE FALL

Phosphorus Free  
**Winterizer** **18-0-12**  
Plus Weed Control





# Phosphorus Free Winterizer Plus Weed Control

## 18-0-12

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Harmful if inhaled or absorbed through skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

#### FIRST AID

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.  
**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.  
**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

#### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled. Do not apply more than 2 broadcast applications per treatment site per year with a minimum of 30 days between applications.

This product is a stable mixture of three herbicides. It is for use on all Residential Turf.

This product may be used on Bahia, common Bermuda, Bluegrass, Centipede, Fescue, Rye, St. Augustine and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as:

Alder	Dock	Nettle	Stinging Nettle
Annual Yellow	Dogbane	Orange Hawkweed	Stinkweed
Sweet Clover	Dogfennel	Oxalis	Stitchwort
Artichoke	Elderberry	Oxeye Daisy	Strawberry Clover
Aster	English Daisy	Parsley-piert	Sumac
Austrian Fieldcress	Fall Dandelion	Parsnip	Sunflower
Bedstraw	False Dandelion	Pearwort	Sweet Clover
Beggarick	False Flax	Pennycress	Tall Nettle
Biden	False Sunflower	Pennywort	Tall Vervain
Bindweed	Fiddleneck	Peppergrass	Tansy Mustard
Bird Vetch	Field Bindweed	Pepperweed	Tansy Ragwort
Bitterweed	Field Pansy	Pigweed	Tanweed
Bitter Wintercress	Flea Bane (Daisy)	Pineywoods Bedstraw	Tarweed
Black-eyed Susan	Flixweed	Plains Coreopsis	Thistle
Black Medic	Florida Betony	Plantain	Tick Trefoil
Black Mustard	Florida Pusley	Poison Hemlock	Toadflax
Blackseed Plantain	Frenchweed	Poison Ivy	Trailing Crownvetch
Blessed Thistle	Galinsoga	Poison Oak	Tumble Mustard
Blue Lettuce	Garlic Mustard	Pokeweed	Tumble Pigweed
Blue Vervain	Goathead	Goathead	Tumbleweed
Box Elder	Goatsbeard	Povertyweed	Velvet Leaf
Bracted Plantain	Goldenrod	Prickly Lettuce	Veronica
Brassbuttons	Ground-Ivy	Prickly Sida	Vervain
Bristly Oxtongue	Gumweed	Primrose	Vetch
Broadleaf Dock	Hairy Bittercress	Prostrate Knotweed	Virginia Buttonweed
Broadleaf Plantain	Hairy Fleabane	Prostrate Pigweed	Virginia Creeper
Broomweed	Hawkweed	Prostrate Spurge	Virginia Pepperweed
Buckhorn	Healall	Prostrate Vervain	Wavyleaf Bullthistle
Buckhorn Plantain	Heartleaf Drymary	Puncture Vine	Western Clematis
Bulbous Buttercup	Hedge Blindweed	Purslane	Western Salsify
Bull Nettle	Hedge Mustard	Ragweed	White Clover
Bull Thistle	Hemp	Red Clover	White Mustard
Burdock	Henbit	Redroot Pigweed	Wild Mustard
Burning Nettle	Hoary Cress	Red Sorrel	Wild Aster
Bur Ragweed	Hoary Plantain	Redstem Fillaree	Wild Buckwheat
Burweed	Hoary Vervain	Rough Cinquefoil	Wild Carrot
Buttercup	Honeysuckle	Rough Fleabane	Wild Four-O-Clock
Canada Thistle	Hop Clover	Round Leafed Marigold	Wild Garlic
Carolina Geranium	Horsenettle	Rush	Wild Geranium
Carpetweed	Horsetail	Russian Pigweed	Wild Lettuce
Catchweed Bedstraw	Indiana Mallow	Russian Thistle	Wild Marigold
Catsear	Ironweed	St. Johnswort	Wild Onion
Catnip	Jewelweed	Scarlet Pimpernel	Wild Parsnip
Chickweed	Jimsonweed	Scotch Thistle	Wild Radish
Chicory	Kochia	Sheep Sorrel	Wild Rape
Cinquefoil	Knawel	Sheepspurse	Wild Strawberry
Clover	Knotweed	Slender Plantain	Wild Sweet Potato
Cockle	Lambsquarter	Smallflower Galinsoga	Wild Vetch
Cocklebur	Lespedeza	Smartweed	Willow
Coffeebean	Locoweed	Smooth Dock	Witchweed
Coffeeweed	Lupine	Smooth Pigweed	Woolly Momingglory
Common Chickweed	Mallow	Sneezeweed	Woodsorrel
Common Mullein	Marshelder	Southern Wild Rose	Woolly Croton
Common	Matchweed	Sowthistle	Woolly Plantain
Sowthistle	Mexicanweed	Spanishneedle	Wormseed
Corn Chamomile	Milk Vetch	Spatterdock	Yarrow
Creeping Jenny	Milkweed Bloodflower	Speedwell	Yellow Rocket
Crimson Clover	Mugwort	Spiny Amaranth	Yellowflower
Croton	Morningglory	Spiny Cocklebur	Pepperweed
Cudweed	Mouseear Chickweed	Spotted Catsear	and other broadleaf
Curly Dock	Musk Thistle	Spotted Knapweed	weeds
Curly Indigo	Mustard	Spotted Spurge	
Dandelion	Narrowleaf Plantain	Spurge	
Dead Nettle	Narrowleaf Vetch	Spurweed	

#### WHEN TO APPLY

Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a

rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur. NOTE: When applying to Southern grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

#### HOW TO APPLY

This 18 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is equal to one application.

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

#### SUGGESTED SPREADER APPLICATION

This 18 pound bag should be used on 5,000 square feet (3.6 lbs. per 1,000 square feet). To determine how much product you need for your lawn, multiply your lawn's length by its width to get total square feet - then subtract non-lawn areas.


The following settings are approximate for applying contents of this package at a suggested rate of 3.6 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for a slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 0.9 lb. per 250 square feet.

Be sure to close spreader openings when loading, turning or when stopped. Most fertilizers will stain sidewalks, patios, swimming pools and other concrete surfaces. Remove spillage promptly by sweeping or vacuuming.

#### BROADCAST SPREADER SETTINGS (One Pass Application)

Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.


	3.6# Rate
Cyclone B1-84	4
Earthway 2000A	16
GreenThumb 2000A	16
Ortho 6000	4
Precision SB50	6 3/4
Republic 540/75450	5
Scotts Edgeguard, Speedygreen	6
Scotts Easygreen	27



#### DROP SPREADER SETTINGS (One Pass Application)

First apply 2 border strips along the edge of each end of the lawn (see Path B). Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.

	3.6# Rate
Earthway 7350	18
GreenThumb 7350SU	18
Ortho 5000	7
Precision FH22	9
Republic 530/75350	7
Scotts Accugreen	6 1/2
Scotts PF4	5 1/2



#### GUARANTEED ANALYSIS

Total Nitrogen (N)	18.00%
18.00% Urea Nitrogen*	
Soluble Potash (K <sub>2</sub> O)	12.00%
Plant nutrients derived from urea, potassium chloride, and polymer coated sulfur coated urea.	
<b>*3.00% Slowly Available Nitrogen from Polymer Coated Sulfur Coated Urea</b>	<b>F483</b>
Information regarding the contents and levels of metals in this product is available on the internet at <a href="http://www.regulatory-info-ap.com">http://www.regulatory-info-ap.com</a>	

#### STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizer, seed, or other pesticides.

**CONTAINER DISPOSAL:** None refillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled -** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

#### WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

#### LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded. (RV091406N)

#### Distributed by:

**Free Flow Fertilizer**  
P. O. Box 119, Maumee, Ohio 43537

©2009 T.A. Fortify is a registered trademark of The Andersons, Inc.



Visit our website at:  
[www.fortifyyourhome.com](http://www.fortifyyourhome.com)

**FRT183DS1**  
**FRT183DS1B**

**UPC**  
**0 31865 31416 2**  
**FPO**