



15-5-10 Fertilizer Plus 1.0% RONSTAR®

Net Wt. 1,000 Lb.
(453.59 kg)

FEB 13 2011

Regi.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2008-2010 LIC. NO.

- Provides preemergent control or suppression of crabgrass, *Poa annua*, goosegrass, and foxtail
- Provides a light nitrogen feeding

FOR SALE TO, USE AND STORAGE BY PROFESSIONAL TURF MAINTENANCE PERSONNEL ONLY
NOT FOR USE ON HOME LAWNS

Guaranteed Analysis (15-5-10)		F292
Total Nitrogen (N)	15.00%	
1.96% Ammoniacal Nitrogen		
13.04% Urea Nitrogen*		
Available Phosphate (P ₂ O ₅)	5.00%	
Soluble Potash (K ₂ O)	10.00%	
Sulfur (S)	2.43%	
2.43% Free Sulfur		
Plant nutrients derived from diammonium phosphate, urea, potassium chloride, polymer coated sulfur-coated urea.		
Chlorine (Cl) Max	8.46%	
6.00% Slowly available nitrogen from polymer coated sulfur coated urea.		

ACTIVE INGREDIENT:	
Oxadiazon	1.0%
Other Ingredients	99.0%
Total	100.0%
EPA Reg. No. 9198-203	CAS # 19666-30-9
EPA EST. NO. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A	
UNDERLINED LETTER IS FIRST LETTER USED IN RUN CODE ON BAG.	
This product and/or its use is covered by one or more of the following US patents and foreign counterparts thereof: US Pat No 5,219,465; 5,300,135	
Information regarding the contents and levels of metals in this product is available on the internet at http://www.regulatory-info-ap.com/ .	
The Andersons Golf Products logo is a trademark of The Andersons Agriservices, Inc.	
Ronstar is a registered trademark of Rhône-Poulenc.	



LICENSED

PERIOD 2011-2013 LIC. NO.

KEEP OUT OF REACH OF CHILDREN
WARNING
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING. May be fatal if inhaled. Causes substantial but temporary eye injury. Wear goggles or safety glasses when applying this product. Causes skin irritation. Do not get in eyes, on skin, or on clothing. Wear protective clothing including long-sleeved shirt and gloves when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT
IF SWALLOWED: Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. **IF IN EYES:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention. **IF ON SKIN:** Wash with plenty of soap and water. Get medical attention. **IF INHALED:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS
This pesticide is toxic to fish. Use with care when applying to areas adjacent to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.



LICENSED

PERIOD 2014-2016 LIC. NO.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
*Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
For use on established or sprigged bermudagrass, zoysiagrass, Seashore paspalum (see precautions), and St. Augustinegrass ornamental turf for preemergent control or suppression of sprouting:

crabgrass	yellow foxtail	<i>Poa annua</i>
goosegrass	green foxtail	
- *Not currently registered in California.

HOW TO USE

- Apply prior to or after sprigging 200 to 300 lbs. of product per acre (2.0 to 3.0 lbs. ai/acre) for suppression or control of listed annual grassy weeds in bermudagrass, zoysiagrass, and St. Augustinegrass.
- After application of this product on newly sprigged bermudagrass, zoysiagrass, and St. Augustinegrass, wait until the grass has filled in and the stolons have rooted down before application of other PRETURF herbicide, combination fertilizer/herbicide or growth regulator products.
- On Seashore paspalum make application 10-14 days after sprigging.
- For established turf, apply at 200 to 300 lbs. of product per acre (2.0 to 3.0 lbs. ai/acre).

WHEN TO USE

Use this product to control or suppress the listed annual grassy weeds when establishing new turf areas or when treating established turf that has been thinned from winter injury, insects, diseases or other factors.

SUGGESTED SPREADER SETTINGS

SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS		
			200 Lbs. Prod/acre (2.0 lbs ai/A) 0.69 lbs./NM	250 Lbs. Prod/acre (2.5 lbs ai/A) 0.86 lbs./NM	300 Lbs. Prod/acre (3.0 lbs ai/A) 1.03 lbs./NM
50 lbs. treats 10,890 ft ² at the 200 lbs. product/acre (2.0 lbs. ai/A)					
50 lbs. treats 8,712 ft ² at the 250 lbs. product/acre (2.5 lbs. ai/A)					
50 lbs. treats 7,260 ft ² at the 300 lbs. product/acre (3.0 lbs. ai/A)					
Scotts Rotaries					
R-7, R-7X*	3 mph	12 ft	K	L 1/2	N
R-8A, Accu Pro & SR-2000 (cone 4)	3 mph	13 ft	M 1/2	N 1/2	P
Lely Models					
(See below)	4.5 mph	40 ft	71	81	8 1/2 1
Vicon (02 models)	4.5 mph	50 ft	40	45	49
Vicon (03 models)	4.5 mph	36 ft	33	37	42

*For the R-7X model, use adjustable pattern slide 1/4 closed.
Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.

PRECAUTIONS

- Avoid contact with flowers and shrubs.
- This product should not be used on established putting greens and tees because of the varied cultural practices used.
- Do not apply product to moist turfgrass foliage.
- This product may cause discoloration of bermudagrass and St. Augustinegrass. This discoloration is temporary and will normally be outgrown within 2 to 3 weeks.
- On Seashore paspalum wait 10-14 days after sprigging before application. An application made close to the time of sprigging may cause objectionable injury.
- Irrigate immediately after application for best results.
- Seeding into treated areas should be delayed until 4 months after treatment.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.
STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Manufactured by: The Andersons Lawn Fertilizer Division, Inc., P.O. Box 119, Maumee, OH 43537

Attention: This product contains a chemical known to the State of California to cause cancer.

J1-(BULKB)

AGC8684-1BK

PERIOD 2014-2016 LIC. NO. **8054.5**

DEPT. OF AGRICULTURE

LICENSED

Period 2014-2016 Lic. No. 8054.5