



LICENSED

PERIOD 2014-2016 LIC. NO. 8054.63

128392




Early Spring

Crabgrass Preventer + LAWN FERTILIZER 27-0-5



For a thick, green crabgrass free lawn



Early Spring

Controls Crabgrass all season long

Extended Greening from Controlled Release Nitrogen

Covers up to 16,000 square feet  
NET WT 45 LBS (20.41 Kg)

27-0-5



# Crabgrass Preventer + LAWN FERTILIZER

A selective preemergence herbicide and fertilizer product for the control of crabgrass and most other annual weed grasses on lawns in established Bluegrass (perennial), Ryegrass (perennial), Fescue, Bahiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass and Zoysiagrass.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See back panel for Additional Precautionary Statements

EPA Reg. No. 9198-120  
EPA Est. No. 9198-OH-1M, 9198-OH-2S, 9198-AL-001A  
Underlined letter used is first letter of run code on end of bag.

ACTIVE INGREDIENT:

|  |         |
|--|---------|
| *Dithiopyr, 3,5-pyridinedicarboxylic acid, 2-(difluoromethyl)-4-(2-methylpropyl) |         |
| -6-(trifluoromethyl)-S,S-dimethyl ester  | 0.103%  |
| INERT INGREDIENTS  | 99.897% |
| TOTAL  | 100.00% |



Crabgrass Preventer + LAWN FERTILIZER 27-0-5

Early Spring





Early Spring

# Crabgrass Preventer + LAWN FERTILIZER 27-0-5



## 4 Season Cycle Fertilizer Application Guide

| Turf Zone & Variety                         | February                   | March                      | April              | May                | June                               | July                               | August             | September          | October | November | December |
|---|----------------------------|----------------------------|--------------------|--------------------|------------------------------------|------------------------------------|--------------------|--------------------|---------|----------|----------|
| <b>NORTHERN ZONE</b><br>Cool Season Grasses |                            | 27-0-5 Crabgrass Preventer |                    | 29-0-3 Weed & Feed |                                    | 25-0-5 Fertilizer + Insect Control |                    | 24-0-12 Winterizer |         |          |          |
|   |                            |                            |                    |                    | 30-0-3 Lawn Fertilizer             |                                    |                    |                    |         |          |          |
| <b>TRANSITIONAL ZONE</b>                    |                            | 27-0-5 Crabgrass Preventer |                    | 29-0-3 Weed & Feed |                                    | 25-0-5 Fertilizer + Insect Control |                    | 24-0-12 Winterizer |         |          |          |
|   |                            |                            |                    |                    | 30-0-3 Lawn Fertilizer             |                                    |                    |                    |         |          |          |
| <b>SOUTHERN ZONE</b><br>Warm Season Grasses | 27-0-5 Crabgrass Preventer |                            | 29-0-3 Weed & Feed |                    | 25-0-5 Fertilizer + Insect Control |                                    | 24-0-12 Winterizer |                    |         |          |          |
|   |                            |                            |                    |                    | 30-0-3 Lawn Fertilizer             |                                    |                    |                    |         |          |          |



U.S. Climate Zones:

**Early Spring**

Crabgrass Preventer + LAWN FERTILIZER 27-0-5

**Late Spring**

Weed & Feed LAWN FERTILIZER 29-0-3

**Summer**

Insect Control + LAWN FERTILIZER 25-0-5

**Fall**

Winterizer LAWN FERTILIZER 24-0-12

**Four Seasons**

Lawn Fertilizer 30-0-3

**TURF VARIETIES BY ZONE:**

- Bluegrass, Ryegrass, Fine & Tall Fescues
- Bluegrass, Ryegrass, Fine & Tall Fescues, Bermudagrass, Zoysiagrass
- Bermudagrass, Zoysiagrass, Centipedegrass, Bahiagrass, St. Augustine

Abnormal climatic conditions can affect appropriate timing.

**GUARANTEED ANALYSIS:**

Total Nitrogen (N) ..... 27.00%  
27.00% Urea Nitrogen\*

Soluble Potash (K<sub>2</sub>O) ..... 5.00%

Plant nutrients derived from urea, potassium chloride and polymer coated sulfur coated urea.  
\*6.75% slowly available urea nitrogen from polymer coated sulfur coated urea F292

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info.com>

### PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals Keep Out of Reach of Children **CAUTION**

Causes eye irritation. Harmful if inhaled. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

|   |
|---|
| <b>FIRST AID</b>  |
| If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.                                     |
| If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.                                   |
| If swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person. |
| If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.   |
| Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8851 for emergency medical treatment information.  |

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply when weather conditions favor drift from treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

New York Use Restrictions: Do not apply more than 5.6 pounds of this product per 1,000 square feet of turfgrass area per year. Do not make more than one application per year.

Use only according to label instructions.

**GENERAL INFORMATION**

This product is a combination product that provides both crabgrass control and the required nutrients for optimum turfgrass growth all in a single product.

This product contains Dimension®, a crabgrass preventer that can be applied up to 4 weeks later than other crabgrass preventers and provides up to 4 months control of crabgrass and other grass and broadleaf weeds prior to their establishment in lawns and ornamental turfgrasses. This product should be applied to lawns and ornamental turfgrasses that are well established (See "Use Precautions" section below for a list of the tolerant turfgrasses). This product will not harm nearby established ornamentals when used according to label directions.

**USE PRECAUTIONS**

This product should only be applied to lawns and ornamental turfgrasses that are well established. Use of this product on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury.

This product can be applied to the following turfgrasses when applied as directed:

| Cool-Season Turfgrasses | Warm-Season Turfgrasses |
|-------------------------|-------------------------|
| Creeping Bentgrass      | Bahiagrass              |
| Kentucky Bluegrass      | Bermudagrass            |
| Fine Fescues            | Buffalograss            |
| Tall Fescues            | Carpetgrass             |
| Perennial Ryegrass      | Centipedegrass          |
|                         | St. Augustinegrass      |
|                         | Zoysiagrass             |

This product should be applied in the spring or fall prior to the weed seed germination in established lawns and ornamental turfgrasses.

This product can be applied up to 4 weeks later than other crabgrass preventers and still provide season-long weed control of crabgrass. Dimension® herbicide provides preemergence control of crabgrass and other weeds when applied prior to their germination, and it also provides postemergence control of crabgrass. The postemergence control provided by this product is limited to crabgrass only and will provide control of crabgrass when applied up to 4 weeks after crabgrass germination. Application of this product later than 4 weeks after crabgrass germination will result in reduced control of crabgrass.

For control of other grass and broadleaf weeds, this product must be applied as a preemergent treatment. Only crabgrass can be controlled with delayed applications after germination.

**Tips for Improved Weed Control:** For best results, apply this product within a few days after mowing and delay further mowings after the application of this product until the lawn or ornamental turfgrass has been watered or received significant rainfall to dissolve the fertilizer granules. Improved weed control may be observed when treated lawn or ornamental turfgrass is watered or receives rainfall within a few days after application of this product.

Reseeding, Overseeding or Sprigging of turfgrasses into areas that have been previously treated with this product should be delayed until 16 weeks after application. Reseeding, overseeding, or sprigging up to 16 weeks after application may prevent the establishment of newly reseeded, overseeded or sprigged turfgrasses.

- DO NOT apply more than 11 pounds of this product per 1,000 square feet of turfgrass area during the year (1,000 square feet area = 100 feet x 10 feet).
- DO NOT apply to flowers, vegetables, or fruit trees.
- DO NOT use clippings from treated turf for mulching around flowers, vegetables or fruit trees.
- Keep people and pets off treated areas until dust has settled.

**APPLICATION DIRECTIONS**

This product will provide control of crabgrass and other grasses and broadleaf weeds listed below when applied prior to weed seed germination in established lawns and ornamental turfgrasses.

| Grass Weeds          |                    | Broadleaf Weeds   |                   |
|----------------------|--------------------|-------------------|-------------------|
| Annual Bluegrass     | Annual Ryegrass    | Buttercup Oxalis  | Spotted Spurge    |
| Bermudagrass         | Bentgrass          | Carolina Geranium | Chickweed         |
| Crabgrass            | Crowfootgrass      | Common Lespedeza  | Common Purslane   |
| Goosegrass           | Green Foxtail      | Common Purslane   | Corn Speedwell    |
| Kleuygrass (seeding) | Perennial Ryegrass | Creeping Oxalis   | Herbit            |
| Smutgrass (seeding)  | Yellow Foxtail     | Parsley-piert     |                   |
|                      |                    | Pineappleweed     | Yellow Woodsorrel |

**Application Timings:**  
This product may be applied either in the spring or in the fall for the control of weeds.

When this product is applied in the fall, a sequential or second application should be made the following spring to provide season-long control of weeds.

**Length of Control:**  
This product will provide up to 4 months of weed control if applied according to label directions.

The actual length of control will vary depending on use rate, weather conditions, condition of the turfgrass, intensity of weed pressure, and the location or "Region" within the United States that this product is being applied.

Please check the map below to determine which "Region" of the U.S. that applies to your particular use area.

Use "Regions" within the United States:



**Recommended Use Rates:**

| Region of U.S. | Recommended Use Rates for This Product<br>(Pounds of this product to apply per 1,000 square feet of turfgrass) |                    |
|----------------|--|--------------------|
|                | 2-3 Months Control   | 3-4 Months Control |
| North          | 2.8  | 4.0                |
| South          | 4.3  | 5.0                |

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage and disposal. **STORAGE:** Keep container closed and away from food, feedstuffs and domestic water supplies. **DISPOSAL:** If empty: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**IMPORTANT: READ BEFORE USE:**

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

**WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY:**

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

**LEGAL RIGHTS:**

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/STATE/JURISDICTION TO STATE/JURISDICTION.

### Suggested Spreader Settings

Important: It is important to follow the directions as indicated on this quality lawn care product as over-application may cause injury to the lawn. Measure the size of the area to be treated and do not over or under apply. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Calibrate spreader prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly. Close spreader when filling, stopping or turning. Do not hand spread.

#### BROADCAST SPREADER SETTINGS (One Pass Application)



For best results, apply in a circular path starting on the outside and working toward the center. A little overlap of product will ensure complete coverage.

|   | 2.8# Rate | 4.0# Rate | 4.3# Rate | 5.0# Rate |
|---|-----------|-----------|-----------|-----------|
| Agri-Fab 100  | 4         | 4 3/4     | 5         | 6         |
| Cyclone B1-B4   | 3 1/2     | 4 1/2     | 5         | 6         |
| Earthway: 1950, 2000A, 2000AP, 2030, 2400                   | 12        | 14        | 14 1/2    | 15 1/2    |
| Green Thumb: 2000A GT2030, GT2170                           |           |           |           |           |
| GTH2030 P-PLUS  | 12        | 14        | 14 1/2    | 15 1/2    |
| Precision: SB4300   | 4 1/4     | 5 1/2     | 5 3/4     | 7         |
| Red Devil: SB4300   | 4         | 4 3/4     | 5         | 6 1/4     |
| Republic: EZ 540/75450                                      | 4 1/4     | 5         | 5 1/4     | 6 1/2     |
| Scotts: Basic, Deluxe, EdgeGuard, LawnPro, SG3000, Standard | 4 1/2     | 5 3/4     | 6         | 6 1/2     |
| Scotts Easygreen EG3  | 27        | 28 1/2    | 29        | 31        |

#### DROP SPREADER SETTINGS (One Pass Application)

First apply 2 border strips along the edge of each end of the lawn (see Path B). Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.

|                       | 2.8# Rate | 4.0# Rate | 4.3# Rate | 5.0# Rate |
|-----------------------|-----------|-----------|-----------|-----------|
| Earthway 7350, 7350SU | 14        | 16        | 16 1/2    | 17 1/2    |
| GreenThumb 7350SU     | 14        | 16        | 16 1/2    | 17 1/2    |
| Precision DS 1000     | 8         | 9 1/2     | 9 3/4     | 10 1/2    |
| Red Devil DS 1000     | 7         | 8         | 8 1/4     | 9 1/2     |
| Republic 530/75350    | 5 1/2     | 7         | 7 1/4     | 8 1/2     |
| Scotts AG3000         | 5 1/2     | 7         | 7 1/4     | 8 1/2     |



Manufactured by:  
The Andersons  
Lawn Fertilizer Division, Inc.  
P.O. Box 119  
Maumee, OH 43537  
©2013 T.A.

Made in USA  
**GTH276DM45**  
GTH276DM45B

