



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO. **8071.10**



GARDEN *TECH*

Sevin®

Ready-To-Use

BUG KILLER



Japanese Beetles



Cabbage Worms



Ticks



Ants

Kills Over 100 Insect Pests on Ornamentals, Flowers, Fruits and Vegetable Gardens

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Back Panel For Precautionary Statements
PRECAUCION AL USUARIO: Si usted no le ingles,
no use este producto hasta que la etiqueta le haya
sido explicada ampliamente.

ACTIVE INGREDIENT:
Carbaryl (1-naphthyl
N-methylcarbamate)..... 0.126%
INERT INGREDIENTS:... 99.874%
TOTAL..... 100.000%

Net Contents 32 fl. oz. (946 ml)

EPA Reg. No. 432-1211-71004
EPA Est. No. 67572-GA-1

GARDEN TECH
Sevin

READY-TO-USE BUG KILLER

For Vegetable Pests



Alfalfa Caterpillar, Armyworm, Aster Leafhopper, Bean Leaf Beetle, Cucumber Beetles, Cutworms, European Corn Borer, Flea Beetles, Green Cloverworm, Imported Cabbage Worm, Japanese Beetle, Lace Bugs, Leafhoppers, Melonworm, Squash Bugs, Sweet Potato Weevil, Thrips, Tomato Hornworm & many more listed inside.

For Pests Around Buildings



Ants, Bees, Brown Dog Ticks, Carpenter Ants, Centipedes, Cockroaches, Crickets, Earwigs, Firebrats, Fleas, Millipedes & many more listed inside.

For Ornamental Pests



Ants, Armyworm, Blister Beetle, Browntail Moth, Flea Beetles, Gall Midges, Grasshoppers, June Beetles, Leafrollers, Mealybugs, Oak Leafminers, Pine Sawfly, Rose Aphid, Rosslug, Springtails, Thorn Bug, Ticks & many more listed inside.

Peel Back Tab for Additional Information

Distributed by:
TechPac, LLC
P.O. Box 24830
Lexington, KY 40524



PRECAUTIONARY STATEMENTS
Hazard to Humans and Domestic Animals
CAUTION

Harmful if swallowed or inhaled. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

USER SAFETY REQUIREMENTS:

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Wear a hat and eye protection when making overhead applications.
- Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow children or pets to contact treated area until sprays have dried.

FIRST AID

This product is an N-methyl carbamate and is a cholinesterase inhibitor.

If Swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
---------------------	---

(continued)

NOTES:

NOTES:

14

FIRST AID (continued)

If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If in Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on Skin or Clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **1-800-420-9347** for emergency treatment information.

NOTE TO PHYSICIAN

Carbaryl is an N-methyl carbamate insecticide which is a cholinesterase inhibitor. Specific antidote consists of atropine. Narcotics and other sedatives should not be used. Drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

3

BEE CAUTION

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

PLANT RESPONSE PRECAUTIONS

Do not use on Boston Ivy, Virginia creeper and Maidenhair fern as injury will result. During early season, it may also injure Virginia and Sand pines.

DO NOT PLANT ROTATIONAL FOOD AND FEED CROPS NOT LISTED ON THIS OR OTHER CARBARYL LABELS IN CARBARYL TREATED SOIL.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE CONTAINER BEFORE USE

Hold sprayer 8 to 12 inches from the area to be treated. Spray until surface is wet. Apply when air is calm to avoid drift and contact with eyes and skin. This product is not recommended to be applied in mid-day sun or when temperatures exceed 85°F as temporary leaf burn may occur. Start spraying at the farthest corner of the treatment area and work backward to avoid contact with wet surfaces. Allow spray to dry in treated areas before reentering. For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

STORAGE AND DISPOSAL

STORAGE: Store unused product in an area out of reach of children and animals. Do not store in areas where temperatures frequently exceed 100° F.

DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

IMPORTANT: READ BEFORE USE (continued)

to make any warranties beyond those contained herein or to modify the warranties contained herein. TECHPAC, LLC. DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT TECHPAC, LLC.'S ELECTION, THE REPLACEMENT OF PRODUCT.

GardenTech® is a registered trademark of Gulfstream Home & Garden, Inc.

Sevin® is a registered trademark of Bayer for Carbaryl Insecticides

If you have a medical emergency, need consultation on diagnosis, therapy and analysis please call (800) 420-9347

If you have questions relating to product use or application, please call (800) 969-7200

If you have a product spill or transportation emergency call (800) 424-9300

For Material Safety Data Sheets please go to www.gardentech.com, or call (800) 969-7200

Distributed by:
TechPac, LLC.
P.O. Box 24830
Lexington, Ky 40524

PESTS

Ants	Firebrats
Bees	Fleas
Brown Dog Ticks	Millipedes
Carpenter Ants	Scorpions
Centipedes	Silverfish
Cockroaches	Spiders
Crickets	Wasps
Earwigs	

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of TechPac, LLC. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TECHPAC, LLC. MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of TechPac, LLC. is authorized

VEGETABLE CROPS

For control of all the vegetable pests listed below, apply this product in adequate volume to effectively cover both the upper and lower surfaces of the plant. **NOTE:** This product is not registered for use on celery and sweet potatoes in California.

PHI refers to the minimum interval in days that must be observed between the date of the last application and date of harvest. **APP** refers to the maximum number of applications per season.

CROPS	PHI	APP
Asparagus	1	3
Crowder peas, Southern peas	3	6
Corn, Sweet	2	8
Broccoli, Brussels sprouts, Cabbage, Cauliflower, Chinese Cabbage, Kohlrabi	14	3
Eggplant, Tomato	3	4
Black-eyed peas, Cowpeas, Cucumber, Green beans, Lima beans, Melons, Navy beans, Okra, Peas, Peppers, Pumpkin, Snap beans, Squash	3	6
Carrots, Dandelion, Garden beet roots, Garden beet tops, Horseradish, Parsnips, Potatos, Radishes, Rutabagas, Salsify roots, Turnip roots	14	3
Sweet Potatoes	7	4
Celery, Collards, Endive (Escarole), Hanover salads, Head lettuce, Kale, Leaf lettuce, Mustard greens, Parsley, Salsify roots, Spinach, Swiss chard, Turnip tops	14	3
Peanuts	14	5
Dried beans	21	6

PESTS

Alfalfa caterpillar	Lima bean pod borer
Apache cicada	Lygus bugs
Armyworms	Melonworm
Asparagus beetle	Mexican bean beetle
Aster leafhopper	Pea leaf weevil
Bean leaf beetle	Pea weevil
Chinch bug	Rednecked peanutworm
Colorado potato beetle	Sap beetles
Corn earworm	Southwestern corn borer
Corn rootworm adult	Spittlebugs
Cowpea curculio	Squash bugs
Cucumber beetle	Stink bugs
Cutworms	Sweet potato weevil
European corn borer	Three cornered alfalfa hopper
Fall armyworm	Thrips
Flea beetles	Tomato hornworm
Grasshoppers	Tomato pinworm
Green cloverworm	Tortoise beetle
Harlequin bug	Velvetbean caterpillar
Imported cabbageworm	Webworms
Japanese beetle	Western bean cutworm
Lace bugs	Whitefringed beetle (adults)
Leafhoppers	Yellowstriped armyworm

PESTS (continued)

Psyllids	Spruce Needleminer
Puss Caterpillar	Subtropical Pine Tip Moth
Redhumped Oakworm	Tent Caterpillar
Rose Aphid	Thorn Bug
Rose Chafer	Thrips (exposed)
Roseslug	Ticks
Saddled Prominent	Walnut Caterpillar
Sawflies (exposed)	Webworms
Scale Insects	Western Hemlock Looper
Sowbugs	Western Spruce Budworm
Spiny Elm Caterpillar	Willow Leaf Beetles
Springtails	Yellow Poplar Weevil
Spruce Budworm	

PEST CONTROL AROUND BUILDINGS AND OTHER OUTDOOR AREAS

This product may be used as a spot treatment around buildings such as homes, apartments, warehouses, barns and municipal and recreational areas to control the pests listed below. Thoroughly wet the outside perimeter of dwellings and other areas where pests tend to congregate.

NOTE: Staining may occur on certain surfaces such as stucco, brick, cinder block and wood. Therefore, applications of this product to surfaces where a noticeable residue or discoloration is objectionable should be avoided. Do not apply to carpets or draperies as staining may occur. Care should also be exercised to avoid spotting of wallpaper and fabrics.

Do not use this product in commercial food areas of food handling establishments, restaurants or other places where food is prepared or processed. Do not use in serving areas while food is exposed.

PESTS

Ants	Holly Budmoth
Apple Aphid	Holly Leafminer
Armyworm	Jackpine Budworm
Azalea Leafminer	Japanese Beetle
Bagworms	Jeffrey Pine Needleminer
Birch Leafminer	June Beetles
Blister Beetle	Plant Bugs
Boxelder Bug	Lace Bugs
Boxwood Leafminer	Leathoppers
Browntail moth	Leafrollers
Cankerworms	Locust Borer
Catalpa Sphinx	Maple Leafcutter
Chiggers	Mealybugs
Cooley Spruce Gall	Mimosa Webworm
Elm Leaf Aphid	Nantucket Pine Tip Moth
Elm Spanworm	Oak Leafminers
Eriophyid Mites	Oak Leaf Skeletonizer
European Pine Shoot Moth	Oakworm Complex
Fall Armyworm	Oleander Caterpillar
Flea Beetles	Olive Ash Borer
Fuller Rose Beetles	Orange-striped Oakworm
Gall Midges	Orange Tortrix
Gall Wasps	Periodical Cicada
Grasshoppers	Pine Sawfly
Green Striped Mapleworm	Pine Spittlebug
Gypsy Moth	Pitch Pine Tip Moth
Hackberry Nipplegall Makers	Poinsettia Hornworm

(continued)

FRUIT CROPS

This product may be used as a spot treatment for localized pest infestations on the tree fruits listed below. Do not use on large trees or where pest populations are generally inaccessible to spray deposits, as inadequate control will result. Direct spray toward upper and lower leaf surfaces and between fruit clusters, and on small limbs and trunks to the point of runoff where pests appear.

Repeat applications as necessary up to a total of eight times for tree fruits and five times for small fruits but not more often than once every seven days.

NOTE: To avoid undesired apple thinning, delay use until at least 30 days after full bloom. Carbaryl insecticide may injure early dawn and sunrise strawberry varieties on the Delmarva Peninsula.

The number in parenthesis following each fruit and nut crop grouping refers to the minimum interval that must be observed between the date of the last application and the date of harvest.

TREE FRUITS

Apple, Apricots, Cherries, Nectarines, Peaches, Pears, Plums, Prunes (3 days until harvest)

SMALL FRUITS

Blackberries, Blueberries, Dewberries, Grapes, Raspberries, Strawberries (7 days until harvest)

PESTS

Apple Aphid	European Apple Sawfly
Apple Maggot	European Earwig
Apple Mealybug	European Raspberry Aphid
Apple Pandemis	Eyespotted Budmoth
Apple Rust Mite	Forbes Scale
Apple Sucker	Fruitree Leafroller
Avocado Leafroller	Grape Berry Moth
Bagworms	Grape Leafhopper
Black Cherry Aphid	Grape Leafroller
Black Scale	Green Fruitworm
Blueberry Maggot	Japanese Beetle
Brown Soft Scale	June Beetles
California Red Scale	Leafhoppers
Cherry Fruitworm	Oriental Fruit Moth
Cherry Maggot	Oystershell Scale
Citricola Scale	Peach Twig Borer
Citrus Cutworm	Pear Leaf Blister Mite
Citrus Root Weevil	Pear Psylla
Citrus Snow Scale	Pear Rust Mite
Codling Moth	Periodical Cicada
Cucumber Beetles	Plum Curculio
Cutworms	Prune Leafhopper
Eastern Tent Caterpillar	Raspberry Sawfly
Eightspotted Forester	Red-banded Leafroller

(continued)

PESTS (continued)

Rose Chafer	Variigated Leafroller
Rosy Apple Aphid	Western Grapeleaf Skeletonizer
Saltmarsh Caterpillar	Western Tussock Moth
San Jose Scale	Western Yellowstriped Armyworm
Snowy Tree Cricket	West Indian Sugarcane Borer (adults)
Strawberry Weevil	White Apple Leafhopper
Tarnished Plant Bug	Wooly Apple Aphid
Tentiform Leafminers	Yellow Headed Fireworm
Tussock Moth	

TREES, ORNAMENTALS, SHRUBS AND FLOWERS

This product may be used as a spot treatment for localized pest infestations on trees. Do not use on large trees or where pest populations are generally inaccessible to spray deposits, as inadequate control will result. Direct spray toward the upper and lower leaf surfaces and small trunks, stems and twigs to the point of runoff where pests appear. Repeat applications as necessary up to a total of four times for trees and six times for ornamentals, shrubs and flowers but not more often than once every 7 days.

TREES

Trees (including) Shade Trees, Shelter Belts, Plantations, Parks, Recreational Areas
Ornamentals (including) Roses, Other Herbaceous Plants, Woody Shrubs

**KILLS
JAPANESE
BEETLES**


GARDENTECH

Sevin®

**READY TO USE
BUG KILLER**



**Japanese
Beetles**



**Cabbage
Worms**



Ticks



Ants

**Kills Over 100 Insect Pests on
Ornamentals, Flowers, Fruits and Vegetable Gardens**

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Back Panel For Precautionary Statements

PRECAUCION AL USUARIO:

Si usted no le ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

Net Contents
1 gallon, 1 quart (4.73 L)

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate)..... 0.126%

INERT INGREDIENTS: 99.874%

TOTAL: 100.000%

EPA Reg. No. 432-1211-71004

EPA Est. No. 67572-GA-1

GARDEN TECH **Sevin**® **READY-TO-USE BUG KILLER**

For Vegetable Pests



Japanese Beetles



Grasshoppers



Cutworms



Cabbage Worms

Alfalfa Caterpillar, Armyworm, Aster Leafhopper, Bean Leaf Beetle, Cucumber Beetles, Cutworms, European Corn Borer, Flea Beetles, Green Cloverworm, Imported Cabbage Worm, Japanese Beetle, Lace Bugs, Leafhoppers, Melonworm, Squash Bugs, Sweet Potato Weevil, Thrips, Tomato Hornworm & many more listed inside.

For Pests Around Buildings



Ticks



Fleas



Ants



Ants, Bees, Brown Dog Ticks, Carpenter Ants, Centipedes, Cockroaches, Crickets, Earwigs, Firebrats, Fleas, Millipedes & many more listed inside.

For Ornamental Pests



Moths



Grasshoppers



Tent Caterpillars



Japanese Beetles

Ants, Armyworm, Blister Beetle, Browntail Moth, Flea Beetles, Gall Midges, Grasshoppers, June Beetles, Leafrollers, Mealybugs, Oak Leafminers, Pine Sawfly, Rose Aphid, Roseslug, Springtails, Thorn Bug, Ticks & many more listed inside.

Distributed by:
TechPac, LLC
P.O. Box 24830
Lexington, KY 40524



Peel Back Tab for Additional Information

PRECAUTIONARY STATEMENTS
Hazard to Humans and Domestic Animals

CAUTION

Harmful if swallowed or inhaled. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

USER SAFETY REQUIREMENTS:

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Wear a hat and eye protection when making overhead applications.
- Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow children or pets to contact treated area until sprays have dried.

FIRST AID

This product is an N-methyl carbamate and is a cholinesterase inhibitor.

If Swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If in Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on Skin or Clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

(continued)

FIRST AID (continued)

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **1-800-420-9347** for emergency treatment information

NOTE TO PHYSICIAN

Carbaryl is an N-methyl carbamate insecticide which is a cholinesterase inhibitor. Specific antidote consists of atropine. Narcotics and other sedatives should not be used. Drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

BEE CAUTION

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

PLANT RESPONSE PRECAUTIONS

Do not use on Boston Ivy, Virginia creeper and Maidenhair fern as injury will result. During early season, it may also injure Virginia and Sand pines.

DO NOT PLANT ROTATIONAL FOOD AND FEED CROPS NOT LISTED ON THIS OR OTHER CARBARYL LABELS IN CARBARYL TREATED SOIL.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE CONTAINER BEFORE USE

Hold sprayer 8 to 12 inches from the area to be treated. Spray until surface is wet. Apply when air is calm to avoid drift and contact with eyes and skin. This product is not recommended to be applied in mid-day sun or when temperatures exceed 85°F as temporary leaf burn may occur. Start spraying at the farthest corner of the treatment area and work backward to avoid contact with wet surfaces. Allow spray to dry in treated areas before reentering. For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

STORAGE AND DISPOSAL

STORAGE: Store unused product in an area out of reach of children and animals. Do not store in areas where temperatures frequently exceed 100° F.

DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

VEGETABLE CROPS

For control of all the vegetable pests listed below, apply this product in adequate volume to effectively cover both the upper and lower surfaces of the plant. **NOTE:** This product is not registered for use on celery and sweet potatoes in California.

PHI refers to the minimum interval in days that must be observed between the date of the last application and date of harvest. **APP** refers to the maximum number of applications per season.

CROPS	PHI	APP
Asparagus	1	3
Crowder peas, Southern peas	3	6
Corn, Sweet	2	8
Broccoli, Brussels sprouts, Cabbage, Cauliflower, Chinese Cabbage, Kohlrabi	14	3
Eggplant, Tomato	3	4
Black-eyed peas, Cowpeas, Cucumber, Green beans, Lima beans, Melons, Navy beans, Okra, Peas, Peppers, Pumpkin, Snap beans, Squash	3	6
Carrots, Dandelion, Garden beet roots, Garden beet tops, Horseradish, Parsnips, Potatos, Radishes, Rutabagas, Salsify roots, Turnip roots	14	3
Sweet Potatoes	7	4
Celery, Collards, Endive (Escarole), Hanover salads, Head lettuce, Kale, Leaf lettuce, Mustard greens, Parsley, Salsify roots, Spinach, Swiss chard, Turnip tops	14	3
Peanuts	14	5
Dried beans	21	6

VEGETABLE CROPS (continued)**PESTS**

Alfalfa caterpillar	Lima bean pod borer
Apache cicada	Lygus bugs
Armyworms	Melonworm
Asparagus beetle	Mexican bean beetle
Aster leafhopper	Pea leaf weevil
Bean leaf beetle	Pea weevil
Chinch bug	Rednecked peanutworm
Colorado potato beetle	Sap beetles
Corn earworm	Southwestern corn borer
Corn rootworm adult	Spittlebugs
Cowpea curculio	Squash bugs
Cucumber beetle	Stink bugs
Cutworms	Sweet potato weevil
European corn borer	Three cornered alfalfa hopper
Fall armyworm	Thrips
Flea beetles	Tomato hornworm
Grasshoppers	Tomato pinworm
Green cloverworm	Tortoise beetle
Harlequin bug	Velvetbean caterpillar
Imported cabbageworm	Webworms
Japanese beetle	Western bean cutworm
Lace bugs	Whitefringed beetle (adults)
Leafhoppers	Yellowstriped armyworm

FRUIT CROPS

This product may be used as a spot treatment for localized pest infestations on the tree fruits listed below. Do not use on large trees or where pest populations are generally inaccessible to spray deposits, as inadequate control will result. Direct spray toward upper and lower leaf surfaces and between fruit clusters, and on small limbs and trunks to the point of runoff where pests appear.

Repeat applications as necessary up to a total of eight times for tree fruits and five times for small fruits but not more often than once every seven days.

NOTE: To avoid undesired apple thinning, delay use until at least 30 days after full bloom.

Carbaryl insecticide may injure early dawn and sunrise strawberry varieties on the Delmarva Peninsula.

The number in parenthesis following each fruit and nut crop grouping refers to the minimum interval that must be observed between the date of the last application and the date of harvest.

TREE FRUITS

Apple, Apricots, Cherries, Nectarines, Peaches, Pears, Plums, Prunes **(3 days until harvest)**

SMALL FRUITS

Blackberries, Blueberries, Dewberries, Grapes, Raspberries, Strawberries **(7 days until harvest)**

PESTS

Apple Aphid, Apple Maggot, Apple Mealybug, Apple Pandemis, Apple Rust Mite, Apple Sucker, Avocado Leafroller, Bagworms, Black Cherry Aphid, Black Scale, Blueberry Maggot, Brown Soft Scale, California Red Scale, Cherry Fruitworm, Cherry Maggot, Citricola Scale, Citrus Cutworm, Citrus Root Weevil, Citrus Snow Scale, Codling Moth, Cucumber Beetles, Cutworms, Eastern Tent Caterpillar, Eightspotted Forester, European Apple Sawfly, European Earwig, European Raspberry Aphid, Eyespotted Budmoth, Forbes Scale, Fruitree Leafroller, Grape Berry Moth, Grape Leafhopper, Grape Leafroller, Green Fruitworm, Japanese Beetle, June Beetles, Leafhoppers, Oriental Fruit Moth, Oystershell Scale, Peach Twig Borer, Pear Leaf Blister Mite, Pear Psylla, Pear Rust Mite, Periodical Cicada, Plum Curculio, Prune Leafhopper, Raspberry Sawfly, Red-banded Leafroller, Rose Chafer, Rosy Apple Aphid, Saltmarsh Caterpillar, San Jose Scale, Snowy Tree Cricket, Strawberry Weevil, Tarnished Plant Bug, Tentiform Leafminers, Tussock Moth, Variegated Leafroller, Western Grapeleaf Skeletonizer, Western Tussock Moth, Western Yellowstriped Armyworm, West Indian Sugarcane Borer (adults), White Apple Leafhopper, Woolly Apple Aphid, Yellow Headed Fireworm

TREES, ORNAMENTALS, SHRUBS AND FLOWERS

This product may be used as a spot treatment for localized pest infestations on trees. Do not use on large trees or where pest populations are generally inaccessible to spray deposits, as inadequate control will result. Direct spray toward the upper and lower leaf surfaces and small trunks, stems and twigs to the point of runoff where pests appear. Repeat applications as necessary up to a total of four times for trees and six times for ornamentals, shrubs and flowers but not more often than once every 7 days.

TREES

Trees (including) Shade Trees, Shelter Belts, Plantations, Parks, Recreational Areas
Ornamentals (including) Roses, Other Herbaceous Plants, Woody Shrubs

PESTS

Ants, Apple Aphid, Armyworm, Azalea Leafminer, Bagworms, Birch Leafminer, Blister Beetle, Boxelder Bug, Boxwood Leafminer, Browntail moth, Cankerworms, Catalpa Sphinx, Chiggers, Cooley Spruce Gall, Elm Leaf Aphid, Elm Spanworm, Eriophyid Mites, European Pine Shoot Moth, Fall Armyworm, Flea Beetles, Fuller Rose Beetles, Gall Midges, Gall Wasps, Grasshoppers, Green Striped Mapleworm, Gypsy Moth, Hackberry Nipplegall Makers, Holly Budmoth, Holly Leafminer, Jackpine Budworm, Japanese Beetle, Jeffrey Pine Needleminer, June Beetles, Plant Bugs, Lace Bugs, Leafhoppers, Leafrollers, Locust Borer, Maple Leafcutter, Mealybugs, Mimosa Webworm, Nantucket Pine Tip Moth, Oak Leafminers, Oak Leaf Skeletonizer, Oakworm Complex, Oleander Caterpillar, Olive Ash Borer, Orange-striped Oakworm, Orange Tortrix, Periodical Cicada, Pine Sawfly, Pine Spittlebug, Pitch Pine Tip Moth, Poinsettia Hornworm, Psyllids, Puss Caterpillar, Redhumped Oakworm, Rose Aphid, Rose Chafer, Roseslug, Saddled Prominent, Sawflies (exposed), Scale Insects, Sowbugs, Spiny Elm Caterpillar, Springtails, Spruce Budworm, Spruce Needleminer, Subtropical Pine Tip Moth, Tent Caterpillar, Thorn Bug, Thrips (exposed), Ticks, Walnut Caterpillar, Webworms, Western Hemlock Loopers, Western Spruce Budworm, Willow Leaf Beetles, Yellow Poplar Weevil

PEST CONTROL AROUND BUILDINGS AND OTHER OUTDOOR AREAS

This product may be used as a spot treatment around buildings such as homes, apartments, warehouses, barns and municipal and recreational areas to control the pests listed below. Thoroughly wet the outside perimeter of dwellings and other areas where pests tend to congregate.

NOTE: Staining may occur on certain surfaces such as stucco, brick, cinder block and wood. Therefore, applications of this product to surfaces where a noticeable residue or discoloration is objectionable should be avoided. Do not apply to carpets or draperies as staining may occur. Care should also be exercised to avoid spotting of wallpaper and fabrics.

Do not use this product in commercial food areas of food handling establishments, restaurants or other places where food is prepared or processed. Do not use in serving areas while food is exposed.

PESTS

Ants, Bees, Brown Dog Ticks, Carpenter Ants, Centipedes, Cockroaches, Crickets, Earwigs, Firebrats, Fleas, Millipedes, Scorpions, Silverfish, Spiders, Wasps
--

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of TechPac, LLC. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TECHPAC, LLC. MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of TechPac, LLC. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TechPac, LLC. disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT TECHPAC, LLC. ELECTION, THE REPLACEMENT OF PRODUCT.

GardenTech® is a registered trademark of Gulfstream Home & Garden, Inc.

Sevin® is a registered trademark of Bayer CropScience for Carbaryl Insecticides

If you have a medical emergency, need consultation on diagnosis, therapy and analysis please call (800) 420-9347

If you have questions relating to product use or application, please call (800) 969-7200

If you have a product spill or transportation emergency call (800) 424-9300

For Material Safety Data Sheets please go to www.gardentech.com, or call (800) 969-7200

Distributed by:
TechPac, LLC.
P.O. Box 24830
Lexington, Ky 40524





PRECAUTIONARY STATEMENTS
Hazard to Humans and Domestic Animals

CAUTION

Harmful if swallowed or inhaled. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

USER SAFETY REQUIREMENTS:

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Wear a hat and eye protection when making overhead applications.
- Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow children or pets to contact treated area until sprays have dried.

FIRST AID

This product is an N-methyl carbamate and is a cholinesterase inhibitor.

If Swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If in Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on Skin or Clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **1-800-420-9347** for emergency treatment information.

NOTE TO PHYSICIAN

Carbaryl is an N-methyl carbamate insecticide which is a cholinesterase inhibitor. Specific antidote consists of atropine. Narcotics and other sedatives should not be used. Drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

If you have a medical emergency, need consultation on diagnosis, therapy and analysis please call (800) 420-9347
If you have questions relating to product use or application, please call (800) 969-7200
If you have a product spill or transportation emergency call (800) 424-9300
For Material Safety Data Sheets please go to www.gardentech.com, or call (800) 969-7200

Distributed by:
TechPac, LLC.
P.O. Box 24830
Lexington, Ky 40524

