



LICENSED

PERIOD **2015-2017** LIC. NO.

8071.19



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2012-2014** LIC. NO.

----- COPY TOLERANCE

File Path / Date / Time Central Garden JOBS/Central Garden Jobs WIP/071190/071190_A/050_Production Art/MPLS_071190_A_Label_K.ai 12/15/2011 1:38:34 PM

Component #:	NA
Description:	Garden Tech RTU Insecticide & Miticide 32oz
UPC #:	N/A
Consumer Unit Size:	W: 4 IN H: 4.834 IN.
Program:	Adobe Illustrator CS4
Printer:	N/A
Creation Date:	12.04.2011
Rev. Date:	12.15.2011

Ink Colors

- Black
- Yellow
- Magenta
- Cyan
- PMS 361
- PMS 186

UPC Check

Initial _____ Date _____
Schawk Minneapolis

Numerical Scan Only Pass Fail

Proofread/QC

Proofreader _____ Date _____
Initial

Quality Control _____ Date _____
Initial

SCHAWK!

2626 2nd Street NE,
Minneapolis, MN
55418 USA
612 789 8514
schawk.com

The color on this copy will not match the final printed package.
Schawk Minneapolis has checked this artwork for accuracy. Final approval is the client's responsibility.
Please double check for errors before reproduction.

GARDENTECH Worry Free^{Brand}

READY TO USE



tent caterpillars



ticks



fleas



mites

**No mixing!
No Measuring!
Kills on Contact!**

PRODUCT FACTS

Kills	Over 250 types of insects — including Ants (Excluding Pharaoh and Carpenter Ants), Aphids, Beetles (including Japanese and Cucumber beetles), Caterpillars (including Armyworms, Cabbage Looper, Corn Earworm, Cutworm, Diamondback Moth, Hornworm, Imported Cabbageworm, Lesser Peachtree Borer, Oriental Fruit Moth, Peach Twig Borer, Tomato Fruitworm, Tomato Pinworm), Leafhoppers, Mites & other listed Insects
Where To Use	Vegetables, Fruits, Herbs, Spices, Roses, Flowers, Shrubs, Evergreen and Shade Trees
Amount To Use	Spray lightly covering upper and lower leaf surfaces.
When To Use	At the first signs of insect infestation. Can be used up to day of harvest. Kills during dormant stage.

Peel Back Tab for Additional Information

FOR OUTDOOR RESIDENTIAL USE ONLY

EPA Reg. No. 1021-1801-71004
EPA Est. 67572-GA-1a, 67572-GA-2a
Subscript used is first letter of run code on container.

Distributed by: TechPac, LLC.
2030 Powers Ferry Road, Ste 370
Atlanta, GA 30339



092711A-12/11

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **For information regarding medical emergencies or pesticide incidents, call 1-800-420-9347.**

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

2

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use this product in or on electrical equipment because of shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE EACH USE

USE INFORMATION

SHAKE WELL BEFORE USING.

The product may be used outdoors as a contact spray for flying and crawling pests commonly found in and around the garden and landscaping. This product contains Pyrethrum, derived from a Chrysanthemum flower. Pyrethrum will "flush" hiding insects. This product may be used throughout the growing season; or as a pre-harvest treatment. It is acceptable for use up to the day of harvest on edible plants. This product must come into contact with insects in order to kill them.

3

Phytotoxicity Note: Plant safety is an important consideration when using insecticides. However, it is not possible to evaluate the potential adverse effects of this product on numerous plant varieties that may react differently to insecticides in different growth stages or under varying environmental conditions. Before making widespread applications of this product treat a limited number of plants and observe for adverse effects over a one to 10 day period.

HOW TO APPLY: Wearing household latex or rubber gloves, use this ready-to-use bottle to apply product. Water plant thoroughly. Turn spray nozzle to the on position, and hold spray nozzle about 12" from the plants. Spray thoroughly covering, contacting insects on upper and lower leaf surfaces until surfaces are slightly wet. Avoid over-saturating. (Many insects hide on the underside of the leaf.) Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward. Allow liquid to dry in treated areas before re-entering.

Wash hands thoroughly after handling and before eating or smoking. Do not allow spray to drift to non-target plants or water supplies. Wash all fruits and vegetables before eating. Avoid contamination of food and feedstuffs.

WHEN TO APPLY: Use at first signs of insect infestation, preferably before damage and feeding occurs. Do not wait until plants are heavily infested. Spray may be applied up to and including day of harvest.

For Best Results: Apply in the twilight hours of the evening or just before sunrise to avoid possible harm to honey bees. Honey bees normally forage during the daylight hours, when the temperatures reach or exceed 55 degrees F.

WHERE TO USE: On Vegetables, Fruit and Nut Trees, Fruits, Nuts, Roses, Flowers, Ornamentals, Herbs, Spices, Shrubs, Bushes, Evergreens, Shade Trees. Do not use on lawns or large trees. For trees taller than 10 feet, consider hiring a licensed professional.

Use Restrictions

Aerial applications are prohibited.

Do not reapply within 3 days except under extreme pest pressure.

In case of extreme pest pressure, do not reapply within 24 hours.

Do not apply more than 1 time per day.

Do not apply when bees are active in the area.

Do not make applications during rain.

Do not apply if rain is expected within 24 hours.

Do not apply if wind causes drift.

This product is not for use on plants being grown for sale or other commercial use.

Do not allow adults, children, or pets to enter the treated area until sprays have dried.

Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift.

Do not water the treated area to the point of run-off.

Do not use treated article until spray has dried.

Do not wet plants to the point of runoff or drip.

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted use:

- Application to vegetation.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.

Do not allow the product to enter any drain during or after application.

INSECTS KILLED**On Fruit & Nut Trees:**

Apple Aphid, Caterpillars, Codling Moth, filbertworm, fruit fly, green fruitworm, leafhoppers, leafrollers, lesser peach tree borer, mired bug, navel orangeworm, oriental fruit moth, peach twig borer, pear psylla, plum curculio, rose chafer, spotted tentiform leafminer, tarnished plant bug, thrips, and walnut husk fly.

On Vegetables & Melons:

Aphids, Armyworms, Artichoke Plume Moth, Asparagus Beetle, Bean Beetles, Cabbage Looper, Colorado potato beetle, Corn Earworm, Corn Rootworm (adult), Cucumber Beetle (adult), Cutworms, Diamondback Moth, European Corn Borer, Flea beetles, Hornworms, Imported Cabbage Looper, Leaf Hoppers, Leaf Miners, Melon Worm, Onion Maggot (adult), Onion Thrips, Pepper Weevil, Pickleworm, Plantbugs (including Lygus, Squash, and Stink bugs), Potato Psyllid, Potato Tuberworm, Rindworm, Squash Vine Borer, Tomato Fruitworm, and Tomato Pinworm.

On Ornamentals (including Roses):

Asparagus beetle, Beet Armyworm, Cabbage Looper, Japanese Beetle, Lygus bug, and Omnivorous Leafroller.

On and Around Ornamental Roses, Flowers, Evergreens, Trees, Shade Trees, Bushes and Shrubs:

Adelgids, Ants (Acrobat, Argentine, Bigheaded, Cornfield, Crazy, Field, Ghost, Harvester, Odorous House, Pavement, and Thief; EXCLUDING Pharaoh and Carpenter ants), Aphids, Apple Aphids, Apple Maggots, Armyworms, Asian Cerambycid Beetles, Asian Lady Beetles, Asian Pine Tip Moths, Australian Sod Flies, Azalea Leafminers, Bagworms, Beetles, Billbugs, Birch Leafminers, Black Fungus Beetles, Black Turfgrass Ataenius (adult), Black Turpentine Beetles, Black Vine Weevils, Blister Beetles, Blow Flies, Blue Grass Billbug (adult), Boll Weevils, Boxelder Bugs, Boxwood Leafminers, Brown Dog TICKs, Brown Spider Beetles, Brown Spruce Longhorn Beetles, Browntail Moths, Budworms, Cadelles, California Oakworms, Campylomma Bugs, Cankerworms, Carrion Beetles, Catalpa Sphinx,

Caterpillar Ants, Caterpillars (including Tent), Catorama beetles, Chinch Bugs, Chrysanthemum Leaf Miners, Cicadas, Clearwing Borers, Climbing Cutworms, Clover Mites, Clover Weevils, Cockroaches (Smokey Brown, American, German), Codling Moths, Cone Beetles, Cone Worms, Conifer Spider Mites (adult), Corn Borers, Corn Earworms, Cottonwood Borers, Cottonwood Leaf Beetles, Crane Flies, Crickets, Cutworms (exposed), Dermesid Beetles, Diamondback Larvae, Dichondra Flea Beetles, Dingy Cutworm Adult, Douglass Fir Midge, Douglass Fir Tussock Moth, Dung Beetles, Earwigs, Eastern Tent Caterpillars, Elm Leaf Aphids, Elm Spanworms, Elmleaf Beetles, Emerald Ash Borer, Eriophyid Mites, European Pine Shoot Moths, European Pine Tip Moths, European Red Mites, Fall Armyworms, Fall Cankerworms, Fall Webworms, Field Crickets, Firebrats, Fireworms, Flatheaded Borers, Fleas, Flea Beetles, Forest Tent Caterpillars, Fruit Flies, Fruitree Leafrollers, Fuller Rose Beetles, Fungus Beetles, Fungus Gnats, Gall Midges, Gall Wasps, Giant Root Aphids, Glassy-winged Sharpshooters, Ground Beetles, Grasshoppers, Green Apple Aphids, Green Bugs, Greenhouse Thrips, Greenhouse Whiteflies, Greenstriped Mapleworms, Grubs, Gypsy Moths (adult and larvae), Hackberry Nipplegall Makers, Hairy Fungus Beetles, Harlequin Bugs, Heliothis spp., Holly Budmoths, Holly Leafminers, Hoplia Beetles, Hornworms, Hyperodes Weevils (adult), Inchworms, Iris Borer, Jackpine Budworms, Japanese Beetles, Jeffery Pine Needleminers, June Beetles, Katydid, Lace Bugs, Lawn Moths (Sod Webworms), Leaf Tiers, Leafhoppers, Leafminers, Leafrollers, Lesser Peach Tree Borers, Locust Borers, Loopers, Love Bugs, Lygus Bugs, Madrone Thrips, Maggots, Maple Leafcutters, McDaniel Spider Mites, Mealybugs, Millipedes, Mimosa Webworms, Plant Mites, Mole Crickets (nymphs and young adults), Moths Murrmidus Beetles, Mushroom Flies, Nantucket Pine Tip Moths, Oak Leaf Skeletonizers, Oak Leafminers, Oakworm complex, Oleander Caterpillars, Olive Ash Borers, Orange Tortrix, Orangestriped Oakworms, Pacific Mites, Palmetto Bugs, Peach Twig Borer, Pear Psylla, Pear Sawflies, Pearleaf Blister Mites, Periodical Cicadas, Phorid Flies, Pillbugs, Pine Bark Beetles, Pine Engraver, Pine Sawflies, Pine Spittlebugs, Pink Scavenger Caterpillars, Pinworms, Pitch Pine Tip Moths, Plant Bugs, Plum Curculia, Poinsettia Hornworms, Ponderosa Pine Needle Miner, Psyllids, Puss Caterpillars, Raspberry Horntail, Red Gum Lerp Psyllid, Red Lilly Beetles, Red Oak Borers,

Redbanded Leafrollers, Redhumped Oakworms, Rootworms, Rose Aphids, Rose Chafers, Rose Curculios, Rose Leafhoppers, Rose Slugs, Rosy Apple Aphids, Royal Palm Bugs, Saddled Prominent Caterpillar, Sawflies (exposed), Scale (Crawlers), Sciarid Flies, Seedbugs, Silverfish, Small Fruit Flies, Southern Pine Beetles, Southern Red Mite (adult), Southern Chinch Bugs, Sowbugs, Spider Beetles, Spider Mites, Spiders, Spiny Elm Caterpillars, Spittlebugs, Springtails, Spruce Budworms, Spruce Needleminers, Stalk Borers, Strawberry Mites, Subtropical Pine Tip Moths, Sweet Potato Whiteflies, Sycamore Leaf Beetle, Tarnished Plant Bug, Tent Caterpillars, Thorn Bugs, Thrips (exposed), Ticks, Trogoderma Beetles, Two-spotted Spider Mites, Viburnum Leaf Beetles, Walnut Caterpillars, Wasps, Waterbugs, Webworms (including Garden, Sod, Tropical), Weevils, Western Flower Thrips, Western Hemlock Loopers, Western Pine Tip Moth, Western Spruce Budworms, White Apple Leafhoppers, White Apple Leafrollers, White Grubs (Japanese Beetle, European Chafer, Southern Chafer), Willow Leaf Beetles, Woolly Apple Aphids, and Yellow Poplar Weevils.

FOR USE ON ROSES, FLOWERS, TREES, SHRUBS, AND VINES:

Vegetables (including in home greenhouses): Arrugula, Beets, Broccoli, Brussels Sprouts, Cabbage, Carrots, Chinese Cabbage (Bok Choy), Chinese Mustard Cabbage (Gai Choy), Cabbage (Napa), Cauliflower, Cucumbers, Collards, Cilantro, Celery, Eggplant, Edible Gourds, Endive, Japanese Radish (Daikon), Lettuce (Head and Leafy), Okra, Melons (including hybrids-Cantaloupe, Casaba, Crenshaw, Honeydew, Muskmelon), Parsley, Parsnips, Potato, Peppers (including Bell, Chili, Cooking, Pimentos, Sweet), Pumpkins, Rhubarb, Rutabaga, Radishes, Swiss Chard, Squash (Winter and Summer), Spinach, Tomatillo, Tomatoes, Watermelons, and Zucchini.

Small Fruits and Berries: Blackberries, Blueberries, Currants, Dewberries, Elderberries, Gooseberries, Grapes, Huckleberries, Olallie Berries, Raspberries (Black and Red), Strawberries, and Youngberries.

Fruits: Apples, Apricots, Crabapples, Cherries (Sweet and Sour), Citrus Hybrids, Grapefruit, Loquat, Lemons, Limes, Nectarines, Oranges (Sweet and Sour), Oriental Pears, Pears, Pummelos, Peaches, Plums, Prunes, and Quince.

For Use on All Ornamental Plants such as Roses, Hyacinth, Gladiolus, Freesia. All Annuals such as Cosmos, Celosia, Ageratum, Impatiens. All Perennials such as Gazania, Cineraria, Chrysanthemum, Zinnia. All Shrubs such as Mountain Laurel, Juniper, Privet, Mock Orange. All Shade Trees such as Oak, London Plane, Maple, Sycamore, Mulberry, and All Vines such as Trumpet Creeper, Creeping Lantana, Wisteria, Honeysuckle.

For the Control of: Ants, Aphids, Armyworms, Asparagus Beetles, Blister Beetles, Brown Marmorated Stink Bugs (BMSB), Cabbage Loopers, 12-Spotted Cucumber Beetles, Colorado Potato Beetles, Corn Earworms, Crickets, Cross-striped Cabbageworms, Diamondback Moth Larvae, Fireworms, Flea Beetles, Fruit Worms, Fruit Tree Leafrollers, Grape Leafhoppers, Gypsy Moths (larvae and adults), Harlequin Bugs, Tomato Hornworms, Imported Cabbageworms, Leafhoppers, Leafrollers, Leaf-tiers, Mexican Bean Beetles, Potato Leafhoppers, Psyllids, Stink Bugs, Thrips, Vinegar Flies, and Webworms: Water plant thoroughly. Turn spray nozzle to the on position, and hold spray nozzle about 12" from the plants. Spray thoroughly covering, contacting insects on upper and lower leaf surfaces until surfaces are slightly wet. Do not over-saturate. Do not water the treated area to the point of run-off. Spray when insects or their damage first appears. Do not wait until plants are heavily infested. Spray may be applied up to and including day of harvest.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage

Store in a cool, dry area. Always store pesticides in the original container. Store in an upright position only. Store with spray nozzle securely closed. Do not store with spray nozzle below level of product in container. Store away from food and pet food.

Pesticide Disposal and Container Handling

Nonrefillable container. Do not reuse or refill this container.

If empty:

Place in trash or offer for recycling if available.

If partly filled:

Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Sevin® is a registered trademark of Bayer.

GardenTech® is a registered trademark of Gulfstream Home and Garden, Inc.

Worry Free® Brand is a registered trademark of Central Garden & Pet.

If you have questions relating to product use or application, please call


866-945-5033.

If you have a product spill or transportation emergency, please call **800-424-9300**.

For Material Safety Data Sheets, please call **866-945-5033**.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

NOTES



File Path / Date / Time Central Garden JOBS/Central Garden Jobs WIP/071190/071190_B/050_Production Art/MPLS_071190_B_Label_K.ai 12/09/2011 3:38:34 PM

Component #:	NA
Description:	Garden Tech RTU Insecticide & Miticide 32oz
UPC #:	6-13499-04002-2
Consumer Unit Size:	W: 4 IN H: 4.834 IN.
Program:	Adobe Illustrator CS4
Printer:	N/A
Creation Date:	12.04.2011
Rev. Date:	00.00.0000

Ink Colors
<input checked="" type="checkbox"/> Black

UPC Check		Proofread/QC	
Initial Schawk Minneapolis	Date	Proofreader Initial	Date
Numerical Scan Only	<input type="checkbox"/> Pass <input type="checkbox"/> Fail	Quality Control Initial	Date

SCHAWK! 2626 2nd Street NE,
Minneapolis, MN
55418 USA
612 789 8514
schawk.com

The color on this copy will not match the final printed package.
Schawk Minneapolis has checked this artwork for accuracy. Final approval is the client's responsibility.
Please double check for errors before reproduction.