



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2014-2016** LIC. NO.

8071.4



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2011-2013** LIC. NO.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2008-2010** LIC. NO.

KILLS JAPANESE BEETLES

GARDEN TECH

Sevin®

Ready-To-Spray

BUG KILLER

Japanese Beetles Cabbage Worms Ticks Ants

Kills Over 100 Listed Insects on Vegetables, Fruits, & Ornamentals

Net Contents 32 fl. oz. (946 ml)

KEEP OUT OF REACH OF CHILDREN
CAUTION
See Back Panel For Precautionary Statements

ACTIVE INGREDIENT
Carbaryl (1-naphthyl N-methylcarbamate), 22.5% by wt
INERT INGREDIENTS: 77.5% by wt.
Contains 2 Pounds Carbaryl per Gallon
EPA Reg. No. 264-334-71004 EPA Est. No. 67572-GA-1

file: 24177-3_front label
folder: 24177-3_TecPac_70TEC-032334-04
colors: 4cp, pms 072, pms 186, green match pms 356
size: 3.5" w x 5.75" h

created: 01-15-01 bm
alt: 04-07-08 kc

Sevin® GARDEN TECH Ready-To-Spray BUG KILLER

For Vegetable Insects



Japanese Beetles



Cutworms



Horn/Cabbage Worms



Leafhopper

Alfalfa Caterpillar, Armyworm, Aster Leafhopper, Bean Leaf Beetle, Blister Beetle, Colorado Potato Beetle, Cucumber Beetles, Cutworms, Diamondback Moth, European Corn Borer, Fall Armyworm, Flea Beetles, Grape Colaspis, Green Cloverworm, Harlequin Bug, Imported Cabbage Worm, Japanese Beetle, Lace Bugs, Leafhoppers, Melonworm, Mexican Bean Beetle, Pickleworm, Rednecked Peanutworm, Spittlebugs, Squash Bugs, Sweet Potato Weevil, Tarnished Plant Bugs, Three Cornered Alfalfa Hopper, Thrips, Tomato Fruitworm, Tomato Hornworm, Tomato Pinworm, Tortoise Beetles, Webworms, Western Bean Cutworm, & many more.

For Ornamental Insects



Moths



Grasshoppers



Tent Caterpillars



Japanese Beetles



Cicadas

Ticks, Ants, Azalea Leafminer, Blister Beetle, Brown Tail Moth, Cutworms, Elm Leaf Aphid, Eriophyid Mites, Fall Armyworm, Fuller Rose Beetle, Plant Bugs, Scale Insects (crawlers), Tent Caterpillars, & many more listed inside.

Distributed by:
TechPac, LLC
P.O. Box 24830
Lexington, KY 40524



Peel Back Tab for Additional Information

created: 09-02-98 ds
alt: 04-07-08 kc
size: 3" W X 5.75" H

file: 6062-6_book art
folder: 6062-6_TecPac_707EC-032334-05
colors: 4 cp, pms186
barcode: upca, 80%, -39

FIRST AID

IF SWALLOWED:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.

**For MEDICAL Emergencies Call 24 Hours A Day 1-800-420-9347.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.**

NOTE TO PHYSICIAN: Carbaryl is a N-methyl carbamate that inhibits cholinesterase. Atropine is antidotal. Do not use 2-PAM (pyridine-2-aldoxime methiodide), opiates or cholinesterase inhibiting drugs. Probable mucosal damage may contraindicate the use of gastric lavage. To aid in the confirmation of diagnosis, urine samples should be obtained within 24 hours of exposure and immediately frozen. Analysis will be arranged by TechPac, LLC. Consultation on therapy can be obtained at all hours by calling the appropriate emergency number 1-800-420-9347 (U.S.A.).

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (& DOMESTIC ANIMALS) CAUTION

HARMFUL IF SWALLOWED, ABSORBED THROUGH THE SKIN OR INHALED. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Remove contaminated clothing and wash clothing before reuse.

USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when mixing and applying this product.
- Wear a hat and eye protection when making overhead applications.
- Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Avoid contamination of food and feed.
- Do not allow children or pets to contact treated area until sprays have dried.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters.

BEE CAUTION: MAY KILL HONEYBEES IN SUBSTANTIAL NUMBERS. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops and weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Contact your Cooperative Agricultural Extension Service for further information.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Strictly observe label directions and cautions. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. The area being treated must be vacated by unprotected persons. Keep unprotected persons out of treated areas until sprays have dried. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Shake all containers prior to use. Do not reuse empty containers or measuring devices for other purposes. Apply when insects or damage appear. Repeat as necessary to maintain control, unless spray interval is specified.

Apply when air is calm to avoid drift and contact with eyes and skin.

Start spraying at the farthest corner of the treatment area and work backward to avoid contact with wet surfaces. Allow spray to dry in treated areas before reentering.

For trees taller than 10 feet, consider hiring a licensed professional.

Spray thoroughly to wet upper and lower leaf surfaces, stems and branches.

Do not repeat applications more than once a week.

STORAGE AND DISPOSAL

Do not contaminate food or feed by storage or disposal

STORAGE:

Store unused product in original container only. In cool, dry area out of reach of children and animals, preferably in a locked storage area. Do not store in areas where temperatures frequently exceed 100°F. After use, replace cap on insecticide container.

DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.


HOSE END SPRAYER INSTRUCTIONS

Shake before using.

HOW TO USE:



1. Attach garden hose to spray nozzle.
2. Turn on water supply.



4. Point nozzle toward plants and turn water control knob to "On". Product will automatically mix with water. Spray upper and lower leaf surfaces thoroughly.



3. Bend safety tab down and turn control to "water".



GENERAL CAUTIONS AND RESTRICTIONS**PLANT RESPONSE PRECAUTIONS**

Application to wet foliage or during periods of high humidity may cause injury to tender foliage.

Do not use on Boston Ivy, Virginia creeper and maidenhair fern as injury may result. Carbaryl may also injure Virginia and sand pines.

The use of adjuvants may increase the potential for crop injury to sensitive crops.

NOTE: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics, and carpet. Spray deposits on painted or stained surfaces or finishes (i.e., cars, houses, trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid applications to surfaces where visible spray residues are objectionable. Staining of fabrics and carpet may be permanent.

VEGETABLES EXCEPT ASPARAGUS AND SWEET CORN

CROP	PEST	FL. OZ. OF THIS PRODUCT PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Broccoli Brussels Sprouts Cauliflower Cabbage Celery Chinese Cabbage Collards Cowpeas Cucumbers Dandelion Eggplant Endive	Alfalfa caterpillar Armyworm Aster leafhopper Bean leaf beetle Blister beetle Colorado Potato beetle Cucumber beetles Cutworms Diamondback moth	European corn borer Fall armyworm Flea beetles Grape colapsis Green cloverworm Harlequin bug Imported cabbageworm Japanese beetle	1 1/2 For optimum control of squash bugs time sprays for early morning or late afternoon. When disease transmission is suspected, monitor plants following application and retreat if reinfestation occurs. For cutworm control, this product is most effective against species which feed on the upper portions of the plant. For optimum control of thrips, ensure adequate coverage for the underside of leaves.

(continued)

VEGETABLES EXCEPT ASPARAGUS AND SWEET CORN *(continued)*

CROP	PEST	FL. OZ. OF THIS PRODUCT PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Fresh and Dried Beans Beans (<i>Phaseolus</i> species including snap, navy and kidney) Fresh and Dried Peas (<i>Pisum</i> species) Kale Kohlrabi Lettuce (head and leaf) Lentils Melons Mustard Greens Parsley Peppers Pumpkins Tomatoes	Lacebugs Leafhoppers Melonworm Mexican bean beetle Pickleworm Rednecked peanutworm Spittlebugs Squash bugs Sweet potato hornworm Sweet potato weevil Tarnished plant bugs Three cornered alfalfa hopper Thrips Tomato fruitworm Tomato hornworm Tomato pinworm Tortoise beetles Velvetbean caterpillar Webworms Western bean cutworm Whitefringed beetle adults	1 1/2	For optimum control of squash bugs time sprays for early morning or late afternoon. When disease transmission is suspected, monitor plants following application and retreat if reinfestation occurs. For cutworm control, this product is most effective against species which feed on the upper portions of the plant. For optimum control of thrips, ensure adequate coverage for the underside of leaves.

(continued)

VEGETABLES EXCEPT ASPARAGUS AND SWEET CORN *(continued)*

CROP	PEST	FL. OZ. OF THIS PRODUCT PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Root and Tuber Crops: Garden Beets, Carrots, Horseradish, Parsnips, Radishes, Rutabagas, Salsify, Potatoes Southern Peas Spinach Squash Sweet Potatoes Swiss Chard		1 1/2	For optimum control of squash bugs time sprays for early morning or late afternoon. When disease transmission is suspected, monitor plants following application and retreat if reinfestation occurs. For cutworm control, this product is most effective against species which feed on the upper portions of the plant. For optimum control of thrips, ensure adequate coverage for the underside of leaves.

RESTRICTIONS AND PRECAUTIONS: VEGETABLES

- For Broccoli, Brussel Sprouts, Cauliflower, Cabbage, Cucumbers, Eggplant, Kohlrabi, Melons, Peppers, Pumpkins, Tomatoes and Squash do not apply within 3 days of harvest.
- For Fresh Beans and Peas, do not apply within 3 days of harvest. For Dried Beans and Peas, do not apply within 21 days of harvest.
- For Root and Tuber Crops and Sweet Potatoes do not apply within 7 days of harvest.
- For Celery, Chinese Cabbage, Collards, Dandelion, Endive, Kale, Lettuce (head and leaf), Mustard Greens, Parsley, Spinach and Swiss Chard do not apply within 14 days of harvest.
- For all vegetables do not make more than a total of 4 applications per year.

RESTRICTIONS AND PRECAUTIONS: VEGETABLES (continued)

- For all vegetables do not apply more than a total of 9 fluid ounces per 1000 sq.ft. per year.
- Do not apply more often than once every 7 days.

ASPARAGUS

CROP	PEST	FL. OZ. OF THIS PRODUCT PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Asparagus	Apache cicada Asparagus beetle	3	Repeat applications as necessary up to a total of 3 times prior to harvest or a total of 5 times per crop but not more often than once every 7 days. For cutworm control, this product is most effective against species which feed on the upper portions of the plant.
	Cutworms		
		6	Application to ferns or brush growth following harvest of spears: Repeat applications as necessary but not more often than once every 7 days. Do not make more than a total of 5 applications per year to spears and ferns combined.

RESTRICTIONS AND PRECAUTIONS: ASPARAGUS

- Do not apply within 1 day of harvest.
- Do not apply more than a total of 9 fluid ounces per 1000 SQ. FT. before harvest of spears.
- Do not apply more than a total of 15 fluid ounces per 1000 SQ. FT. per year.

SWEET CORN

CROP	PEST	FL. OZ. OF THIS PRODUCT PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Sweet Corn	Armyworm Chinch bugs Corn earworm Corn rootworm adults Fall armyworm	Flea beetles Japanese beetle Sap beetles Southwestern corn borer Leafhoppers	3 OBSERVE BEE CAUTION Repeat applications as necessary up to a total of 8 times but not more often than once every 7 days. Optimum timing and good coverage are essential for effective control.
	European corn borer		3
	Western bean cutworm Cutworms		3

RESTRICTIONS AND PRECAUTIONS: SWEET CORN

- Do not apply within 2 days of harvest of ears or within 14 days of harvest or grazing of forage.
- Do not apply more than a total of 24 fluid ounces per 1000 SQ. FT. per crop.

TREE FRUIT CROPS

On all tree fruit crops, apply in sufficient volume for adequate coverage. This will vary depending on the pest and its severity, the tree condition, size, and density, and other factors.

CITRUS FRUITS

CROP	PEST	FL. OZ. OF THIS PRODUCT PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Citrus Fruits	Apopka weevil (adult) Avocado leafroller California orangedog Citrus cutworm Citrus root weevils (adults) Citrus rust mite Eriophyid mites Fruitree leafroller Fuller Rose Beetle Little leaf notcher (adult)	Orange Tortrix Plant bugs Scale insects (Black scale, brown soft scale, California red scale (except in California), citrus snow scale, yellow scale) Western tussock moth	OBSERVE BEE CAUTION. Repeat applications as necessary up to a total of 8 times but not more often than once every 14 days. Use lower rates for pests attacking leaves. Use higher rates for pests attacking fruit and for higher infestations. For scale control, apply when crawlers are present. For best control of Eriophyid mites including citrus rust mite, apply when pest populations are low.
	California only: California red scale	7 1/2	Do not make more than 1 application per season. Apply when crawlers are present.

(continued)

RESTRICTIONS AND PRECAUTIONS: CITRUS FRUITS (continued)

- For trees taller than 10 feet, consider hiring a licensed professional.
- Do not apply within 5 days of harvest.
- Do not apply more than a total of 30 fluid ounces per 1000 SQ. FT. per crop

STONE FRUIT AND POME FRUIT

CROP	PEST	FL. OZ. OF THIS PRODUCT PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Stone Fruits: Apricots, Cherries, Nectarines, Peaches, Plums, and Prunes Pome Fruits: Apples, Pears, Loquats, Crabapples, and Oriental Pears.	Apple aphid Apple maggot Apple mealybug Apple pandemis Apple rust mite Bagworms Black cherry aphid California pearslug (pear sawfly) Cherry fruitworm Cherry maggot (Cherry fruit fly) Codling moth Cucumber beetles Eastern tent caterpillar Eyespotted bud moth European apple sawfly European earwig Fruittree leafroller Green fruitworm Gypsy moth Japanese beetle June beetle Lesser appleworm Lesser peachtree borer Lygus bugs Mealy plum aphid Orange tortrix	4 1/2	OBSERVE BEE CAUTION. Repeat applications as necessary up to a total of 3 times per crop but not more often than once every 7 days. An additional application at the dormant or delayed dormant timing may be made. For optimum scale control, apply when crawlers are present. For lesser peachtree borer, best results have been found by thoroughly spraying limbs and tree trunks at weekly intervals during moth flight. For psylla control, apply when eggs hatch or young nymphs are present.

(continued)

STONE FRUIT AND POME FRUIT (continued)

CROP	PEST	FL. OZ. OF THIS PRODUCT PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Stone Fruits: Apricots, Cherries, Nectarines, Peaches, Plums, and Prunes Pome Fruits: Apples, Pears, Loquats, Crabapples, and Oriental Pears. (continued)	Oriental fruit moth Peach twig borer Pearleaf blister mite Pear psylla Pear rust mite Periodical cicada Plum curculio Prune leafhopper Redbanded leafroller Rose chafer Rosey apple aphid Scale insects (Brown soft scale, Forbes scale, Lecanium scale Olive scale, Oystershell scale, San Jose scale) Tarnished plant bug Tentiform leafminers Tussock moth Variegated leafroller White apple leafhopper Woolly apple aphid Yellowheaded fireworm	4 1/2	OBSERVE BEE CAUTION. Repeat applications as necessary up to a total of 3 times per crop but not more often than once every 7 days. An additional application at the dormant or delayed dormant timing may be made. For optimum scale control, apply when crawlers are present. For lesser peachtree borer, best results have been found by thoroughly spraying limbs and tree trunks at weekly intervals during moth flight. For psylla control, apply when eggs hatch or young nymphs are present.

(continued)

RESTRICTIONS AND PRECAUTIONS: STONE FRUIT AND POME FRUIT *(continued)*

- For trees taller than 10 feet, consider hiring a licensed professional.
- Do not apply within 3 days of harvest.
- Do not apply more than a total of 21 fluid ounces per 1000 SQ. FT. per crop.
- Do not apply more than a total of 13 1/2 fluid ounces per 1000 SQ. FT. during the production season.

TREES AND ORNAMENTALS

CROP	PEST	FL. OZ. OF THIS PRODUCT PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Trees, Ornamentals (Roses, Flowers, Other Herbaceous Plants), Woody Plants and Shrubs	Ants Apple aphid Armyworm Ash whitefly Azalea leafminer Bagworms Balsam twig aphid Birch leafminer Blister beetle Boxelder bug Boxwood leafminer Brown tail moth Cankerworms Catalpa sphinx Chiggers Cooley spruce gall adelgid Cutworms Cypress tip moth Douglas-fir tussock moth Eastern spruce gall adelgid Elm leaf aphid Elm leaf beetle Elm spanworm Eriophyid mites European pine shoot moth Fall armyworm Flea beetle	1 1/2	Observe directions on applicator label. Do not apply more than once every 7 days.

(continued)

TREES AND ORNAMENTALS *(continued)*

CROP	PEST	FL. OZ. OF THIS PRODUCT PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Trees, Ornamentals (Roses, Flowers, Other Herbaceous Plants), Woody Plants and Shrubs <i>(continued)</i>	Fuchsia gall mite Fuller rose beetle Gall midges Gall wasps Greenstriped mapleworm Grasshoppers Gypsy Moth Hackberry nipplegall maker Holly bud moth Holly leafminer Jackpine budworm Japanese beetle Jeffrey pine needleminer June beetles Lace bugs Leafhoppers Leafrollers Locust borer Maple leafcutter	Mealy bugs Mimosa webworm Nantucket pine tip moth Oak leafminers Oak moth Oak skeletonizer Oakworm complex Oleander caterpillar Olive ash borer Orange-striped oakworm Periodical cicada Pine looper Pine sawfly Pine spittlebug Pitch pine tip moth Spruce budworm Plant bugs	1 1/2 Observe directions on applicator label. Do not apply more than once every 7 days.

(continued)

TREES AND ORNAMENTALS *(continued)*

CROP	PEST	FL. OZ. OF THIS PRODUCT PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Trees, Ornamentals (Roses, Flowers, Other Herbaceous Plants), Woody Plants and Shrubs <i>(continued)</i>	Poinsettia hornworm Psyllids Puss caterpillar Redhumped oakworm Rose aphid Rose chafer Rose slug Saddled prominent Sawflies (exposed) Scale insects (crawlers) Sowbugs Spiney elm caterpillar Springtails Spruce needleminer	Subtropical pine tip moth Tent caterpillars Thorn bug Trips (exposed) Ticks Walnut caterpillar Webworms Western hemlock looper Western spruce budworm Willow leaf beetles Woolly gall aphid Yellow poplar weevil	1 1/2 Observe directions on applicator label. Do not apply more than once every 7 days.

RESTRICTIONS AND PRECAUTIONS: TREES AND ORNAMENTALS

- For trees taller than 10 feet, consider hiring a licensed professional.

CONTROL OF SPECIFIC PESTS ACROSS MULTIPLE SITES

CROP/SITE	PEST	FL. OZ. OF THIS PRODUCT PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Ticks Which Vector Lyme Disease All crops/sites on this label	<i>Ixodes</i> spp. (Deer tick, Bear tick, Black legged tick) <i>Amblyomma</i> spp. (Lone star tick)	1 1/2 to 3 fl. oz. per 1000 sq. ft.	Use the high rate for heavy tick infestations. Use higher spray volumes for dense ground cover or heavy leaf litter. Target applications for nymphal control in late spring or early summer. Control of adult tick can be obtained with late summer and fall applications. Do not use spot treatments. Treat entire area and perimeter areas where exposure to ticks may occur. Ticks may be reintroduced from surrounding areas on host animals. Retreat as necessary but not more often than once every 7 days, to maintain adequate control levels.

(continued)

CONTROL OF SPECIFIC PESTS ACROSS MULTIPLE SITES *(continued)*

CROP/SITE	PEST	FL. OZ. OF THIS PRODUCT PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Imported Fire Ants All crops/sites on this label	Imported fire ants	1 1/2 fl. oz. per mound.	MOUND DRENCH APPLICATION: Apply approximately 2 gallons of the hose-end spray solution over the surface of the mound. Thoroughly wet mound and surrounding areas to a 4 ft diameter (12 sq. ft.). Do not disturb mound prior to treatment. Direct flow at mound to disrupt and allow solution to flow into ant tunnels. For best results apply in cool weather (65-80°F) or in early morning or late evening hours. Repeat application if mound activity resumes after 7 days. Treat new mounds as they appear.
Adult Mosquito Control All crops/sites on this label	Mosquitoes (adults)	1 1/2 fl. oz. per 1000 sq. ft.	OBSERVE BEE CAUTION. Treat shrubbery and areas where adult mosquitoes congregate. Treat when adult mosquitoes are active in early mornings or late evenings. Repeat applications as necessary but not more often than once every 7 days.

(continued)

CONTROL OF SPECIFIC PESTS ACROSS MULTIPLE SITES *(continued)*

CROP/SITE	PEST	FL. OZ. OF THIS PRODUCT PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Nuisance Pest Control	Ants Bees Brown dog tick Centipedes Cockroaches Crickets Darkling beetle (lesser mealworm or litter beetle) Earwigs Firebrats Fleas Millipedes Scorpions Silverfish Spiders Wasps	12 fl. oz. per 1000 sq. ft. as a perimeter treatment	PERIMETER TREATMENT DIRECTIONS: Observe plant response precautions. Apply as a coarse wet spray in a band 6 to 10 feet wide around the outside perimeter of buildings. Confine applications to areas immediately adjacent to the building. Direct application to the structure should be minimal and restricted to cracks, crevices, and areas where insects tend to congregate. For use in and around flower beds and ornamental plantings, apply as a coarse wet spray in and around plant beds. May be applied to outdoor sleeping quarters of pets. Do not treat animals. Do not allow animals to use treated area until sprays have dried. Do not treat more often than once every 7 days.

RESTRICTIONS AND PRECAUTIONS:

- DO NOT ALLOW PUBLIC USE OF TREATED AREAS DURING APPLICATIONS OR UNTIL SPRAYS HAVE DRIED.
 - NOTE: Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics, and carpet. Spray deposits on painted or stained surfaces of finishes (i.e., cars, house trailers, boats, etc.) should be removed immediately by washing to prevent discoloration. Avoid application to surfaces where visible spray residues are objectionable. Staining of fabrics and carpet may be permanent.
- CAUTION: May kill shrimp and crabs. Do not use in areas where these are important resources.

Important: Read Before Use

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of TechPac, LLC. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TECHPAC, LLC. MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of TechPac, LLC. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TECHPAC, LLC. DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT TECHPAC, LLC.'S ELECTION, THE REPLACEMENT OF PRODUCT.

If you have a medical emergency, need consultation on diagnosis, therapy and analysis please call (800) 420-9347.

If you have questions relating to product use or application, please call (800) 969-7200.

If you have a product spill or transportation emergency call (800) 424-9300.

For Material Safety Data Sheets please go to www.gardentech.com, or call (800) 969-7200.
Sevin® is a registered trademark of Bayer for Carbaryl Insecticide.
Garden Tech® is a registered trademark of Gulfstream Home & Garden, Inc.