T-12740

Department of Agriculture
STATE OF HAWAII
LICENSED
PERIOD 2014-2016 LIC. NO.
8079.107

Department of Agriculture
STATE OF HAWAII
LICENSED
PERIOD 2011-2013 LIC. NO.

18" LENGTH

F
GRAPHICS ORIENTATION

3" GUSSET

18" LENGTH

B

Sta-Green

PHOSPHORUS - FREE
WEED & FEED
Controls a wide spectrum of Broadleaf weeds

Use on lawns to kill the toughest weeds like clover and dandelion
Feeds for 3 months guaranteed!*

COVERS 5,000 SQ FT

Kills over 250 Weeds
as listed

Sta-Green®

PHOSPHORUS - FREE

WEED & FEED

Controls a wide spectrum of Broadleaf weeds

Controla una amplia variedad de malezas de hoja ancha

Use on lawns to kill the toughest weeds like clover and dandelion

Feeds for 3 months guaranteed!*

Utilice en céspedes para eliminar malezas resistentes como tréboles y dientes de león ¡Nutrición garantizada durante 3 meses!*

- Kills the root
  Mata la raíz
- Starts working on contact
  Comienza a funcionar al contacto
- Quick greening and extended feeding*
  Proporciona un color verde rápido y una alimentación continua*

ACTIVE INGREDIENTS:

2,4-D, 2-ethylhexyl ester	1.108%
Mecoprop-p acid	0.167%
Dicamba acid	0.071%
OTHER INGREDIENTS:	98.654%
TOTAL	100.000%

THIS PRODUCT CONTAINS:
14.70 lb. 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735%. CAS 1928-43-4.
3.34 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.167%. CAS 16484-77-8.
1.42 lb. 3,6-dichloro-o-anisic acid equivalent per ton or 0.071%. CAS 1918-00-9.
Isomer Specific by AOAC Methods

KEEP OUT OF REACH OF CHILDREN
CAUTION
See Back Panel For First Aid And Additional Precautionary Statements

COVERS 5,000 SQ FT
MONEY BACK GUARANTEED† • NET WT 13 LB (5.90 KG)

28-0-4
SEE BACK PANEL FOR ANALYSIS DESCRIPTION

COVERS 5,000 SQ FT

Use on lawns to kill the toughest weeds like clover and dandelion
Feeds for 3 months guaranteed!*

PHOSPHORUS - FREE
WEED & FEED
Controls a wide spectrum of Broadleaf weeds

Sta-Green

Sta-Green®

PHOSPHORUS - FREE
WEED & FEED
Controls a wide spectrum of Broadleaf weeds

Kills over 250 Weeds
as listed

COVERS 5,000 SQ FT • NET WT 13 LB (5.90 KG)

WITH Pennington Coated® TECHNOLOGY
GREENS QUICKLY, FEEDS FOR MONTHS

Pennington Coated – Slow Release Technology
*For a Long Lasting Green Lawn With No Burning**

Sta-Green®

Pennington Coated – Slow Release Technology
*For a Long Lasting Green Lawn With No Burning**

WITH Pennington Coated® TECHNOLOGY
GREENS QUICKLY, FEEDS FOR MONTHS

PHOSPHORUS FREE – Helping Our Water Sources

28 N nitrogen – 0 P phosphorus – 4 K potassium
For lush, green plant growth
For overall plant health and vigor

GUARANTEED ANALYSIS F525 **28-0-4**

Total Nitrogen (N)	28.00%
27.13% Urea Nitrogen*	
0.87% Ammoniacal Nitrogen	
Soluble Potash (K_2O)	4.00%
Sulfur (S)	0.99%
0.99% Combined Sulfur (S)	

Derived from: Urea, Polymer Coated Urea, Ammonium Sulfate, Muriate of Potash.
*Part of the nitrogen in this product has been polymer coated to provide 5.70% coated slow release nitrogen.

STOP READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If in Eyes:
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:
- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

INSTRUCTIONS: Sta-Green Weed and Feed is a combination of three proven weed killers -- 2,4-D, mecoprop-p, and dicamba. This herbicide controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 28-0-4 fertilizer provides a supply of plant nutrients.

USE PRECAUTIONS

Do not use on dichondra, carpetgrass, St. Augustinegrass, creeping bentgrass, or turfgrass where desirable clovers are present.
Do not apply this product to newly seeded grasses until well-established or after the second or third mowing.
Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area and do not apply this product within the dripline of, or over surface roots of trees, shrubs, and other ornamentals.
Reseeding or overseeding areas may occur three (3) weeks after the application of this product.
Do not use this product on or near desirable plants and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of the roots of desirable trees and shrubs, since injury may result.
Do not apply directly to or near water, storm drains, gutters, sewers or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

HOW MUCH TO APPLY

For use on: Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, zoysiagrass, St. Augustinegrass, centipedegrass, hybrid bermudagrass, and colonial bentgrass. Apply 2.6 pounds of product per 1,000 square feet of turfgrass. A second application may be made in approximately 30 days. The maximum number of broadcast applications per treatment site is 2 per year.

This product controls the following list of growing weeds:

Amaranth, purple
Amaranth, slender
Aster, heath
Aster, white heath
Aster, purple
Aster, white prairie
Artichoke
Austrian fieldcress
Bedstraw, Catchweed
Bedstraw, smooth
Bedstraw, Northern
Bedstraw, pineywoods
Beggarticks, hairy
Betony, Florida
Bindweed, Field
Bindweed, Hedge
Bitter Wintercress
Bittercress, hairy
Bitterweed
Black-eyed Susan
Black Medic
Blood flower milkweed
Blue lettuce
Brassbuttons
Bristly oxtongue
Broomweed
Buckhorn
Bullnettle
Burclover
Burdock
Burweed, lawn (spurweed)
Buttercup, bulbous
Buttercup, creeping
Buttonweed, common (poorjoe)
Buttonweed, virginia
Carolina geranium (wild)
Carpetweed
Catnip
Catsear, spotted (false dandelion)
Chamber bitter
Chickweed, common
Chickweed, mouseear
Chickweed, sticky
Chicory
Cinquefoil
Clover, crimson
Clover, alyce
Clover, hop
Clover, rabbit foot
Clover, red
Clover, strawberry
Clover, yellow sweet
Clover, white sweet
Clover, white
Cockle
Cocklebur
Cocklebur, oriental
Cocklebur, spiny
Common mullein
Corn Spurry
Creeping Beggarweed
Creeping Jenny (creeping charlie)
Cupid shaving Brush
Cudweed
Daisy, English
Daisy, oxeye
Dandelion, common
Dandelion, false
Day flower
Deadnettle
Dichondra
Dock, broadleaf
Dock, curly
Dogbane
Dogfennel
Dollarweed (Pennywort)
Dove weed
Elderberry
Falseflax
False sunflower
Fiddleneck
Filaree, redstem
Filaree, whitestem
Fleabane, daisy (annual)
Fleabane, rough
Galinsoga, hairy
Galinsoga, smallflower
Goldenrod
Ground ivy
Groundsel
Gumweed
Hairy fleabane
Hawkweed, meadow
Hawkweed, mouseear
Hawkweed, orange
Hawkweed, yellow
Healall
Heartleaf drymary
Hemp
Henbit
Hoary cress
Horsenettle
Horseweed
Jimsonweed
Johnny-Jumpup violet
Knawel
Knotweed, prostrate
Kochia
Lambsquarters
Lespedeza
Mallow, alkali
Mallow, bristly
Mallow, common
Mallow, dwarf
Mallow, venice
Marcela
Matchweed
Mexicanweed
Moneywort
Morningglory, bigroot
Morningglory, ivyleaf
Morningglory, mexican
Morningglory, red
Morningglory, tall
Morningglory, wolly
Mugwort Mustard, ball
Mustard, black
Mustard, blue
Mustard, hedge
Mustard, indian,
Mustard, tansy
Mustard, tumble
Mustard, white
Mustard, wormseed
Nettle, stinging (burning)
Nettle, tall
Nettle, wood
Parsley-piert
Parsnip
Pearlwort
Pennycress (frenchweed)
Pepperweed, perennial
Pepperweed, virginia (Peppergrass)
Pigweed, prostate
Pigweed, redroot
Pigweed, russian
Pigweed, smooth
Pigweed, tumble
Pineapple weed
Plains coreopsis (tickseed)
Plantain, blackseed
Plantain, bracted
Plantain, broadleaf
Plantain, buckhorn
Plantain, hoary
Plantain, narrowleaf
Plantain, slender
Plantain, woolly
Poison ivy
Poison oak
Pokeweed
Prairie Sunflower
Prickly lettuce (compass plant)
Prickly sida
Prostrate knotweed
Puncturevine
Purslane, common
Pusley, brazil
Pusley, florida
Pusley, large flower
Ragweed, bur
Ragweed, common
Ragweed, lanceleaf
Ragweed, western
Redstem filaree
Rough cinquefoil
Scarlet pimpernel
Shepardspurse
Smartweed, ladysthumb
Smartweed, pale
Smartweed, pennsylvania
Smooth chaff-flower
Smoothdock
Sorrel, red (sheep)
Sowthistle, annual
Sowthistle, spiny
Spanish needles
Speedwell, bi-lobed
Speedwell, common
Speedwell, corn
Speedwell, ivyleaf
Speedwell, persian
Speedwell, purslane
Speedwell, slender
Speedwell, snow
Speedwell, thymeleaf
Spiny amaranth
Spurge, spotted
Spurge, prostrate
Spurweed
Star of Bethlehem
Strawberry, India mock
Tansy ragwort
Tanweed
Thistle, blessed
Thistle, bull
Thistle, canada
Thistle, flodman (prairie)
Thistle, musk
Thistle, platte
Thistle, plumeless
Thistle, russian
Thistle, scotch
Thistle, wavyleaf
Thistle, yellow (horrible)
Trailing crownvetch
Velvetleaf
Vervain, blue
Vervain, hoary
Vervain, prostrate
Vervain, tall
Vetch, bird
Vetch, hairy
Vetch, milk
Vetch, narrowleaf
Vetch, wild
Virginia creeper
Western clematis
Western salsify
Wild aster
Wild buckweat
Wild carrot
Wild four-o-clock
Wild garlic
Wild lettuce
Wild marigold
Wild mustard (yellow rocket)
Wild onion
Wild parsnip
Wild radish
Wild rape
Wild strawberry
Wild sweet potato
Wild violet
Woodsorrel, creeping
Woodsorrel, violet
Woodsorrel, yellow (Oxalis)
Woolly croton
Wormseed
Yarrow, common
Yarrow, western
Yarrow, yellow

WHEN TO APPLY

Apply when broadleaf weeds are actively growing in spring or early fall.
Apply when the air is calm to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery.
Possible retreatment may be necessary if lawn is heavily overrun by weeds, if adverse low moisture conditions prevail, or if weeds are in a state of poor growth.

APPLICATION INSTRUCTIONS

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
3. Apply when broadleaf weeds are young and actively growing, preferably in the morning when dew is on the grass. At the time of application, moisture on the weed leaves from dew, rainfall or irrigation/watering aids in control. If grass is not moist at time of application, sprinkle lightly with water before applying product.
4. Avoid applying when temperatures are consistently above 90°F.
5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

HOW TO APPLY – SPREADERS AND CALIBRATION

Always apply with a rotary or drop spreader to distribute the granules evenly. Never apply by hand.

Coverage – To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to 5000 square feet.

Important: It is important to follow directions as indicated on this quality lawn care product as over-application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owner's Manual. Never apply by hand.

SINGLE PASS DISTRIBUTION:
- For rotary spreaders: Operate at intervals corresponding to the swath width.
- For drop spreaders: Overlap the wheel tracks to ensure complete coverage.
- For best results, apply this product uniformly and avoid streaks, skips or overlaps. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping.

SPREADER SETTINGS
SPREADER SETTINGS(SPANISH)

BROADCAST SPREADER (BROADCAST SPREADER)		DROP SPREADER (DROP SPREADER)	
Ortho Rotary	1¾	Republic EZ	5
True Temper	3¾	Ortho Drop	5
EV-N-SPRED	14	Precision Drop	6
Precision Rotary	5	Scotts AccuGreen	6¾
Republic EZ Grow	5¾	Scotts AccuGreen 1000	6¾
Scotts EdgeGuard	5¾	Scotts AccuGreen 3000	6¾
Scotts Easy Green	24¾	Sta-Green DS4500	2
Scotts SpeedyGreen 1000	5¾		
Scotts SpeedyGreen 2000	5¾		
Scotts SpeedyGreen 3000	5¾		
Sta-Green SB5500	5½		
Sta-Green SB4500	5½		

CALIBRATION IINSTRUCTIONS AND SPREADER SETTINGS:
Calibrate the spreader according the the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended rate under actual application conditions.

To provide proper distribution, calibrate spreader before application.
• For 2.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 385 square feet and adjust to apply at the rate of 1 pound of product per 385 square feet.

BROADCAST SPREADERS: For best results, apply in a circular path starting on the outside and working toward the center overlapping slightly in insure complete coverage.

DROP SPREADERS: Apply 2 strips along each end. Then apply along length of lawn, overlapping slightly and shutting off the spreader before each turn at edge of border strips.

EPA Reg. No. 2217-884-59144
EPA Est. No. 59144-GA-1(A), 32802-MO-1(B), 37351-CO-1(C), 8378-IN-1(D), 9404-FL-1(E), 961-PA-01(F), 961-IL-01(G), 35497-OR-1(H), 49991-TX-2(Q)
Letter in parenthesis is first letter of lot number.

PHOSPHORUS FREE – Helping Our Water Sources

ALTO LEA LA ETIQUETA COMPLETA PRIMERO. CUMPLA CON TODAS LAS PRECAUCIONES Y SIGA LAS INSTRUCCIONES DETALLADAMENTE.

DECLARACIONES DE PRECAUCIÓN

RIESGOS PARA SERES HUMANOS Y ANIMALES DOMÉSTICOS

PRECAUCIÓN. Provoca una irritación moderada de los ojos. Evite el contacto con los ojos o la ropa. Lávese bien con agua y jabón después de manipular este producto y antes de comer, beber, masticar chicle, usar tabaco o usar el inodoro.

PRIMEROS AUXILIOS

Si entra en contacto con los ojos:
- Mantenga los ojos abiertos y enjuague lenta y suavemente con agua durante 15 a 20 minutos.
- Si lleva lentes de contacto puestos, quíteselos luego de los primeros 5 minutos, luego continúe enjuagando los ojos.
- Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.

Cuando llame a un centro de control de envenenamientos o a un médico, o busque tratamiento, tenga a mano el envase o la etiqueta del producto. También puede ponerse en contacto con el 1-800-424-9300 para obtener información de tratamiento médico de emergencia.

Riesgos ambientales

Este producto es tóxico para los peces e invertebrados acuáticos y puede tener efectos adversos en plantas que no eran el objetivo del tratamiento. Para proteger el medio ambiente, no deje que el pesticida entre o escurra hacia desagües pluviales, zanjas de desagüe, alcantarillas ni aguas superficiales. Aplicar este producto en un clima tranquilo, cuando no haya pronóstico de lluvia durante las siguientes 24 horas, ayudará a asegurar que el viento o la lluvia no soplen o laven el pesticida del área de tratamiento. Barra el producto que caiga sobre una entrada de garaje, acera o calle llevándolo de vuelta al área tratada del césped o jardín para evitar que se escurra hacia aguas superficiales o sistemas de desagüe.

Este producto químico tiene propiedades y características asociadas con agentes químicos detectados en agua del subsuelo. El uso de este producto químico en áreas donde el suelo es permeable, especialmente donde las napas de agua son de poca profundidad, puede ocasionar contaminación del agua del subsuelo. La aplicación alrededor de una cisterna o pozo puede ocasionar contaminación del agua potable o agua del subsuelo.

INSTRUCCIONES DE USO:

Se considera una violación a la Ley Federal usar este producto con fines que no sean los indicados en la etiqueta.

PRECAUCIONES Y RESTRICCIONES:
- No aplique el producto de ninguna forma que pueda entrar en contacto con personas o mascotas, ya sea en forma directa o indirecta. Mantenga a las personas y mascotas fuera del área durante la aplicación. No permita que las personas o mascotas ingresen al área tratada hasta que el polvo se haya asentado.
- No lo use en césped que se cultiva para la venta u otro uso comercial como tepe, para la producción comercial de semillas ni para fines de investigación.

INSTRUCCIONES: El herbicida y abono Sta-Green es una combinación de tres herbicidas comprobados: 2,4-D, mecoprop-p y dicamba. Este herbicida controla una gran variedad de malezas de césped, incluyendo dientes de león, pamplina, sanguinaria, llantén, ortiga muerta, hierba de golondrina y muchas otras que a menudo se encuentran en los céspedes domésticos. Además, un fertilizante 28-0-4 proporciona un suministro de nutrientes para las plantas.

PRECAUCIONES DE USO

No lo use sobre céspedes oreja de ratón, alfombra, San Agustín, agrostis o sobre céspedes donde haya tréboles que se desean conservar.
No aplique este producto sobre céspedes recién sembrados hasta que estén bien establecidos o después de una segunda o tercera poda.
No use este producto sobre flores, hortalizas, cubiertas vegetales, jardines o plantaciones, incluyendo arbustos y árboles. No exceda la dosis indicada para cada área y no aplique este producto dentro de la línea de goteo o sobre la superficie de las raíces de árboles, arbustos y otras plantas ornamentales.
Se puede volver a sembrar o sembrar varias veces después de tres (3) semanas desde la aplicación de este producto.
No use este producto en o cerca de plantas deseadas, y evite esparcir gránulos en sistemas de raíces expuestos o brotes espontáneos (rizomas) dentro de la línea de goteo de las raíces de los árboles y arbustos deseados, ya que podría provocar daños.
No lo aplique directamente en agua, desagües de aguas pluviales, canaletas, cloacas o zanjas de desagüe ni cerca de éstos. No lo aplique a menos de 7,62 metros de ríos, estanques de peces, lagos, arroyos, embalses, pantanos, estuarios, bahías y océanos. No lo aplique bajo condiciones de viento. Aplique este producto directamente en el césped y barra el producto que caiga sobre la entrada de garaje, la acera o la calle llevándolo de vuelta al césped. Para evitar que el producto escurra, no riegue en exceso las áreas tratadas ni aplique cuando esté lloviendo o cuando se esperen lluvias dentro del día.

CUÁNTO APLICAR

Para usar en: Césped azul de Kentucky, festuca spp., ballico perenne, césped Bermuda común, césped zoysia, césped de San Agustín, césped ciempiés, césped Bermuda híbrido y césped agrostis colonial. Aplique 1,18 kg del producto por cada 92,90 m^2 de césped. Se puede realizar una segunda aplicación en aproximadamente 30 días. El número máximo de aplicaciones por dispersión por sitio de tratamiento es de 2 por año.

CUÁNDO APLICAR

Aplique cuando las hierbas de hojas anchas estén en pleno crecimiento en primavera o a principios del otoño.
Aplique cuando no haya brisa para asegurar una cobertura uniforme y evitar que se esparzan los gránulos sobre flores, hortalizas y arbustos ornamentales.
Podría ser necesario un nuevo tratamiento si el césped está excesivamente invadido por malezas; si prevalecen condiciones adversas de humedad baja o si las malezas están en un período de crecimiento pobre.

INSTRUCCIONES PARA LA APLICACIÓN

1. Corte el césped a altura normal 1 a 2 días antes de la aplicación.
2. Riegue bien el césped al menos 1 a 2 días antes de la aplicación para mantener la humedad hasta el siguiente riego (consulte el paso 5).
3. Aplique cuando las malezas de hojas anchas estén jóvenes y en pleno crecimiento, de preferencia por la mañana cuando hay rocío sobre el césped. En el momento de la aplicación, la humedad en las hojas de las hierbas proveniente del rocío, lluvia o irrigación/regado ayuda en el control. Si el césped no está húmedo en el momento de la aplicación, rocíelo ligeramente con agua antes de aplicar el producto.
4. Evite aplicar el producto cuando las temperaturas sean constantes sobre los 32,2° C.
5. No moje las hojas de las hierbas por 1 ó 2 días después de la aplicación. Tras ese período, se debe realizar un riego riguroso.

CÓMO APLICAR: ESPARCIDORES Y CALIBRACIÓN

Aplique siempre con un esparcidor giratorio o por voleo para distribuir los gránulos uniformemente. Nunca aplique manualmente.
Cobertura – Para determinar cuánto necesita aplicar, multiplique el largo por el ancho del césped. Esto es igual al área total en metros cuadrados. Reste todas las áreas no pertenecientes al césped (casa, jardines, etc.). Una bolsa cubre hasta 464,52 m^2.
Importante: Es importante seguir las instrucciones como se indica en este producto para el cuidado del césped de calidad, ya que aplicar más de lo debido podría dañarlo. Las graduaciones del esparcidor son aproximadas. Se pueden producir diferencias debido al estado del esparcidor, la velocidad con la que se aplica y la escabrosidad del área tratada. Asegúrese de revisar el esparcidor antes de comenzar y haga ajustes de ser necesario. Consulte el Manual del propietario. Nunca aplique manualmente.
INSTRUCCIONES DE CALIBRACIÓN Y GRADUACIONES DEL ESPARCIDOR:
Calibre el esparcidor de acuerdo con las instrucciones del fabricante. Las graduaciones iniciales del esparcidor podrían requerir un ajuste para entregar las dosis recomendadas bajo condiciones de aplicación reales.
Para una distribución adecuada, calibre el esparcidor antes de la aplicación.
• Para 1,18 kg del producto por cada 92,90 m2 de césped: Calibre su esparcidor en 35,77 m2 y ajuste para aplicar una dosis de 453,59 g de producto por cada 35,77 m2.

ALMACENAMIENTO Y ELIMINACIÓN

No contamine el agua, los alimentos o el forraje al almacenarlo o eliminarlo.
ALMACENAMIENTO DEL PESTICIDA: Almacénelo en su envase original y mantenga las bolsas que contengan producto sin utilizar en un área de almacenaje cerrada, que no esté al alcance de los niños y los animales domésticos.
ELIMINACIÓN DEL PESTICIDA Y MANIPULACIÓN DEL ENVASE: El envase no se puede volver a rellenar. No vuelva a usar ni rellene este envase. **Si está vacío:** Colóquelo en la basura o, si está disponible, ofrézcalo para reciclaje. **Si está parcialmente lleno:** Llame a su agencia local de desechos sólidos para obtener instrucciones sobre cómo eliminar el producto. Nunca deseche el producto sin usar en desagües en interiores o exteriores.

Sta-Green® *ANNUAL FEEDING PROGRAM*

A good fertilization program can keep your lawn green and vigorous. Sta Green has a wide selection of fertilizers to help you throughout the year.
Un buen programa de fertilización puede mantener el césped verde y vigoroso. Sta Green cuenta con una amplia selección de fertilizantes para brindarle ayuda durante todo el año.

STA-GREEN CRABGRASS PREVENTERS

Use these pre-emergent products to control crabgrass and many other weeds in your lawn before they germinate. Crabgrass preventers are available with or without fertilizer. Use these products in early spring before weeds begin to emerge.
Use estos productos preemergentes a fin de controlar la digitaria y muchas otras malezas en su césped antes de que germinen. Existen productos para prevenir la digitaria con y sin fertilizante. Utilice estos productos a comienzos de la primavera antes de que comiencen a brotar las malezas.

STA-GREEN LAWN FERTILIZER

A high nitrogen fertilizer with iron that feeds any established lawn up to 4 months. Use this product anytime during the active growing season.
Un fertilizante con alto contenido de nitrógeno y hierro que abona cualquier césped establecido hasta por 4 meses.

STA-GREEN WEED & FEED

This post-emergent product controls a wide variety of existing broadleaf weeds while it feeds your desirable grass. Use weed and feeds when weeds are actively growing.
Este producto postemergente controla una amplia variedad de malezas de hojas anchas a la vez que abona el césped que usted desea.

STA-GREEN SOUTHERN WEED & FEED

This fertilizer is especially formulated to control a wide variety of broadleaf weeds in Centipede and St. Augustine lawns. Use weed and feeds when weeds are actively growing.
Este fertilizante está especialmente formulado para controlar una amplia variedad de malezas de hojas anchas en céspedes de San Agustín y cienpiés.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.
PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.
PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If Empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Keep this product off sidewalks, patios, driveways, etc. This fertilizer may stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.

When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials. Information regarding the contents and levels of metals in this product is available on the Internet at **http://www.aapfco.org/metals.htm.**

LIMITED WARRANTY AND DISCLAIMER
FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

†MONEY BACK GUARANTEE
If you are not completely satisfied with this product, please mail your name, address and proof of purchase (package UPC) and a brief explaination for seeking a refund.

5M = 0-21496-00266-8

Covers up to 5,000 sq ft

LOOK! For more information, visit us on our website at
www.Sta-Green.com
¡BUSQUE! Para obtener más información, visite nuestro sitio web

Manufactured for
Gro Tec, Inc.
P. O. Box 290
Madison, GA 30650

Net Wt 13 lb (5.90 kg)

12.75" WIDTH
12.875" STEP

Kills over 250 Weeds
as listed

Sta-Green®

PHOSPHORUS - FREE
WEED & FEED

Controls a wide spectrum of Broadleaf weeds

Controla una amplia variedad de malezas de hoja ancha

Use on lawns to kill the toughest weeds like clover and dandelion

Feeds for 3 months guaranteed!*

Utilizado en céspedes para eliminar malezas resistentes como tréboles y dientes de león ¡Nutrición garantizada durante 3 meses!*

- Kills the root
  Mata la raíz
- Starts working on contact
  Comienza a funcionar al contacto
- Quick greening and extended feeding*
  Proporciona un color verde rápido y una alimentación continua*

ACTIVE INGREDIENTS:

2,4-D, 2-ethylhexyl ester	1.108%
Mecoprop-p acid	0.167%
Dicamba acid	0.071%
OTHER INGREDIENTS:	98.654%
TOTAL	100.000%

THIS PRODUCT CONTAINS:
14.70 lb. 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735%. CAS 1928-43-4.
3.34 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.167%. CAS 16484-77-8.
1.42 lb. 3,6-dichloro-o-anisic acid equivalent per ton or 0.071%. CAS 1918-00-9.
Isomer Specific by AOAC Methods

KEEP OUT OF REACH OF CHILDREN
CAUTION
See Back Panel For First Aid And Additional Precautionary Statements

COVERS 15,000 SQ FT
MONEY BACK GUARANTEED† • NET WT 39 LB (17.69 KG)

28-0-4
SEE BACK PANEL FOR ANALYSIS DESCRIPTION

Sta-Green
PHOSPHORUS-FREE WEED & FEED
Controls a wide spectrum of Broadleaf weeds
Use on lawns to kill the toughest weeds like clover and dandelion
Feeds for 3 months guaranteed!*
COVERS 15,000 SQ FT

Sta-Green
PHOSPHORUS - FREE
WEED & FEED
Controls a wide spectrum of Broadleaf weeds
Kills over 250 Weeds as listed
COVERS 15,000 SQ FT • NET WT 39 LB (17.69 KG)

Net Wt 39 lb

WITH Pennington Coated® TECHNOLOGY — GREENS QUICKLY, FEEDS FOR MONTHS

Pennington Coated – Slow Release Technology
*For a Long Lasting Green Lawn With No Burning**

PHOSPHORUS FREE – Helping Our Water Sources

28 N (nitrogen) - **0 P** (phosphorus) - **4 K** (potassium)
N: For lush, green plant growth
K: For overall plant health and vigor

GUARANTEED ANALYSIS F525 **28-0-4**

Total Nitrogen (N)	28.00%
27.13% Urea Nitrogen*	
0.87% Ammoniacal Nitrogen	
Soluble Potash (K_2O)	4.00%
Sulfur (S)	0.99%
0.99% Combined Sulfur (S)	

Derived from: Urea, Polymer Coated Urea, Ammonium Sulfate, Muriate of Potash.
*Part of the nitrogen in this product has been polymer coated to provide 5.70% coated slow release nitrogen.

STOP READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If in Eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:
- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

INSTRUCTIONS: Sta-Green Weed and Feed is a combination of three proven weed killers -- 2,4-D, mecoprop-p, and dicamba. This herbicide controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 28-0-4 fertilizer provides a supply of plant nutrients.

USE PRECAUTIONS

Do not use on dichondra, carpetgrass, St. Augustinegrass, creeping bentgrass, or turfgrass where desirable clovers are present.
Do not apply this product to newly seeded grasses until well-established or after the second or third mowing.
Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area and do not apply this product within the dripline of, or over surface roots of trees, shrubs, and other ornamentals.
Reseeding or overseeding areas may occur three (3) weeks after the application of this product.
Do not use this product on or near desirable plants and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of the roots of desirable trees and shrubs, since injury may result.
Do not apply directly to or near water, storm drains, gutters, sewers or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

HOW MUCH TO APPLY

For use on: Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, zoysiagrass, St. Augustinegrass, centipedegrass, hybrid bermudagrass, and colonial bentgrass. Apply 2.6 pounds of product per 1,000 square feet of turfgrass. A second application may be made in approximately 30 days. The maximum number of broadcast applications per treatment site is 2 per year.

This product controls the following list of growing weeds:

Amaranth, purple	Chicory	Galinsoga, hairy	Mustard, hedge	Ragweed, bur	Thistle, wavyleaf
Amaranth, slender	Cinquefoil	Galinsoga, smallflower	Mustard, indian,	Ragweed, common	Thistle, yellow (horrible)
Aster, heath	Clover, crimson	Goldenrod	Mustard, tansy	Ragweed, lanceleaf	Trailing crownvetch
Aster, white heath	Clover, alyce	Ground ivy	Mustard, tumble	Ragweed, western	Velvetleaf
Aster, purple	Clover, hop	Groundsel	Mustard, white	Redstem filaree	Vervain, blue
Aster, white prairie	Clover, rabbit foot	Gumweed	Mustard, wormseed	Rough cinquefoil	Vervain, hoary
Artichoke	Clover, red	Hairy fleabane	Nettle, stinging (burning)	Scarlet pimpernel	Vervain, prostrate
Austrian fieldcress	Clover, strawberry	Hawkweed, meadow	Nettle, tall	Shepardspurse	Vervain, tall
Bedstraw, Catchweed	Clover, yellow sweet	Hawkweed, mouseear	Nettle, wood	Smartweed, ladysthumb	Vetch, bird
Bedstraw, smooth	Clover, white sweet	Hawkweed, orange	Parsley-piert	Smartweed, pale	Vetch, hairy
Bedstraw, Northern	Clover, white	Hawkweed, yellow	Parsnip	Smartweed, pennsylvania	Vetch, milk
Bedstraw, pineywoods	Cockle	Healall	Pearlwort	Smooth chaff-flower	Vetch, narrowleaf
Beggarticks, hairy	Cocklebur	Heartleaf drymary	Pennycress (frenchweed)	Smoothdock	Vetch, wild
Betony, Florida	Cocklebur, oriental	Hemp	Pepperweed, perennial	Sorrel, red (sheep)	Virginia creeper
Bindweed, Field	Cocklebur, spiny	Henbit	Pepperweed, virginia (Peppergrass)	Sowthistle, annual	Western clematis
Bindweed, Hedge	Common mullein	Hoary cress	Pigweed, prostate	Sowthistle, spiny	Western salsify
Bitter Wintercress	Corn Spurry	Horsenettle	Pigweed, redroot	Spanish needles	Wild aster
Bittercress, hairy	Creeping Beggarweed	Horseweed	Pigweed, russian	Speedwell, bi-lobed	Wild buckweat
Bitterweed	Creeping Jenny (creeping charlie)	Jimsonweed	Pigweed, smooth	Speedwell, common	Wild carrot
Black-eyed Susan	Cupid shaving Brush	Johnny-Jumpup violet	Pigweed, tumble	Speedwell, corn	Wild four-o-clock
Black Medic	Cudweed	Knawel	Pineapple weed	Speedwell, ivyleaf	Wild garlic
Blood flower milkweed	Daisy, English	Knotweed, prostrate	Plains coreopsis (tickseed)	Speedwell, persian	Wild lettuce
Blue lettuce	Daisy, oxeye	Kochia	Plantain, blackseed	Speedwell, purslane	Wild marigold
Brassbuttons	Dandelion, common	Lambsquarters	Plantain, bracted	Speedwell, slender	Wild mustard (yellow rocket)
Bristly oxtongue	Dandelion, false	Lespedeza	Plantain, broadleaf	Speedwell, snow	Wild onion
Broomweed	Day flower	Mallow, alkali	Plantain, buckhorn	Speedwell, thymeleaf	Wild parsnip
Buckhorn	Deadnettle	Mallow, bristly	Plantain, hoary	Spiny amaranth	Wild radish
Bullnettle	Dichondra	Mallow, common	Plantain, narrowleaf	Spurge, spotted	Wild rape
Burclover	Dock, broadleaf	Mallow, dwarf	Plantain, slender	Spurge, prostrate	Wild strawberry
Burdock	Dock, curly	Mallow, venice	Plantain, woolly	Spurweed	Wild sweet potato
Burweed, lawn (spurweed)	Dogbane	Marcela	Poison ivy	Star of Bethlehem	Wild violet
Buttercup, bulbous	Dogfennel	Matchweed	Poison oak	Strawberry, India mock	Woodsorrel, creeping
Buttercup, creeping	Dollarweed (Pennywort)	Mexicanweed	Pokeweed	Tansy ragwort	Woodsorrel, violet
Buttonweed, common (poorjoe)	Dove weed	Moneywort	Prairie Sunflower	Tanweed	Woodsorrel, yellow (Oxalis)
Buttonweed, virginia	Elderberry	Morningglory, bigroot	Prickly lettuce (compass plant)	Thistle, blessed	Woolly croton
Carolina geranium (wild)	Falseflax	Morningglory, ivyleaf	Prickly sida	Thistle, bull	Wormseed
Carpetweed	False sunflower	Morningglory, mexican	Prostrate knotweed	Thistle, canada	Yarrow, common
Catnip	Fiddleneck	Morningglory, red	Puncturevine	Thistle, flodman (prairie)	Yarrow, western
Catsear, spotted (false dandelion)	Filaree, redstem	Morningglory, tall	Purslane, common	Thistle, musk	Yarrow, yellow
Chamber bitter	Filaree, whitestem	Morningglory, wolly	Pusley, brazil	Thistle, platte	
Chickweed, common	Fleabane, daisy (annual)	Mugwort Mustard, ball	Pusley, florida	Thistle, plumeless	
Chickweed, mouseear	Fleabane, rough	Mustard, black	Pusley, large flower	Thistle, russian	
Chickweed, sticky		Mustard, blue		Thistle, scotch	

WHEN TO APPLY

Apply when broadleaf weeds are actively growing in spring or early fall.
Apply when the air is calm to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery.
Possible retreatment may be necessary if lawn is heavily overrun by weeds, if adverse low moisture conditions prevail, or if weeds are in a state of poor growth.

APPLICATION INSTRUCTIONS

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
3. Apply when broadleaf weeds are young and actively growing, preferably in the morning when dew is on the grass. At the time of application, moisture on the weed leaves from dew, rainfall or irrigation/watering aids in control. If grass is not moist at time of application, sprinkle lightly with water before applying product.
4. Avoid applying when temperatures are consistently above 90°F.
5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

HOW TO APPLY – SPREADERS AND CALIBRATION

Always apply with a rotary or drop spreader to distribute the granules evenly. Never apply by hand.

Coverage – To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to 5000 square feet.

Important: It is important to follow directions as indicated on this quality lawn care product as over-application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owner's Manual. Never apply by hand.

SINGLE PASS DISTRIBUTION:
- For rotary spreaders: Operate at intervals corresponding to the swath width.
- For drop spreaders: Overlap the wheel tracks to ensure complete coverage.
- For best results, apply this product uniformly and avoid streaks, skips or overlaps. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping.

SPREADER SETTINGS
SPREADER SETTINGS(SPANISH)

BROADCAST SPREADER (BROADCAST SPREADER)		DROP SPREADER (DROP SPREADER)	
Ortho Rotary	1¾	Republic EZ	5
True Temper	3¾	Ortho Drop	5
EV-N-SPRED	14	Precision Drop	6
Precision Rotary	5	Scotts AccuGreen	6¾
Republic EZ Grow	5¾	Scotts AccuGreen 1000	6¾
Scotts EdgeGuard	5¾	Scotts AccuGreen 3000	6¾
Scotts Easy Green	24¾	Sta-Green DS4500	2
Scotts SpeedyGreen 1000	5¾		
Scotts SpeedyGreen 2000	5¾		
Scotts SpeedyGreen 3000	5¾		
Sta-Green SB5500	5½		
Sta-Green SB4500	5½		

CALIBRATION IINSTRUCTIONS AND SPREADER SETTINGS:
Calibrate the spreader according the the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended rate under actual application conditions.

To provide proper distribution, calibrate spreader before application.
- For 2.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 385 square feet and adjust to apply at the rate of 1 pound of product per 385 square feet.

BROADCAST SPREADERS: For best results, apply in a circular path starting on the outside and working toward the center overlapping slightly in insure complete coverage.

DROP SPREADERS: Apply 2 strips along each end. Then apply along length of lawn, overlapping slightly and shutting off the spreader before each turn at edge of border strips.

EPA Reg. No. 2217-884-59144
EPA Est. No. 59144-GA-1(A), 32802-MO-1(B), 37351-CO-1(C), 8378-IN-1(D), 9404-FL-1(E), 961-PA-01(F), 961-IL-01(G), 35497-OR-1(H), 49991-TX-2(Q)
Letter in parenthesis is first letter of lot number.

WITH Pennington Coated® TECHNOLOGY — GREENS QUICKLY, FEEDS FOR MONTHS

Pennington Coated – Slow Release Technology
*For a Long Lasting Green Lawn With No Burning**

PHOSPHORUS FREE – Helping Our Water Sources

ALTO LEA LA ETIQUETA COMPLETA PRIMERO. CUMPLA CON TODAS LAS PRECAUCIONES Y SIGA LAS INSTRUCCIONES DETALLADAMENTE.

DECLARACIONES DE PRECAUCIÓN

RIESGOS PARA SERES HUMANOS Y ANIMALES DOMÉSTICOS

PRECAUCIÓN. Provoca una irritación moderada de los ojos. Evite el contacto con los ojos o la ropa. Lávese bien con agua y jabón después de manipular este producto y antes de comer, beber, masticar chicle, usar tabaco o usar el inodoro.

PRIMEROS AUXILIOS

Si entra en contacto con los ojos:	• Mantenga los ojos abiertos y enjuague lenta y suavemente con agua durante 15 a 20 minutos. • Si lleva lentes de contacto puestos, quíteselos luego de los primeros 5 minutos, luego continúe enjuagando los ojos. • Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.

Cuando llame a un centro de control de envenenamientos o a un médico, o busque tratamiento, tenga a mano el envase o la etiqueta del producto. También puede ponerse en contacto con el 1-800-424-9300 para obtener información de tratamiento médico de emergencia.

Riesgos ambientales

Este producto es tóxico para los peces e invertebrados acuáticos y puede tener efectos adversos en plantas que no eran el objetivo del tratamiento. Para proteger el medio ambiente, no deje que el pesticida entre o escurra hacia desagües pluviales, zanjas de desagüe, alcantarillas ni aguas superficiales. Aplicar este producto en un clima tranquilo, cuando no haya pronóstico de lluvia durante las siguientes 24 horas, ayudará a asegurar que el viento o la lluvia no soplen o laven el pesticida del área de tratamiento. Barra el producto que caiga sobre una entrada de garaje, acera o calle llevándolo de vuelta al área tratada del césped o jardín para evitar que se escurra hacia aguas superficiales o sistemas de desagüe.

Este producto químico tiene propiedades y características asociadas con agentes químicos detectados en agua del subsuelo. El uso de este producto químico en áreas donde el suelo es permeable, especialmente donde las napas de agua son de poca profundidad, puede ocasionar contaminación del agua del subsuelo. La aplicación alrededor de una cisterna o pozo puede ocasionar contaminación del agua potable o agua del subsuelo.

INSTRUCCIONES DE USO:

Se considera una violación a la Ley Federal usar este producto con fines que no sean los indicados en la etiqueta.

PRECAUCIONES Y RESTRICCIONES:
- No aplique el producto de ninguna forma que pueda entrar en contacto con personas o mascotas, ya sea en forma directa o indirecta. Mantenga a las personas y mascotas fuera del área durante la aplicación. No permita que las personas o mascotas ingresen al área tratada hasta que el polvo se haya asentado.
- No lo use en césped que se cultiva para la venta u otro uso comercial como tepe, para la producción comercial de semillas ni para fines de investigación.

INSTRUCCIONES: El herbicida y abono Sta-Green es una combinación de tres herbicidas comprobados: 2,4-D, mecoprop-p y dicamba. Este herbicida controla una gran variedad de malezas de césped, incluyendo dientes de león, pamplina, sanguinaria, llantén, ortiga muerta, hierba de golondrina y muchas otras que a menudo se encuentran en los céspedes domésticos. Además, un fertilizante 28-0-4 proporciona un suministro de nutrientes para las plantas.

PRECAUCIONES DE USO

No lo use sobre céspedes oreja de ratón, alfombra, San Agustín, agrostis o sobre céspedes donde haya tréboles que se desean conservar.
No aplique este producto sobre céspedes recién sembrados hasta que estén bien establecidos o después de una segunda o tercera poda.
No use este producto sobre flores, hortalizas, cubiertas vegetales, jardines o plantaciones, incluyendo arbustos y árboles. No exceda la dosis indicada para cada área y no aplique este producto dentro de la línea de goteo o sobre la superficie de las raíces de árboles, arbustos y otras plantas ornamentales.
Se puede volver a sembrar o sembrar varias veces después de tres (3) semanas desde la aplicación de este producto.
No use este producto en o cerca de plantas deseadas, y evite esparcir gránulos en sistemas de raíces expuestos o brotes espontáneos (rizomas) dentro de la línea de goteo de las raíces de los árboles y arbustos deseados, ya que podría provocar daños.
No lo aplique directamente en agua, desagües de aguas pluviales, canaletas, cloacas o zanjas de desagüe ni cerca de éstos. No lo aplique a menos de 7,62 metros de ríos, estanques de peces, lagos, arroyos, embalses, pantanos, estuarios, bahías y océanos. No lo aplique bajo condiciones de viento. Aplique este producto directamente en el césped y barra el producto que caiga sobre la entrada de garaje, la acera o la calle llevándolo de vuelta al césped. Para evitar que el producto escurra, no riegue en exceso las áreas tratadas ni aplique cuando esté lloviendo o cuando se esperen lluvias dentro del día.

CUÁNTO APLICAR

Para usar en: Césped azul de Kentucky, festuca spp., ballico perenne, césped Bermuda común, césped zoysia, césped de San Agustín, césped ciempiés, césped Bermuda híbrido y césped agrostis colonial. Aplique 1,18 kg del producto por cada 92,90 m^2 de césped. Se puede realizar una segunda aplicación en aproximadamente 30 días. El número máximo de aplicaciones por dispersión por sitio de tratamiento es de 2 por año.

CUÁNDO APLICAR

Aplique cuando las hierbas de hojas anchas estén en pleno crecimiento en primavera o a principios del otoño.
Aplique cuando no haya brisa para asegurar una cobertura uniforme y evitar que se esparzan los gránulos sobre flores, hortalizas y arbustos ornamentales.
Podría ser necesario un nuevo tratamiento si el césped está excesivamente invadido por malezas; si prevalecen condiciones adversas de humedad baja o si las malezas están en un período de crecimiento pobre.

INSTRUCCIONES PARA LA APLICACIÓN

1. Corte el césped a altura normal 1 a 2 días antes de la aplicación.
2. Riegue bien el césped al menos 1 a 2 días antes de la aplicación para mantener la humedad hasta el siguiente riego (consulte el paso 5).
3. Aplique cuando las malezas de hojas anchas estén jóvenes y en pleno crecimiento, de preferencia por la mañana cuando hay rocío sobre el césped. En el momento de la aplicación, la humedad en las hojas de las hierbas proveniente del rocío, lluvia o irrigación/regado ayuda en el control. Si el césped no está húmedo en el momento de la aplicación, rocíelo ligeramente con agua antes de aplicar el producto.
4. Evite aplicar el producto cuando las temperaturas sean constantes sobre los 32,2° C.
5. No moje las hojas de las hierbas por 1 ó 2 días después de la aplicación. Tras ese período, se debe realizar un riego riguroso.

CÓMO APLICAR: ESPARCIDORES Y CALIBRACIÓN

Aplique siempre con un esparcidor giratorio o por voleo para distribuir los gránulos uniformemente. Nunca aplique manualmente.

Cobertura – Para determinar cuánto necesita aplicar, multiplique el largo por el ancho del césped. Esto es igual al área total en metros cuadrados. Reste todas las áreas no pertenecientes al césped (casa, jardines, etc.). Una bolsa cubre hasta 464,52 m^2.

Importante: Es importante seguir las instrucciones como se indica en este producto para el cuidado del césped de calidad, ya que aplicar más de lo debido podría dañarlo. Las graduaciones del esparcidor son aproximadas. Se pueden producir diferencias debido al estado del esparcidor, la velocidad con la que se aplica y la escabrosidad del área tratada. Asegúrese de revisar el esparcidor antes de comenzar y haga ajustes de ser necesario. Consulte el Manual del propietario. Nunca aplique manualmente.

INSTRUCCIONES DE CALIBRACIÓN Y GRADUACIONES DEL ESPARCIDOR:
Calibre el esparcidor de acuerdo con las instrucciones del fabricante. Las graduaciones iniciales del esparcidor podrían requerir un ajuste para entregar las dosis recomendadas bajo condiciones de aplicación reales.

Para una distribución adecuada, calibre el esparcidor antes de la aplicación.
- Para 1,18 kg del producto por cada 92,90 m2 de césped: Calibre su esparcidor en 35,77 m2 y ajuste para aplicar una dosis de 453,59 g de producto por cada 35,77 m2.

ALMACENAMIENTO Y ELIMINACIÓN

No contamine el agua, los alimentos o el forraje al almacenarlo o eliminarlo.
ALMACENAMIENTO DEL PESTICIDA: Almacénelo en su envase original y mantenga las bolsas que contengan producto sin utilizar en un área de almacenaje cerrada, que no esté al alcance de los niños y los animales domésticos.
ELIMINACIÓN DEL PESTICIDA Y MANIPULACIÓN DEL ENVASE: El envase no se puede volver a rellenar. No vuelva a usar ni rellene este envase. **Si está vacío:** Colóquelo en la basura o, si está disponible, ofrézcalo para reciclaje. **Si está parcialmente lleno:** Llame a su agencia local de desechos sólidos para obtener instrucciones sobre cómo eliminar el producto. Nunca deseche el producto sin usar en desagües en interiores o exteriores.

Sta-Green® ANNUAL FEEDING PROGRAM

A good fertilization program can keep your lawn green and vigorous. Sta Green has a wide selection of fertilizers to help you throughout the year.
Un buen programa de fertilización puede mantener el césped verde y vigoroso. Sta Green cuenta con una amplia selección de fertilizantes para brindarle ayuda durante todo el año.

STA-GREEN CRABGRASS PREVENTERS

Use these pre-emergent products to control crabgrass and many other weeds in your lawn before they germinate. Crabgrass preventers are available with or without fertilizer. Use these products in early spring before weeds begin to emerge.
Use estos productos preemergentes a fin de controlar la digitaria y muchas otras malezas en su césped antes de que germinen. Existen productos para prevenir la digitaria con y sin fertilizante. Utilice estos productos a comienzos de la primavera antes de que comiencen a brotar las malezas.

STA-GREEN LAWN FERTILIZER

A high nitrogen fertilizer with iron that feeds any established lawn up to 4 months. Use this product anytime during the active growing season.
Un fertilizante con alto contenido de nitrógeno y hierro que abona cualquier césped establecido hasta por 4 meses.

STA-GREEN WEED & FEED

This post-emergent product controls a wide variety of existing broadleaf weeds while it feeds your desirable grass. Use weed and feeds when weeds are actively growing.
Este producto postemergente controla una amplia variedad de malezas de hojas anchas a la vez que abona el césped que usted desea.

STA-GREEN SOUTHERN WEED & FEED

This fertilizer is especially formulated to control a wide variety of broadleaf weeds in Centipede and St. Augustine lawns. Use weed and feeds when weeds are actively growing.
Este fertilizante está especialmente formulado para controlar una amplia variedad de malezas de hojas anchas en céspedes de San Agustín y cienpiés.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.
PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.
PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If Empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Keep this product off sidewalks, patios, driveways, etc. This fertilizer may stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.

When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials. Information regarding the contents and levels of metals in this product is available on the Internet at **http://www.aapfco.org/metals.htm.**

LIMITED WARRANTY AND DISCLAIMER
FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

†MONEY BACK GUARANTEE
If you are not completely satisfied with this product, please mail your name, address and proof of purchase (package UPC) and a brief explaination for seeking a refund.

15M = 0-21496-00267-5

Covers up to 15,000 sq ft

LOOK! For more information, visit us on our website at
www.Sta-Green.com
¡BUSQUE! Para obtener más información, visite nuestro sitio web

Manufactured for
Gro Tec, Inc.
P. O. Box 290
Madison, GA 30650

Net Wt 39 lb (17.69 kg)