STATE OF HAWAII
Department of Agriculture
ACCEPTED
LICENSE NO.
8079.109

Expert GARDENER®

CRABGRASS PREVENTER
PLUS FERTILIZER

CONTROLS CRABGRASS AND BROADLEAF WEEDS BEFORE THEY GERMINATE

Provides lush, green grass
Ofrece un césped verde y tupido

- Prevent crabgrass and fertilize your lawn in one application
  Impida el crecimiento de digitarias y fertilice su césped en una sola aplicación
- Up to 5 month weed and grass control
  Hasta 5 meses de control de malezas y césped

ACTIVE INGREDIENTS:	
Prodiamine: (CAS 29091-21-2)	0.29%
INERT INGREDIENTS:	99.71%
Total	100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION
See back panel for further precautions and First Aid information.

COVERS 5,000 SQ FT
MONEY BACK GUARANTEED† • NET WT 15.43 LB (7.01 KG)

30-0-4
SEE BACK PANEL FOR ANALYSIS DESCRIPTION

CRABGRASS PREVENTER PLUS FERTILIZER 30-0-4

CRABGRASS PREVENTER PLUS FERTILIZER 30-0-4

Expert GARDENER®

CRABGRASS PREVENTER
PLUS FERTILIZER

COVERS 5,000 SQ FT
NET WT 15.43 LB (7.01 KG)

30-0-4

PHOSPHORUS FREE – Helping Our Water Sources

SIN FÓSFORO: Cuida Nuestras Napas de Agua

READ THE ENTIRE "DIRECTIONS FOR USE" AND "WARRANTY STATEMENT" BEFORE USING THIS PRODUCT. IF TERMS ARE NOT ACCEPTABLE, RETURN THE UNOPENED PRODUCT CONTAINER AT ONCE TO THE SELLER.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION:

PRECAUCIONES
RIESGOS PARA SERES HUMANOS Y ANIMALES DOMÉSTICOS
ATENCIÓN:

FIRST AID

ENVIRONMENTAL HAZARDS

DIRECTIONS FOR USE

INSTRUCCIONES DE USO

STORAGE AND DISPOSAL

SPREADER INSTRUCTIONS

OTHER RECOMMENDED Expert LAWN PRODUCTS

Net Wt 15.43 lb (7.01 kg)

WEB 39.0"

18"

1.5"

3"

CUTOFF 13.5"

SGS Southern Graphic Systems	CUSTOMER	PRINTER	PRINTER #
SGS Charlotte, 9201-A Forsythe Park Drive, Charlotte, NC 28273, Phone: 704-587-3899, Fax: 704-587-3898	GRO TEC/Pennington	Plastic Pkg	
		LOCATION: Hickory, NC	END USER #
	DESCRIPTION: Grotec Inc_ Expert Gardener 007007-Crabgrass Preventer 5m 3/cs - 15.43 lb_P57638	JOB # 3136265_Pt8	OTHER: XXX
		DATE: 11.23.10	CSR: Chris Walker
		WEB: 41	CODE: UPC - A 0-21496-00700-7
	DRAWING # XX-XX-XX / SPEC# XX-XX-XX / DIE# XX-XX-XX	DISTORT: 99.07% Around	CUTOFF: 13.5
		MAG: 100%/80%	BWA: -0.003
		CYL: 27	ACR: 1 / ARO: 2
		L/S: 120	

MAC- SJ; Suz, SJ; Suz; CW-->, Suz
QC-
Final Color QC-

PRINT COLORS:
BLACK LINE, BLACK SCREEN, BLUE 286 BKG, GREEN 349, BLUE 286 BUMP, PRO CYAN, PRO MAGENTA, YELLOW LINE, COMBO YELLOW, WHITE

Placed Images:
shutterstock 16588846 ICC cc5.eps

ATTENTION:
Due to the bright, clean colorants used to create some of the PANTONE MATCHING SYSTEM Colors, not all PANTONE colors will be accurately simulated on this proof. The above swatches should be used as a reference only. Please refer to a PANTONE® Color Formula Guide for precise color communication.

PHOSPHORUS FREE – Helping Our Water Sources

30	0	4
N (nitrogen)	P (phosphate)	K (potassium)
For lush, green plant growth		For overall plant health and vigor

F525

GUARANTEED ANALYSIS 30-0-4

Total Nitrogen (N) 30.00%
30.00% Urea Nitrogen
Soluble Potash (K_2O) 4.00%

Derived from: Urea, and Muriate of Potash.

STOP READ THE ENTIRE "DIRECTIONS FOR USE" AND "WARRANTY STATEMENTS" BEFORE USING THIS PRODUCT. IF TERMS ARE NOT ACCEPTABLE, RETURN THE UNOPENED PRODUCT CONTAINER AT ONCE TO THE SELLER.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow this product to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather will help to ensure that wind does not blow this product off the treated area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

FIRST AID

If in eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	• Call a poison control center or physician immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or physician. • Do not give anything by mouth to an unconscious person.
If inhaled	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

Hot line number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens). It controls susceptible weeds by inhibiting weed seeds germination and root development. When used according to directions, it will control: barnyardgrass; bluegrass, annual (*Poa Annua*); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass[1]; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass[3]; shepherd's purse[2]; signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

1 In many areas a single application of 5.2 to 11.9 pounds/1000 square feet will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 5.2 to 7.9 pounds/1000 square feet, followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).
2 For this weed, apply in late summer, fall or winter prior to germination.
3 Suppression only.

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing .92 pounds of nitrogen per 1,000 square feet and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

This product may stain driveways, walks, patios, pools, etc. Sweep off thoroughly immediately after each use.

WHEN TO APPLY TO ESTABLISHED TURF

This product must be applied prior to germination of weed seeds, as it will not control established weeds.



APPLICATION DIRECTIONS

Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days after application.

USE PRECAUTIONS

The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil). Do not apply to newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa Annua*) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

RATES OF APPLICATION

This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)

Turf Species	Minimum Rate	Maximum Annual Rate[1]
Bahiagrass, Bermudagrass[2] Centipedegrass, Zoysiagrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf-type)	2.75 lbs/1000 sq ft	11.9 lbs/1000 sq ft per year
Buffalograss Kentucky Bluegrass Perennial Ryegrass	2.75 lbs/1000 sq ft	7.9 lbs/1000 sq ft per year
Creeping Red Fescue	2.75 lbs/1000 sq ft	6 lbs/1000 sq ft per year
Creeping Bentgrass	2.75 lbs/1000 sq ft	5.2 lbs/1000 sq ft per year

1 These are the maximum rates per calendar year.
2 May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 4.1 pounds/1000 square feet. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/ reseeding results.

USE RATE	MONTHS BEFORE OVERSEEDING North	Transition	South
2.75 lbs/1000 sq ft	4	4	4
5.2 lbs/1000 sq ft	5	4	4
6 lbs/1000 sq ft	6	5	5
6.3 lbs/1000 sq ft	---	6	6
7.9 lbs/1000 sq ft	---	7	7
9 lbs/1000 sq ft	---	---	9
10.3 lbs/1000 sq ft	---	---	10
11.9 lbs/1000 sq ft	---	---	12

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

SPREADER USE INSTRUCTIONS

1. Determine size of area to be treated – measure length and width in feet, then multiply to determine coverage area in square feet (length x width). Next, subtract non-lawn areas.
2. Use the spreader setting chart to determine the appropriate setting.
3. Apply evenly over dry grass, avoiding streaks, skips and overlaps – thorough coverage is important. On hot days (above 80°F), water the lawn to rinse particles off the grass.

Note: The spreader settings listed at right are approximate. It is important to check calibration of spreader frequently to ensure spreader is functioning properly and applying product evenly. To check calibration, accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust spreader setting if necessary in order to ensure correct rate of application.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back onto your lawn.

SPREADER SETTINGS

BROADCAST SPREADER	
Scotts SpeedyGreen	5
Scotts Basic	5
Scotts S Lawn Pro	5
Scotts Edge Guard Deluxe	5
Scotts Edge Guard Mini	5
Precision Rotary	3.75
EV-N-SPRED	11.5
DROP SPREADER	
Scotts AccuGreen	6.5
Earthway	9.5

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store in original container in a locked storage area out of reach of children and domestic animals. **IF EMPTY:** Nonrefillable container: Do not reuse or refill this container. Offer for recycling, if available. **IF PARTLY FILLED:** Call your local solid waste agency for disposal instructions. Never place any unused product down any indoor or outdoor drain or toilet.

Information regarding the contents and levels of metals in this product is available on the Internet at http://www.aapfco.org/metals.htm

†MONEY BACK GUARANTEE
If you are not completely satisfied with this product, please mail your name, address and proof of purchase (package UPC) and a brief explanation for seeking a refund.

EPA Reg. No. 32802-70-59144
EPA Est. No. 59144-GA-1(A), 32802-MO-1(B), 37351-CO-1(C), 8378-IN-1(D), 9404-FL-1(E), 961-PA-01(F), 961-IL-01(G), 35497-OR-1(H), 49991-TX-2(Q)
Letter in parenthesis is first letter of lot number.

Net Wt 46.29 lb (21.00 KG)

SIN FÓSFORO: Cuida Nuestras Napas de Agua

ALTO LEA TODAS LAS "INSTRUCCIONES DE USO" Y LA "GARANTÍA" ANTES DE USAR EL PRODUCTO. SI LOS TÉRMINOS NO SON ACEPTABLES, DEVUELVA EL ENVASE DEL PRODUCTO SIN ABRIR DE INMEDIATO AL VENDEDOR.

PRECAUCIONES

RIESGOS PARA SERES HUMANOS Y ANIMALES DOMÉSTICOS

ATENCIÓN: Produce lesiones oculares moderadas (irritación). Su inhalación o absorción por la piel son perjudiciales. Evite el contacto con la piel, los ojos o la vestimenta. Evite aspirar el polvo. Lave bien con agua y jabón después de manipularlo. El contacto prolongado o reiterado del producto con la piel puede provocar reacciones alérgicas a algunas personas.

RIESGOS AMBIENTALES

Para proteger el medio ambiente, no permita el ingreso de este producto o que se escurra al interior de bocas de tormenta, zanjas de drenaje, alcantarillas o espejos de agua. La aplicación del producto en días de buen clima contribuirá a asegurar que el viento no lo arrastre fuera del área tratada. Barrer cualquier cantidad del producto que se deposite en entradas de automóvil, veredas o calles nuevamente hacia el área de césped o el jardín tratados ayudará a prevenir su escurrimiento hacia cuerpos de agua o sistemas de drenaje.

PRIMEROS AUXILIOS

En caso de contacto con los ojos	• Mantenga el ojo abierto y enjuague lenta y delicadamente con agua durante 15-20 minutos. • Quítese los lentes de contacto, si es que usa, después de los primeros 5 minutos. Luego siga enjuagando el ojo. • Llame a un centro de control de intoxicaciones o a un médico para obtener indicaciones para el tratamiento.
En caso de contacto con la piel o la vestimenta	• Quítese las prendas contaminadas. • Enjuague la piel de inmediato con abundante cantidad de agua durante 15-20 minutos. • Llame a un centro de control de intoxicaciones o a un médico para obtener indicaciones para el tratamiento.
En caso de ingestión	• Llame a un centro de control de intoxicaciones o a un médico de inmediato para obtener indicaciones para el tratamiento. • Si la persona puede tragar, haga que beba un vaso de agua. • No induzca el vómito a menos que así se lo indique el centro de control de intoxicaciones o el médico. • No administre nada por boca si la persona está inconsciente.
En caso de inhalación	• Traslade a la persona al aire libre. • Si la persona no respira, llame al 911 o una ambulancia y hágale respiración artificial, preferentemente boca a boca, si fuera posible. • Llame a un centro de control de intoxicaciones o a un médico para obtener más indicaciones para el tratamiento.

Llamados a líneas de emergencia: Tenga el envase del producto o la etiqueta junto a usted al llamar a un centro de control de intoxicaciones o a un médico, o cuando solicite tratamiento. Para obtener información sobre este plaguicida, (incluidas cuestiones relacionadas con la salud, urgencias médicas o incidentes con el plaguicida) , llame a la *National Pesticide Telecommunications Network* (Red Nacional de Telecomunicaciones sobre Plaguicidas) al 1-800-858-7378.

INSTRUCCIONES DE USO

La utilización del producto en contrario a las indicaciones del etiquetado constituye una violación de la legislación federal.

No debe aplicarse al césped cultivado para su venta u otros destinos comerciales en forma de tepes, ni para producción comercial de semillas o destinado a la investigación.

Este producto es un herbicida selectivo de aplicación previa a la aparición que ofrece control residual de numerosas malezas herbáceas y de hoja ancha en céspedes establecidos y ornamentales (excepto *putting greens* de campos del golf). Controla las malezas susceptibles mediante la inhibición de la germinación de sus semillas y del desarrollo de raíces. Si se utiliza de acuerdo con las instrucciones, ofrece control de: zacate de agua (*Echinochloa crus-galli*), zacate azul (*Poa Annua*); anisillo (*Mollugo verticillata*), yerba del pajarero (*Stellaria media*), malezas del género Cerastio (a partir de la semilla), digitarias (*Digitaria sanguinalis y Digitaria ischaemum*), zacate egipcio, malezas del género *Erinochloa*, colas de zorro (malezas del género *Alopecurus*), pata de gallina (*Eleusine indica*)[1], zapatitos (*Lamium amplexicaule*), caminadora (*Rottboellia cochinchinensis*), zacate Johnson (*Sorghum halepense,* a partir de la semilla), arrocillo silvestre (*Echinochloa colona*); sanguinaria (*Polygonum aviculare*), coquia (*Kochia scoparia*), quelite (*Chenopodium album*), malezas del género Eragrostis, malezas del género *Panicum* (variedades *texanum, dichotomiflorum, fasciculatum*), malezas del género *Amaranthus*, verdolaga (*Portulaca oleracea*), golondrina blanca (*Richardia scabra*), cebadilla (*Bromus catharticus*)[3], bolsa de pastor (*Capsella bursa-pastoris*)[2], zacate horquetilla (*Brachiaria plantaginea*), verónica (*Veronica persica*), malezas del género *Leptochloa*; hierba de la golondrina (*Euphorbia maculata*), *Panicum capillare*, oxalis común amarilla (*Oxalis stricta*; a partir de la semilla).

1 En muchas áreas, una única aplicación de entre 5.2 y 11.9 libras/1000 pies^2 es capaz de mantener bajo control la pata de gallina (*Eleusine indica*). Sin embargo, en situaciones de gran abundancia de pata de gallina y/o temporadas prolongadas de crecimiento, puede lograrse un control más efectivo de la maleza mediante una aplicación inicial de entre 5.2 y 7.9 libras/1000 pies^2, seguida por una segunda aplicación entre 60 y 90 días después en dosis que no excedan la cantidad anual máxima (consulte la tabla de "Cantidades mínimas y máximas de aplicación por especie de hierba" en la sección *Cantidades de aplicación* de esta etiqueta).
2 Para esta maleza, aplique a fines del verano, en otoño o en invierno, antes de la germinación.
3 Sólo supresión.

En general, el césped consume entre 1 y 4 libras de nitrógeno cada 1,000 pies^2 por temporada de crecimiento. Si se usa de acuerdo con las instrucciones, este producto complementará su programa habitual de fertilización incorporando 0.92 libras de nitrógeno cada 1,000 pies cuadrados y un herbicida para controlar las malezas mencionadas. Consulte a la oficina local de difusión en caso de dudas sobre programas periódicos de fertilización.
Este producto puede manchar las entradas de automóvil, las veredas, los patios, las piscinas, etc. Barra bien inmediatamente después de cada aplicación.

CUÁNDO APLICAR SOBRE CÉSPED ARRAIGADO

Este producto debe aplicarse antes de la germinación de las semillas de la maleza, ya que no es capaz de controlar malezas arraigadas.

INSTRUCCIONES DE APLICACIÓN

Aplique el producto uniformemente, usando un esparcidor adecuado que haya sido calibrado correctamente. Se logrará un control más efectivo de la maleza si el producto es activado por al menos 0.5 pulgadas de agua de lluvia o riego o removiendo la tierra superficialmente (1-2 pulgadas) antes de la germinación de las semillas de la maleza y dentro de los 14 días posteriores a la aplicación.

PRECAUCIONES DE USO

Las siguientes precauciones de uso son aplicables al uso del producto en céspedes. Es posible que la aplicación de este producto disminuya la densidad del zacate azul que ya haya brotado y sobre los pastos sobresembrados recientemente. No aplique sobre césped sobresembrado dentro de los 60 días posteriores a la siembra o hasta el segundo corte, lo que ocurra después. Es probable que se dañen los almácigos cuyo crecimiento se desea si se aplica el producto antes de que las raíces secundarias del almácigo penetren la segunda pulgada del suelo (no de paja y suelo). No aplique sobre tepes recién colocados hasta el año siguiente. Su aplicación al césped desgastado por sequías, baja fertilidad o daños ocasionados por plagas puede perjudicar el césped. La interrupción de la barrera de herbicida mediante prácticas habituales (como la aireación del césped) puede reducir el nivel de control de la maleza. No aplique en *putting greens* ni en áreas donde la dichondra, *Agrostis capillaris, Agrostis canina* o zacate azul (*Poa Annua*) sean especies deseables. No aplicar a través de ninguna clase de sistema de riego. No aplicar por medios aéreos. No aplicar en áreas de pastoreo ni obtener forraje de las áreas tratadas.

CANTIDADES DE APLICACIÓN

Se recomienda el uso de este producto en las especies de césped indicadas en la siguiente tabla. La cantidad máxima de producto que puede aplicarse por año se indica por cada especie de césped individualmente. No exceda la cantidad máxima anual enumerada a continuación:

INSTRUCCIONES PARA EL USO DE ESPARCIDORES

1. Determine las dimensiones del área que se tratará: mida su longitud y ancho en pies y a continuación multiplíquelos para calcular el área de cobertura en pies cuadrados (longitud x ancho). Posteriormente, reste las áreas que no posean césped.
2. Utilice la tabla de configuración de esparcidores para determinar la configuración correcta.
3. Aplique uniformemente sobre el césped seco; evite las concentraciones en vetas, y las superposiciones y no omita ninguna área: es importante que la cobertura sea total. En días de alta temperatura (más de 80°F), riegue el césped para extraer las partículas que pueda contener.

Nota: Las configuraciones indicadas a la derecha para el esparcidor son aproximadas. Es importante controlar la calibración del esparcidor con frecuencia para asegurar que la máquina funcione correctamente y que el producto se aplique uniformemente. Para controlar la calibración, pese con precisión algunas libras en el esparcidor y aplíquelas sobre un área medida. A continuación, ajuste la configuración del esparcidor si fuera necesario para asegurar que la velocidad de aplicación es la correcta.

No aplique cerca de cuerpos de agua, bocas de tormenta ni zanjas de drenaje. No aplique si se han pronosticado lluvias abundantes. Aplique el producto solamente al césped y barra cualquier cantidad del producto que se deposite en la entrada del automóvil, la vereda o la calle en dirección al césped.

ALMACENAMIENTO Y DESECHO

No contamine el agua, los alimentos ni el pienso al almacenar o desechar el producto. **ALMACENAMIENTO:** Conserve en el envase original en un área de almacenamiento bajo llave, fuera del alcance de los niños y de los animales domésticos. **SI EL ENVASE ESTÁ VACÍO:** Envase no recargable: No vuelva a utilizar o rellenar este envase. Si es posible, ofrézcalo para su reciclaje. **SI EL ENVASE ESTÁ PARCIALMENTE LLENO:** Llame a la agencia local de residuos sólidos para solicitar instrucciones para desecharlo. Nunca vacíe un producto sin usar arrojando su contenido a drenajes interiores o exteriores ni al inodoro.

*Si se utiliza según se indica

OTHER RECOMMENDED Expert GARDENER LAWN PRODUCTS

A good fertilization program can keep your lawn green and vigorous. Expert Gardener has a wide selection of fertilizers to help you throughout the year.
Un buen programa de fertilización puede mantener el césped verde y vigoroso. Expert Gardener cuenta con una amplia selección de fertilizantes para brindarle ayuda durante todo el año.

EXPERT GARDENER CRABGRASS PREVENTERS

Use these pre-emergent products to control crabgrass and many other weeds in your lawn before they germinate. Crabgrass preventers are available with or without fertilizer. Use these products in early spring before weeds begin to emerge.
Use estos productos preemergentes a fin de controlar la digitaria y muchas otras malezas en su césped antes de que germinen. Existen productos para prevenir la digitaria con y sin fertilizante. Utilice estos productos a comienzos de la primavera antes de que comiencen a brotar las malezas.

EXPERT GARDENER LAWN FERTILIZER

A high nitrogen fertilizer with iron that feeds any established lawn up to 3 months. Use this product anytime during the active growing season.
Un fertilizante con alto contenido de nitrógeno y hierro que abona cualquier césped establecido hasta por 3 meses.

EXPERT GARDENER WEED & FEED

This post-emergent product controls a wide variety of existing broadleaf weeds while it feeds your desirable grass. Use weed and feeds when weeds are actively growing.
Este producto postemergente controla una amplia variedad de malezas de hojas anchas a la vez que abona el césped que usted desea.

EXPERT GARDENER SOUTHERN WEED & FEED

This fertilizer is especially formulated to control a wide variety of broadleaf weeds in Centipede and St. Augustine lawns. Use weed and feeds when weeds are actively growing.
Este fertilizante está especialmente formulado para controlar una amplia variedad de malezas de hojas anchas en céspedes de San Agustín y cienpiés.

Warranty
The directions for use of this product are believed to be adequate and must be followed carefully, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result due to such factors as weather conditions, presence or absence of other materials, or the manner of use or application, all of which are beyond the control of Gro Tec, Inc, the manufacturer, or the seller. To the extent consistent with applicable law, the products sold to you are furnished "as is" by Gro Tec, Inc. The manufacturer and the seller are subject only to the manufacturer's warranties, if any, which appear on the label of the product sold to you. Except as warranted by this label Gro Tec, Inc, the manufacturer, or the seller makes no warranties, guarantees, or representations of any kind to the buyer or the user, either express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold or use of the product, including, but not limited to, merchantability, fitness for a particular purpose or use, or eligibility of the product for any particular trade usage. To the extent consistent with applicable law, buyer's or user's exclusive remedy, and Gro Tec, Inc, the manufacturer's or the seller's total liability shall be limited to damages not exceeding the cost of the product. No agent or employee of Gro Tec, Inc, or the seller is authorized to amend the terms of this warranty disclaimer or the product's label or to make a presentation or recommendation different from or inconsistent with the label of this product. To the extent consistent with applicable law, Gro Tec, Inc, the manufacturer, or the seller shall not be liable for consequential, special, or indirect damages resulting from the use, handling, application, storage, or disposal of this product or for damages in the nature of penalties, and the buyer and the user waive any right that they may have to such damages.

Manufactured for
Gro Tec, Inc.
P. O. Box 290 • Madison, GA 30650

0 21496 00704 5

Elena Elisseeva, 2010 Used under license from Shutterstock.com

Expert GARDENER

CRABGRASS PREVENTER PLUS FERTILIZER

COVERS 15,000 SQ FT
NET WT 46.29 LB (21.00 KG)

30-0-4

CONTROLS CRABGRASS AND BROADLEAF WEEDS BEFORE THEY GERMINATE

Provides lush, green grass
Ofrece un césped verde y tupido

- Prevent crabgrass and fertilize your lawn in one application
  Impida el crecimiento de digitarias y fertilice su césped en una sola aplicación
- Up to 5 month weed and grass control
  Hasta 5 meses de control de malezas y césped

ACTIVE INGREDIENTS:
Prodiamine: (CAS 29091-21-2) 0.29%
INERT INGREDIENTS: 99.71%
Total 100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION
See back panel for further precautions and First Aid information.

COVERS 15,000 SQ FT
MONEY BACK GUARANTEED† • NET WT 46.29 LB (21.00 KG)

30-0-4
SEE BACK PANEL FOR ANALYSIS DESCRIPTION

CRABGRASS PREVENTER PLUS FERTILIZER 30-0-4