Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2015-2017 LIC. NO.
8080.3

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2012-2014 LIC. NO.

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2009-2011 LIC. NO.

LICENSED
HAWAII
DEPT. OF AGRICULTURE
Period 2003-2005 No. 8080.3

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2006-2008 LIC. NO.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

CRITICAL CARE Disinfectant, Fungicidal & Virucidal Spray is a colorless, odorless broad spectrum antimicrobial disinfectant and deodorizer. Proven to kill bacteria, fungus and viruses*, **CRITICAL CARE Disinfectant, Fungicidal & Virucidal Spray** should be used on non-porous environmental hard surfaces in homes, hospitals, nursing homes, medical and dental clinics, laboratories, ambulance and patient transfer vehicles, funeral homes, hotels, restaurants, schools, day care facilities, offices, veterinary clinics, animal shelters, kennels, exercise facilities, beauty and barber shops, subways, trains, airplanes, ships, busses and other public transportation vehicles, locker rooms, kitchens and restrooms.

CRITICAL CARE Disinfectant, Fungicidal & Virucidal Spray has been formulated to disinfect hard, non-porous environmental surfaces (painted, glazed tile, plastic, metal, glass, glazed porcelain) and objects such as walls, floors, counters, sinks, toilets, cabinets, tubs, showers, doorknobs, light switch covers, telephones, appliances, stove tops, bed frames, wheelchairs, over-bed tables, examination tables and waste containers, tables, and chairs.

Fungicidal Activity: CRITICAL CARE Disinfectant, Fungicidal & Virucidal Spray is effective against *Trichophyton mentagrophytes*, the Athlete's foot fungus, Use in locker rooms, dressing rooms, showers and bath areas, and exercise facilities.

Deodorizes: CRITICAL CARE Disinfectant, Fungicidal & Virucidal Spray reduces annoying odors caused by bacteria. Use to control odors in hospitals, nursing homes, public restrooms, animal kennels and barn stalls. In private homes, use in kitchen, bathroom, sink rooms and basement.

EnvirOx
Critical Care

Disinfectant, Fungicide & Virucide*
Disinfects and Deodorizes

RESTAURANTS • HOSPITALS • SCHOOLS • HOMES • OFFICES

No harsh chemical smell • No mixing required • No dulling residue

Fast, Easy, Effective

General Information

Critical Care Disinfectant, Fungicidal & Virucidal Spray successfully killed the following organisms under AOAC protocols (In order to ensure that all organisms listed are killed, you must use the contact times as directed in the Application Instructions)

Organism	Kill Time
Pseudomonas aeruginosa [1]	30 seconds
Staphylococcus aureus [1]	30 seconds
Salmonella cholerasuis [1]	30 seconds
Listeria monocytogenes [1]	30 seconds
Vancomycin resistant *Enterococcus faecium* [1]	2 minutes
Methicillin resistant *Staphylococcus aureus* [1]	2 minutes
Escherichia coli 0157:H7 [1]	2 minutes
Trichophyton mentagrophytes (Athlete's Foot Fungus)	10 minutes
HIV type 1-Strain HTLV IIIB [1]	30 seconds
Herpes Simplex Type 1 VR-733 F(1) Strain [2]	1 minute
Influenza A VR-544, Hong Kong Strain [2]	10 minutes
Rhinovirus R37 VR-1147, Strain 151-1 [2]	10 minutes
Polio Type 2, VR-1002 Lansing Strain [2]	10 minutes

1 Evaluated in the presence of 5% organic soil. 2 Evaluated in the presence of 1% organic soil.

Distributed by: EnvirOx LLC
1938 E. Fairchild Street, Danville, IL. 61832
EPA Reg. No. 72977-3-69268
EPA Est. No. 69268-IL-001
Net Vol.:32 oz.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Active Ingredient

Silver*	0.003%
Citric Acid	4.840%
Other ingredients	95.157%
Total	100.000%

* Electrolytically generated silver ions

APPLICATION INSTRUCTIONS

Surfaces that are heavy soiled with organic matter must be pre-cleaned prior to using this product.

General Disinfection: For general disinfection and control of bacteria such as *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Salmonella cholerasuis*, *Listeria monocytogenes*, Vancomycin Resistant *Enterococcus faecium* (VRE), Methicillin Resistant *Staphylococcus aureus* (MRSA) and *Escherichia coli* 0157:H7 the surface must be completely wet with **CRITICAL CARE Disinfectant, Fungicidal & Virucidal Spray** for 2 minutes. The surface may then be wiped dry with a clean towel. When used as directed, **CRITICAL CARE Disinfectant, Fungicidal & Virucidal Spray** provides protection from *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Salmonella cholerasuis* up to 24 hours after initial application.

Fungus Control: For effective control of fungus such as *Trichophyton mentagrophytes*, the surface must be completely wet with **CRITICAL CARE Disinfectant, Fungicidal & Virucidal Spray** for 10 minutes. The surface may then be wiped dry with a clean towel. Re-apply when cleaning or when new growth appears.

Viral Control: To kill Herpes Simplex Type 1 F(1) Strain, Influenza A Virus, Hong Kong strain, Rhinovirus R37 Strain 151-1, Polio Virus Type 2 Lansing Strain the surface must be completely wet with **CRITICAL CARE Disinfectant, Fungicidal & Virucidal Spray** for 10 minutes. The surface may then be wiped clean with a towel.

Kills HIV-1 on pre-cleaned environmental surfaces/objects previously soiled with bloody/body fluids in health care settings (or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV): **Instructions for Cleaning and Decontamination Against HIV** on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids: **Personal Protection:** When handling items soiled with blood/body fluids, use appropriate barrier protection such as latex gloves, gowns, masks or eye coverings. **Cleaning Procedure:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this disinfectant. **Contact time:** Apply **CRITICAL CARE Disinfectant, Fungicidal & Virucidal Spray** to area to be treated. The surface must completely be wet with **CRITICAL CARE Disinfectant, Fungicidal & Virucidal Spray** for 30 seconds. The surface may then be wiped dry with a clean towel. **Disposal of Infectious Materials:** Blood and other body fluids should be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the human body, either into or in contact with the bloodstream, or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not normally penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

Storage and Disposal	
Storage	Do not contaminate water, food or feed by storage or disposal.
Disposal	Do not reuse container. Rinse thoroughly before discarding in trash or recycling.

In Case of Emergency

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact CHEMTREC 1-800-424-9300 for emergency medical treatment information.

1 2003