Department of Agriculture
STATE OF HAWAII
LICENSED
PERIOD 2015-2017 LIC. NO. 8109.32

Department of Agriculture
STATE OF HAWAII
LICENSED
PERIOD 2012-2014 LIC. NO.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER.

KEEP OUT OF REACH OF CHILDREN: Corrosive. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Wear protective eyewear (goggles, face shield or safety glasses). Wear protective clothing and rubber gloves. Avoid contamination of food. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT – FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

For additional medical advice, call the following emergency phone number: 1-800-328-0026.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

PHYSICAL OR CHEMICAL HAZARDS

Combustible: Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Do not store on side. Avoid crossing or inspecting of side walls. Store securely in closed original container. Avoid storage at temperature extremes or in sunlight. Avoid shipping or storing below freezing. If product freezes, thaw at room temperature and shake gently to remix components. Use locked storage in an area that will prevent cross-contamination of other pesticides, fertilizer, food, and feed. Store in locked area inaccessible to children.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture rinsate is a violation of Federal Law. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Discard Rinsate. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Nonrefillable container: Do not reuse or refill this container. Offer for recycling if available.

EPA Reg. No. 67170-1-42964
EPA Est. 3053-IN-1 (L), 1677-IL-2 (J), 1677-MIN-1 (P), 1677-CA-1 (S), 1677-TX-1 (D), 1677-GA-1 (M), 1677-CA-2 (R), 1677-WV-1 (V).
*Supercript, retain to first letter of date code.

This product is effective against the following microorganisms:
Contact time is 10 minutes unless noted as below:

Isolates From AIDS Patients
Aspergillus niger
Candida albicans
Cryptococcus neoformans
Pseudomonas aeruginosa
Staphylococcus aureus
Streptococcus pneumoniae

Gram Positive Clinical Isolates
Enterococcus faecalis
Micrococcus luteus
Staphylococcus aureus
Staphylococcus aureus (Toxic shock)
Staphylococcus epidermidis
Staphylococcus saprophyticus
Streptococcus haemolyticus

Gram Negative Clinical Isolates
Acinetobacter calcoaceticus var. anitratus
Acinetobacter calcoaceticus var. lwoffii
Bordetella bronchiseptica
Brevundimonas diminuta
Burkholderia cepacia
Enterobacter agglomerans
Enterobacter cloacae
Enterobacter gergoviae
Enterobacter liquefaciens
Escherichia coli (Urinary)
Escherichia coli (Wound)
Flavobacterium meningosepticum
Klebsiella oxytoca
Klebsiella pneumoniae
Morganella morganii
Proteus mirabilis
Proteus vulgaris
Pseudomonas aeruginosa
Pseudomonas fluorescens
Pseudomonas pseudomallei
Pseudomonas putida
Pseudomonas stutzeri
Serratia marcescens
Sphingomonas paucimobilis

Other Bacteria
Actinobacillus pleuropneumoniae
Actinomyces pyogenes
Bordetella bronchiseptica
Burkholderia pickettii
Campylobacter jejuni
Corynebacterium pseudotuberculosis
Enterococcus faecalis
Escherichia coli
Escherichia coli strain O157:H7
Escherichia vulneris
Listeria monocytogenes
Pasteurella haemolytica
Pseudomonas aeruginosa
Rhodococcus equi
Salmonella enterica
Salmonella typhi
Staphylococcus aureus
Staphylococcus aureiscolis
Staphylococcus capitis
Staphylococcus hominis
Staphylococcus simulans
Streptococcus equi var. equi
Streptococcus equi var. zooepidemicus
Streptococcus pyogenes
Yersinia enterocolitica

Environmental Fungi
Aspergillus candidus
Aspergillus niger
Penicillium chermesinum
Penicillium oxalicum
Penicillium spinulosum
Discfadium sp.

Antibiotic Resistant Gram Positive Bacteria
Enterococcus faecalis (Vancomycin Resistant-VRE)
Staphylococcus aureus (Methicillin Resistant-MRSA)

Human Viruses
Cytomegalovirus
HBV (Hepatitis B Virus)
HCV (Hepatitis C Virus)
Herpes Simplex type 2 Virus
HIV-1 (AIDS Virus) 30 seconds contact time
Human Coronavirus
Influenza A/ Brazil Virus
Influenza B Virus (Allen strain)
Influenza C Virus (Taylor strain)
Measles Virus
Parainfluenza Virus type 1
Poliovirus type 1 (Chat strain) 30 minutes contact time
Respiratory Syncytial Virus
Rotavirus

Non-Human Viruses
Avian Influenza/Turkey/Wisconsin Virus
Canine Coronavirus
Equine Virus Influenza
Feline Calicivirus
Infectious Bovine Rhinotracheitis Virus (IBR)
Newcastle Disease Virus
Porcine Parvovirus
Pseudorabies Virus
Transmissible Gastroenteritis Virus (TGE)
T1 bacteriophage
T4 bacteriophage
Bovine Viral Diarrhea Virus (BVDV)
Norovirus
Avian Influenza Virus (H5N1)

ECOLAB®

HB Quat

Cleaner • Disinfectant • Detergent • Deodorizer • Fungicide • Virucide*

COMPLETE CLEANING SYSTEM
OASIS®

14673
Net Contents: 2.5 U.S. gal/9.46 L

ACTIVE INGREDIENTS:
Alkyl (60% C14; 30% C12; 5% C12; 5% C16)
dimethyl benzyl ammonium chloride 2.37%
Alkyl (68% C12; 32% C14)
dimethyl ethylbenzyl ammonium chloride 2.37%
INERT INGREDIENTS: 95.26%
TOTAL: 100.00%

KEEP OUT OF REACH OF CHILDREN
DANGER *See side panel for Precautionary Statements and First Aid*

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

To be used in hospitals in the following areas as a disinfectant: operating rooms, patient care rooms & facilities, recovery, anesthesia, ER, radiology, X-ray cat labs, newborn nurseries, orthopedics, respiratory therapy, surgi-centers, labs, blood collection rooms, central supply, housekeeping & janitorial rooms, nursing homes, doctors offices & labs, dentists offices & labs (dental operatories).

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

BACTERICIDAL STABILITY OF USE-DILUTION: Tests confirm that this product, when diluted in 400 ppm hard water and in the presence of 5% soil load, remains effective against *Pseudomonas aeruginosa, Staphylococcus aureus, Salmonella enterica* for up to 64 days when stored in a sealed container at room temperature.

If the use-dilution product becomes visibly dirty or contaminated, the use-dilution must be discarded and a fresh product prepared.

Always use clean, properly labeled dry containers when diluting the product. Bactericidal stability of the use-dilution does not apply to open containers such as buckets or pails. Use-dilution product in open containers must be prepared daily or more often if the solution becomes visibly dirty or diluted or contaminated.

DISINFECTION/FUNGICIDAL/*VIRUCIDAL* DIRECTIONS: Apply use solution to hard inanimate, non-porous surfaces thoroughly wetting surfaces with a cloth, mop, sponge or sprayer. For heavily soiled areas, a preliminary cleaning is required. For spraypr applications use a coarse spray device. Spray 6-8 inches from surface and rub with brush, sponge or cloth. Do not breathe spray.

Add 2 ounces per gallon (16 milliliters per liter) of water to disinfect hard, non-porous surfaces. Treated surfaces must remain wet for 10 minutes. Prepare a fresh solution at least daily or when use dilution becomes diluted or soiled.

At 2 ounces per one gallon (or equivalent use dilution of 1:64) in the presence of 300 ppm hard water (CaCO3). This product was proven to be effective against Hepatitis B Virus with a contact time of 10 minutes.

KILLS HCV, HBV AND HIV-1 ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HCV (Hepatitis C Virus), HBV (Hepatitis B Virus) or HIV-1 (AIDS Virus).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HCV, HBV AND HIV-1 OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

PERSONAL PROTECTION: Specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable latex gloves, gowns, masks or eye coverings.

© Airkem Professional Products, Division of Ecolab Inc.
370 Wabasha Street N, St. Paul 55102-1300 USA
© 2010 Ecolab Inc. Made in U.S.A. 717848/5401/0410
All Rights Reserved

CLEANING PROCEDURE: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids should be autoclaved and disposed of according to Federal, State and local regulations for infectious waste disposal. CONTACT TIME: Leave surfaces wet for 30 seconds for HIV-1 and 10 minutes for HCV and HBV. The contact time for the viruses and bacteria listed on this label is 10 minutes except for Poliovirus Type 1 (Chat strain), which is 30 minutes.

FUNGICIDAL: At 2 ounces per gallon use-level, (or equivalent use dilution) is effective against the pathogenic fungus *Trichophyton mentagrophytes* ("athlete's foot fungus – cause of Ringworm) on inanimate surfaces in the presence of 5% organic soil load and 300 ppm water hardness as CaCO3 in locker rooms, dressing rooms, shower and bath areas and exercise facilities. Contact time — 10 minutes.

This product, in the presence of a — 100% organic soil load, diluted 1:64 (2 ounces per gallon) in 3SS ppm Hard Water, demonstrated efficacy within 10 minutes against the following pathogenic fungus: *Trichophyton mentagrophytes*. Note that the organism referenced in the previous statement is not associated with blood spills. For blood spills, the surface must be thoroughly cleaned before applying this product.

VIRUCIDAL*: When used on inanimate, hard, non-porous, environmental surfaces at 2 ounces per gallon of water for a 10 minute contact time (5% organic soil), except for Poliovirus type 1 (Chat strain) which requires a 30 minute contact time (5% organic soil) and HIV-1 which requires only a 30 second contact time. This product, in the presence of 98% organic soil load, diluted 1:64 (2 ounces per gallon) in 400 ppm Hard Water, demonstrated efficacy within 10 minutes against the following virus: *Human Coronavirus*. Note that the organism referenced in the above statement is not associated with blood spills. For blood spills, the surface must be thoroughly cleaned before applying this product.

General Deodorization: To deodorize, add 2 ounces of this product per gallon of water (or equivalent use dilution of 1:64). Excess material must be wiped up or allowed to air dry.

DISINFECT TOILET BOWLS

NON-ACID TOILET BOWL (AND URINAL) DISINFECTION/CLEANER DIRECTIONS

Remove gross filth prior to disinfection.

From Concentrate: Add 2 ounces (16 milliliters per liter) to the toilet bowl and mix. Brush thoroughly over exposed surfaces and under the rim with a cloth, mop, or sponge. Repeat and allow to stand for 10 minutes and flush.

From Use Solution: Empty toilet bowl or urinal and apply 2 ounces per gallon (16 milliliters per liter) use solution to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer, brush or swab thoroughly and allow to stand for 10 minutes and flush.

For Heavy Duty Use: Empty toilet bowl or urinal and apply 8 ounces per gallon (64 milliliters per liter) use solution to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer, brush or swab thoroughly and allow to stand for 10 minutes and flush.

Cleaning/Deodorizing Directions: Add 1 to 4 ounces per gallon (8 to 32 milliliters per liter) of water to clean and deodorize windows, mirrors and non-food contact glass surfaces. When using a coarse spray device, spray 6-8 inches from surface. Rub with sponge or cloth. Do not breathe spray.

Veterinary Clinics/Animal Life Science Laboratory/Zoos/Pet Shop/Kennels/Breeding and grooming Establishment/Tack Shops

Disinfection Directions: For cleaning and disinfecting the following hard non-porous surfaces: equipment, utensils, instruments, cages, kennels, stables, stalls and catteries. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 2 ounces per gallon of water (or equivalent dilution) for a period of 10 minutes. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals as well as forks, shovels, scrappers, used in removing litter and manure. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, automatic feeders, waterers and other equipment which dispenses food or water with soap or detergent, and rinse with potable water before reuse.

ECOLAB®

HB Quat

Cleaner • Disinfectant • Detergent • Deodorizer • Fungicide • Virucide*

COMPLETE CLEANING SYSTEM
OASIS®

14673
Net Contents: 2.5 U.S. gal/9.46 L

ACTIVE INGREDIENTS:
Alkyl (60% C14; 30% C12; 5% C12; 5% C16)
dimethyl benzyl ammonium chloride 2.37%
Alkyl (68% C12; 32% C14)
dimethyl ethylbenzyl ammonium chloride 2.37%
INERT INGREDIENTS: 95.26%
TOTAL: 100.00%

KEEP OUT OF REACH OF CHILDREN
DANGER See side panel for Precautionary Statements and First Aid

DIELINE #
574129

FILE NAME	DIELINE # / SIZE	CONTAINER COLOR	COLORS (#)	NOTES
717848/54	574129	WHITE	GCMI 226	
PACKAGING SUBSTRATE (LABEL TYPE)			GCMI 90	
POLYSLEEVE				