Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2012-2014 LIC. NO.

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2015-2017 LIC. NO.

8126.119

BAYER ADVANCED™

100% SATISFACTION GUARANTEED

Weed Killer for Lawns

CONCENTRATE

KILLS BROADLEAF WEEDS

MATA LAS MALEZAS DE HOJA ANCHA

¿HABLA ESPAÑOL? VEA EL PANEL TRASERO

- Rainproof in 1 Hour
- Won't harm lawns†

†When used as directed

Kills 200+ Weeds

Active Ingredients:
2,4-D, dimethylamine salt 7.59%
Mecoprop-p, dimethylamine salt ... 1.83%
Dicamba, dimethylamine salt 0.84%
Other ingredients 89.74%
Total 100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION See back panel booklet for additional precautionary statements

Treats up to 16,000 sq ft

NET CONTENTS 32 FL OZ (946mL)

79659326 R.0

Book Placement

3.50”

BAYER ADVANCED

Weed Killer for Lawns

CONCENTRATE

This product kills over 200 types of broadleaf weeds. This formula kills Dandelion, Clover and Chickweed without harming your lawn†. Controlling weeds in your lawn has never been easier.

PRECAUTIONARY STATEMENTS

CAUTION Hazards to Humans and Domestic Animals

- Harmful if absorbed through skin or if swallowed.
- Causes moderate eye irritation.
- Avoid contact with eyes, skin or clothing.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

After using this product:

- Rinse gloves before removing.
- Remove clothing and launder separately before reuse.
- Wash hands thoroughly with soap and water.
- If product gets on clothing, remove clothing as soon as possible and shower.

FIRST AID ✚

IF SWALLOWED	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.
IF IN EYES	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	• Move person to fresh air. • If not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor’s office, bring this product label to show medical personnel.
For medical emergency information, you may call toll-free 1-877-229-3763.
Active Ingredients: 7.59% 2,4-D, dimethylamine salt [CAS # 2008-39-1], 1.83% Mecoprop-p, dimethylamine salt [CAS # 66423-09-4], and 0.84% Dicamba, dimethylamine salt [CAS # 2300-66-5].

ENVIRONMENTAL HAZARDS

- This product is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.
- Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.
- This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

NOTICE: Research and testing have determined that the "Directions For Use" are appropriate for the proper use of this product under expected conditions. The Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

***Money Back Guarantee:** If you are not satisfied with this product, we will gladly refund your original purchase price.
†When used as directed.

Produced for: Bayer Advanced
A Business Unit of Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, NC 27709

EPA Reg. No. 72155-4
EPA Est. No indicated by 3rd and 4th digits of the batch number on this package. (75) = 5905-AR-1 (71) = 2217-KS-2 (39) = 58996-MO-1 (57) = 071106-GA-003

Bayer (reg'd), Bayer Cross (reg'd), and Bayer Advanced™ are trademarks of Bayer.

PRESS TO RESEAL

79720912b R.1

Book Placement 6.375”

8.125”

4.0”

6/28/11

CCL LABEL YOUR GLOBAL PARTNER	Description: BOTPE 40oz. ASB_1_REV2- CONSTRUCTION		NOTES	LEGEND
	Supplied for: Bayer	File No.: CCL4951		Copy Image Area
	Finished Size: 8.125” (H) x 4.0” (W)	Supersedes: CCL4931		Cut/Fold Lines
	Date: 5/6/09	Supplied by: CCL Label Memphis		
	Application: Illustrator CS2	Prepared By: NR	Measurements are in Inches	

MANTENER FUERA DEL ALCANCE DE LOS NIÑOS

DECLARACIONES DE PRECAUCIÓN

PRECAUCIÓN **Riesgos para seres humanos y animales domésticos**

• Nocivo si se absorbe por la piel o se ingiere. • Causa irritación moderada en los ojos. • Evitar el contacto con la piel, los ojos o la ropa. • Al utilizar este producto, usar camisa de mangas largas, pantalones largos, medias, zapatos y guantes de goma.

Después de utilizar este producto:
• Enjuagar los guantes antes de quitárselos. • Quitarse la ropa y lavarla en forma separada antes de volver a usar.
• Lavarse las manos prolijamente con agua y jabón. • Si el producto entra en contacto con la ropa, quitarse la ropa lo antes posible y ducharse.

PRIMEROS AUXILIOS ✚

EN CASO DE INGESTIÓN	• Llamar inmediatamente a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento. • Hacer que la persona beba tragos de agua, si puede tragar. • No inducir el vómito a menos que lo indique el médico o el centro de control de envenenamiento. • No administrar nada por boca a una persona inconsciente.
EN CASO DE CONTACTO CON LOS OJOS	• Mantener los ojos abiertos y enjuagar en forma lenta y suave con agua durante 15 a 20 minutos. • Retirar los lentes de contacto, si los hubiera, después de los primeros 5 minutos, y luego continuar enjuagando los ojos. • Llamar a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento.
EN CASO DE CONTACTO CON LA PIEL O LA ROPA	• Quitar la ropa contaminada. • Enjuagar la piel inmediatamente con abundante agua durante 15 a 20 minutos. • Llamar a un médico o un centro de control de envenenamiento para obtener asesoramiento médico.
EN CASO DE INHALACIÓN	• Trasladar a la persona a un lugar fresco. • Si hay paro respiratorio, llamar al 911 o a una ambulancia, luego proporcionar respiración artificial, preferentemente de boca a boca, si fuera posible. • Llamar a un médico o un centro de control de envenenamiento para mayores consultas sobre el tratamiento.

NOTA: Al llamar a un centro de control de envenenamiento, tener la etiqueta del producto a mano. En caso de solicitar tratamiento en una sala de emergencias o consultorio médico, llevar la etiqueta del producto para mostrarla al personal médico.
Para solicitar información médica de emergencia, llamar al número gratuito 877-229-3763.
Ingredientes activos: 7.59% 2,4-D, sal dimetilamina [CAS Nº 2008-39-1], 1.83% mecoprop-p, sal dimetilamina [CAS Nº 66423-09-4] y 0.84% dicamba, sal dimetilamina [CAS Nº 2300-66-5].

RIESGOS AMBIENTALES

• Este producto es tóxico para los peces e invertebrados acuáticos y puede afectar a plantas que no son el objetivo del tratamiento. No aplicar directamente sobre el agua, o en lugares donde haya agua en la superficie, o en áreas donde haya marea por debajo de la marca media de la altura del agua.
• La deriva y el residuo líquido pueden ser perjudiciales para los organismos acuáticos que se encuentran en aguas adyacentes a las áreas tratadas.
• Para proteger el medio ambiente, no deje que el pesticida se filtre o ingrese a las bocas de tormenta, desagües, alcantarillas o aguas superficiales. Aplicar este producto cuando el clima esté calmo y no haya probabilidades de lluvia para las próximas 24 horas ayudará a garantizar que el viento o la lluvia no dispersen o escurran el pesticida del área tratada. Enjuagar el equipo de aplicación usado sobre el área tratada evitará que ingrese en cursos de agua o sistemas de desagüe.
• Esta sustancia química tiene propiedades y características relacionadas con las sustancias químicas que se encuentran en las aguas subterráneas. El uso de esta sustancia en áreas donde los suelos son permeables, en particular donde la capa freática es poco profunda, puede causar contaminación de las aguas subterráneas. La aplicación en cercanías a un tanque de agua o pozo puede contaminar el agua potable o las aguas subterráneas.

NOTA: Investigaciones y pruebas han determinado que las "Indicaciones de uso" son las adecuadas para el uso apropiado de este producto bajo las condiciones previstas. El Comprador asume la responsabilidad por la falta de rendimiento o seguridad si no es utilizado de acuerdo con las indicaciones.
***Garantía de devolución de dinero:** Si usted no está satisfecho con este producto, gustosamente le reembolsaremos el monto original del precio de compra.

†Cuando se use según las indicaciones de uso.

page 8

READ THE LABEL FIRST!

Indicaciones en español después de las de inglés

OPEN
Resealable Label for Directions & Precautions

READ THE LABEL FIRST

QUICK FACTS

- **Kills Over 200 Broadleaf Weeds****
- **Won't Harm Your Lawn†**
- **Treats Up To 16,000 Sq. Ft.**

KILLS	Over 200 broadleaf weeds such as Dandelion, Clover, Chickweed and other listed weeds ** as listed on this label
WHERE TO USE	**On Lawns** †(When used as directed. Do not use on Carpetgrass, Dichondra or Floratam variety of St. Augustinegrass.)
WHEN TO USE	When weeds are actively growing
HOW MUCH TO USE	**Northern Lawns:** Fescue, Kentucky Bluegrass, and Perennial Ryegrasses **and** **Hardy Southern Lawns:** Bermuda, Buffalo and Zoysia grasses 2 fl. oz. per 500 sq. ft. **Southern Lawns:** Bahia, Centipede and St. Augustinegrass grasses 1 fl. oz. per 500 sq. ft.
☎ 💻	For questions or comments, call toll-free 1-877-BayerAG (1-877-229-3724), or visit us on-line at www.BayerAdvanced.com

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	7.59%
Mecoprop-p, dimethylamine salt	1.83%
Dicamba, dimethylamine salt	0.84%
OTHER INGREDIENTS	89.74%
TOTAL	100.00%

THIS PRODUCT CONTAINS:
0.00427 lb 2,4-D acid equivalent per fl. oz. or 6.30%
0.00102 lb Mecoprop-p acid equivalent per fl. oz. or 1.51%
0.00047 lb Dicamba acid equivalent per fl. oz. or 0.69%
Isomer specific by AOAC Method

EPA Reg. No. 72155-4
EPA Est. No indicated by 3rd and 4th digits of the batch number on this package.
(75) = 5905-AR-1 (71) = 2217-KS-2 (39) = 58996-MO-1
(57) = 071106-GA-003

FOR OUTDOOR RESIDENTIAL USE ONLY

6 87073 04160 6

79720912b R.1

front

6.375”

3.5” 3.5”

6/28/11

CCL LABEL
YOUR GLOBAL PARTNER

Description: BOTPE 40oz. ASB_1_REV2- BOOK OUT			
Supplied for: Bayer	**File No.:** CCL4951	Total number of pages must be divisible by four (4).	Copy Image Area Cut/Fold Lines
Finished Size: 8.125” (H) x 4.0” (W)	**Supersedes:** CCL4931		
Date: 5/6/09	**Supplied by:** CCL Label Memphis	Measurements are in Inches	
Application: Illustrator CS2	**Prepared By:** NR		

NOTES

LEGEND

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Read and follow all label directions.
Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes.
Do not apply through any type of irrigation system.

BEFORE YOU USE
Read and follow these directions when using:

- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn or garden area only.
- Spray when most of the day is expected to remain cooler than 90˚F.
- Do not use this product on or near desirable plants, including within the dripline of desirable trees and shrubs, since injury may result.
- The maximum number of broadcast applications per treatment site is two per year.

HOW TO USE
Shake Well Before Use

- Determine amount of product to use based on grass type and size of area to be treated.
- Prepare product for use according to How To Apply.
- Apply to lawn.

KILLS

Annual yellow sweetclover
Aster
Austrian fieldcress
Bedstraw
Beggarticks
Betony, Florida
Bindweed, field
Bird vetch
Bitter wintercress
Bittercress, hairy
Bitterweed
Black-eyed Susan
Black medic
Black mustard
Blackseed plantain
Blessed thistle
Bloodflower milkweed
Blue lettuce
Blue vervain
Bracted plantain
Brassbuttons
Bristly oxtongue
Broadleaf dock
Broadleaf plantain
Broomweed
Buckhorn
Buckhorn plantain
Bulbous buttercup
Bull thistle
Bullnettle
Burclover
Burdock
Burning nettle
Burweed
Buttercup
Buttonweed
Canada thistle
Carolina geranium
Carpetweed
Catchweed bedstraw
Catnip
Catsear
Chickweed, common
Chickweed, mouseear
Chicory
Cinquefoil
Clover, crimson
Clover, hop
Clover, red
Clover, strawberry
Clover, sweet
Clover, white
Cockle
Cocklebur
Common mullein
Creeping Charlie
Creeping Jenny
Cudweed
Curly dock
Daisy, English
Daisy, fleabane
Daisy, oxeye
Dandelion
Dichondra
Dogbane
Dogfennel
Dollarweed
Elderberry
False dandelion
False flax
False sunflower
Fiddleneck
Florida pusley
Frenchweed
Galinsoga
Goathead
Goldenrod
Ground ivy
Gumweed
Hairy fleabane
Hawkweed
Healall
Heartleaf drymary
Heath aster
Hedge bindweed
Hedge mustard
Hemp
Henbit
Hoary cress
Hoary plantain
Hoary vervain
Horsenettle
Jimsonweed
Knawel
Knotweed
Kochia
Lambsquarters
Lespedeza
Mallow
Matchweed
Mexicanweed

(con't)

page 2

DIRECTIONS FOR USE *(cont'd)*

KILLS (cont'd)

Milk vetch
Morningglory
Mouseear hawkweed
Mugwort
Musk thistle
Mustard
Narrowleaf plantain
Narrowleaf vetch
Nettle
Orange hawkweed
Oriental cocklebur
Oxalis
Parsley-piert
Parsnip
Pearlwort
Pennycress
Pennywort
Pepperweed
Pigweed
Pineywoods bedstraw
Plains coreopsis (tickseed)
Plantain
Poison ivy
Poison oak
Pokeweed
Poorjoe
Prairie sunflower
Prickly lettuce
Prickly sida
Prostrate knotweed
Prostrate pigweed
Prostrate spurge
Prostrate vervain
Puncturevine
Purslane, common
Ragweed
Red sorrel
Redroot pigweed
Redstem filaree
Rough cinquefoil
Rough fleabane
Russian pigweed
Russian thistle
Scarlet pimpernel
Scotch thistle
Sheep sorrel
Shepherdspurse
Slender plantain
Smallflower galinsoga
Smooth dock
Smooth pigweed
Sorrel
Sowthistle
Spanishneedles
Speedwell
Spiny amaranth
Spiny cocklebur
Spiny sowthistle
Spotted catsear
Spotted spurge
Spurweed
Stinging nettle
Strawberry, India mock
Tall nettle
Tall vervain
Tansy mustard
Tanweed
Thistle
Trailing crownvetch
Tumble mustard
Tumble pigweed
Velvetleaf
Venice mallow
Virginia buttonweed
Virginia creeper
Virginia pepperweed
Water pennywort
Wavyleaf bullthistle
Western clematis
Western salsify
White mustard
Wild aster
Wild buckwheat
Wild carrot
Wild four-o'clock
Wild garlic
Wild geranium
Wild lettuce
Wild marigold
Wild mustard
Wild onion
Wild parsnip
Wild radish
Wild rape
Wild strawberry
Wild sweet potato
Wild vetch
Woodsorrel
Woolly croton
Woolly morningglory
Woolly plantain
Wormseed
Yarrow
Yellow rocket
Yellowflower pepperweed and other broadleaf weeds

HOW MUCH TO APPLY

Northern Lawns:
Fescue, Kentucky Bluegrass, and Perennial Rye - 2 fl. oz. (4 Tbs.) per gallon of water

Hardy Southern Lawns:
Bermuda, Buffalo and Zoysia grasses - 2 fl. oz. (4 Tbs.) per gallon of water

Southern Lawns:
Bahia, Centipede, and St. Augustinegrass - 1 fl. oz. (2 Tbs.) per gallon of water

For lawns with mixed grass types, use the lowest rate.

Each gallon of solution treats 500 sq. ft.

TIP: Measure length and width of area to be treated in feet. Multiply these 2 numbers together. This is the area to be treated in square feet (sq. ft.)

Application Restrictions:

- Limited to **2 applications per year** with a **30 day interval** between applications
- **Maximum single application rate** of 8.0 fl. oz./1,000 sq. ft. (1.47 lbs 2,4-D ae, 0.35 lb MCPP-p ae and 0.16 lb dicamba ae per acre per application)
- **Maximum seasonal rate** of 16.0 fl. oz./1,000 sq. ft. (2.94 lbs 2,4-D ae, 0.71 lb MCPP-p ae and 0.33 lb dicamba ae per acre per application)

(con't)

page 3

6.375"

3.5" 3.5"

6/28/11

CCL LABEL YOUR GLOBAL PARTNER	**Description:** BOTPE 40oz. ASB_1_REV2- BOOK IN		**NOTES**	**LEGEND**
	Supplied for: Bayer	**File No.:** CCL4951	Total number of pages must be divisible by four (4).	Copy Image Area Cut/Fold Lines
	Finished Size: 8.125" (H) x 4.0" (W)	**Supersedes:** CCL4931		
	Date: 5/6/09	**Supplied by:** CCL Label Memphis	Measurements are in Inches	
	Application: Illustrator CS2	**Prepared By:** NR		

DIRECTIONS FOR USE *(cont'd)*

HOW TO APPLY

PUMP STYLE SPRAYERS

1. Add the appropriate amounts of water and concentrate to the sprayer tank.
2. Close sprayer, shake well and pressurize.
3. Adjust nozzle to a coarse spray pattern.
4. Large Areas - Apply evenly over area.
   Individual Weeds - Hold tip of wand about 12 inches from weed, aim at center of weed and spray.
5. Occasionally re-pressurize the sprayer if needed to maintain a good spray pattern.

DIAL STYLE HOSE-END SPRAYERS

1. Fill sprayer jar with enough concentrate to cover area to be treated.
2. Close sprayer and set dial.
   TIP: The numbers on Dial-Style Hose-End sprayers are ounces of concentrate per gallon of spray.
3. Attach sprayer to hose and apply.
4. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.

CONCENTRATE-PLUS-WATER HOSE-END SPRAYER

1. Pour concentrate into sprayer jar to the required fluid ounce level.
2. Add the appropriate amount of water to the gallons level.
3. Replace sprayer top and shake well before applying.
4. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.

HOW IT WORKS

This product is a *selective* herbicide – it *selectively* kills weeds in the lawn but won't harm the lawn itself. It is a blend of proven weed killers designed to get broadleaf weeds such as dandelion, clovers and chickweed.

FOR BEST RESULTS

- Apply when weeds are young and actively growing.
- If the lawn is dry, water 24 hours before using this product.
- Rain or watering one hour after application will not wash away effectiveness.
- Apply 1 to 3 days after mowing.
- Certain hard-to-kill weeds may need to be spot-treated in 30 days.
- Do not apply to new lawns until the grass has reached a height of at least 2 inches.
- Bare spots may be seeded 3 to 4 weeks after application.

Do not allow people (other than applicator) or pets on treated area during application.

Do not allow people or pets to re-enter the treated area until spray has dried.

IMPORTANT- Directions for Storage and Disposal

Do not contaminate food or feed by storage and disposal.

PESTICIDE STORAGE

Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING

If Empty:

- Non-refillable container. Do not refill or reuse empty container.
- Place empty container in trash or offer for recycling if available.

If Partly Filled:

- If it is necessary to dispose of unused product please call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions.
- Never place unused product down any indoor or outdoor drain.

¡LEER LA ETIQUETA ANTES DE UTILIZAR EL PRODUCTO!

DATOS RÁPIDOS

- **Mata más de 200 malezas de hoja ancha**
- **No daña el césped†** • **Trata hasta 16,000 pies cuadrados (1,500 m2)**

MATA	Más de 200 malezas de hoja ancha como el diente de león, trébol, capiqui y otras malezas de la lista **según el listado que figura en esta etiqueta.
DÓNDE USAR	**Sobre el césped** †(Cuando se use según las indicaciones de uso. No use sobre césped Carpeta, Dichondra o variedad Floratam del pasto San Agustín.)
CUÁNDO USAR	Cuando las malezas están en crecimiento activo.
CUÁNTO USAR	**Céspedes del norte:** Cañuela, pasto azul de Kentucky y ballicos perennes **y** **Céspedes resistentes del sur:** Césped Bermuda, Búfalo y Zoysia 2 onzas líquidas (60 cc) cada 500 pies cuadrados (45 m2) **Céspedes del sur:** Césped Bahía, Centipede y San Agustín 1 onza líquida (30 cc) cada 500 pies cuadrados (45 m2)
	Para hacer preguntas o comentarios, llamar al número gratuito 1-877-BayerAG (1-877-229-3724) o visítenos en www.BayerAdvanced.com

SÓLO PARA USO RESIDENCIAL EN EXTERIORES

Ingredientes Activos:

2,4-D, sal dimetilamina	7.59%
Mecoprop-p, sal dimetilamina	1.83%
Dicamba, sal dimetilamina	0.84%
OTROS INGREDIENTES	89.74%
Total	100.00%

ESTE PRODUCTO CONTIENE:
0,00427 lb de equivalente ácido de 2,4-Diclorofenoxiacético por fl.oz. o 6,30%
0,00102 lb de equivalente ácido de Mecoprop-p por fl. oz. o 1,51%
0,00047 lb de equivalente ácido de Dicamba por fl. oz. o 0,69%
Isómero específico según los métodos AOAC.

INDICACIONES DE USO

Es una violación de la Ley Federal utilizar este producto de manera contraria a la indicada en la etiqueta. Lea y siga detenidamente las instrucciones de la etiqueta. No aplicar este producto de manera que entre en contacto con personas o mascotas, directamente o a través del escurrimiento. Mantener a las personas y a las mascotas alejadas del área durante la aplicación. No usar sobre césped que se cultive para la venta o uso comercial, como los panes de césped, ni para la producción de semillas comerciales, ni para fines de investigación. No aplicar sobre ningún tipo de sistema de irrigación.

ANTES DE USAR
Leer y seguir estas indicaciones de uso

- No aplicar directamente o cerca del agua, bocas de tormenta, canales de desagüe, alcantarillas o zanjas. No aplicar dentro de los 25 pies de distancia de ríos, estanques, lagos, arroyos, represas, pantanos, estuarios, bahías y océanos. No aplicar cuando hay viento. Para evitar que el producto se escurra, no regar en exceso el área o áreas tratadas hasta el punto de escurrimiento ni aplicar cuando llueve o cuando se esperan lluvias para ese día. Enjuagar el aplicador solamente sobre el área del césped o jardín.
- Rociar cuando se espera que la mayor parte del día tenga una temperatura inferior a 90 ° F (32 ° C).
- No usar este producto sobre plantas favoritas ni en sus cercanías ya que pueden ser dañadas, incluso si se encuentran dentro de la línea de goteo de árboles y arbustos.
- La cantidad máxima de aplicaciones al voleo por sitio de tratamiento es de dos por año.

(continuación)

6.375”

3.5” 3.5”

6/28/11

CCL LABEL YOUR GLOBAL PARTNER	**Description:** BOTPE 40oz. ASB_1_REV2- BOOK IN		**NOTES**	**LEGEND**
	Supplied for: Bayer	**File No.:** CCL4951	Total number of pages must be divisible by four (4).	Copy Image Area Cut/Fold Lines
	Finished Size: 8.125” (H) x 4.0” (W)	**Supersedes:** CCL4931		
	Date: 5/6/09	**Supplied by:** CCL Label Memphis	Measurements are in Inches	
	Application: Illustrator CS2	**Prepared By:** NR		

INDICACIONES DE USO *(continuación)*

CÓMO USAR
Agitar bien antes de usar

- Determinar la cantidad de producto a usar en base al tipo de césped y al tamaño del área a tratar.
- Preparar el producto para utilizar de acuerdo con las indicaciones de "Cómo aplicar".
- Aplicar al césped.

MATA

Trébol amarillo	Bardana menor	Mostaza	Cerraja brava
Aster	Barbasco	Plántago de hoja delgada	Pasto del chancho
Rorippa austriaca	Pilea nummulariifolia	Veza común	Euphorbia maculata
Galium	Lysimachia nummularia	Ortiga	Soliva sessilis
Aceitilla	Vira Vira	Naranja de la vellosilla	Ortiga mayor
Stachys floridana	Romaza	Bardana menor	Fresa de la India
Manto blanco	Margarita menor	Oxalis	Urtica procera
Vicia cracca	Erigeron philadelphicus	Alquimila arvense	Verbena lila
Berro de altura	Margarita mayor	Chirivía	Pamita
Falso berro	Diente de león	Sagina procumbens	Polígono anfibio
Helenium tenuifolium	Dichondra	Carraspique	Cardo
Alma del poeta	Verdolaga de la montaña	Obolaria virginica	Coronilla varia
Lupilina	Manzanilla cimarrona	Lentejilla	Nabo chileno
Mostaza negra	Sombrerito de agua	Bledo	Quelite
Plántago rugelii	Baya del saúco	Pineywoods bedstraw	Malva blanca
Cardo santo	Falso diente de león	Coreopsos tinctoria (pilosa de Bidens)	Hibisco
Hierba de la cucaracha	Camelina	Plántago	Diodia virginiana
Lechuga azul	Falso girasol	Hiedra venenosa	Parra virgen
Verbena azul	Amsinckia menziesii var	Roble venenoso	Lentejilla
Plántago aristata	Botoncillo	Hierba carmín	Centella asiática
Cotula	Carraspique	Diosa teres	Wavyleaf bullthistle
Cardo perruno	Galinsoga	Girasol silvestre	Clemátide
Lengua de vaca	Abrojos	Lactuca serviola	Salsifi
Plántago mayor	Vara de oro	Afata hembra	Mostaza blanca
Gutierrezia dracunculoides	Hiedra de tierra	Centinodia	Aster silvestre
Tritogonia verrucosa	Pegapega	Bledo rojo	Enredadera (Polygonum convolvulus)
Llantén menor	Rama negra	Hierba de golondrina	Zanahoria silvestre
Hierba velluda	Flor de peña	Verbena bracteata	Maravilla
Cardo negro	Cola de ratón	Abrojos	Ajo silvestre
Mala mujer	Cinquillo	Verdolaga	Geranio silvestre
Trébol de carretilla	Aster ericoides	Ambrosía	Lechuga silvestre
Bardana	Campanilla blanca	Rumex acetosella	Caléndula silvestre
Ortiga mayor	Erísimo	Yuyo colorado	Mostaza silvestre
Cadillo	Cáñamo común	Alfilerillo	Cebolla silvestre
Celidonia menor	Ortiga mansa	Potentilla norvegica	Chirivía silvestre
Hierba del soldado	Antonico	Erigeron asperugineus	Rábano silvestre
Calcida vera	Llantén media	Axyris amaranthoides	Colza silvestre
Geranio	Verbena stricta	Cardo ruso	Fresa silvestre
Verdolaga alfombra	Solanum de Carolina	Pimpinela escarlata	Batata silvestre
Galium aparine	Belladona	Cardo escocés	Vicia americana
Hierba gatera	Escleranto	Amargosa	Oxalis montana
Hierba del halcón	Sanguinaria	Bolsa del pastor	Croton capitatus
Capiqui común	Morenita	Plántago heterophylla	Palo de rosa
Cerastium vulgatum	Cenizos	Galinsoga parviflora	Hierba del pastor
Endivia	Lespedeza	Rumex altissimus	Chenopodium aristatum
Tormetilla	Malva	Amaranthus hybridus	Milenrama
Trébol encarnado	Escobilla	Acedera	Hierba de Santa Bárbara
Trifolium agrarium	Caperonia castaneifolia	Achicoria	Mastuerzo
Trébol rojo	Astrágalo	Bidens bipinnata	y otras malezas de hoja ancha
Trébol fresa	Campanilla	Verónica	
Trébol dulce	Oreja de ratón	Bledo espinoso	
Trébol blanco	Artemisa	Cepa caballo	
Berbecho	Cardo pendiente		

CUÁNTO APLICAR

Céspedes del norte: Cañuela, pasto azul de Kentucky y ballico perenne: 2 onzas líquidas (60 cc) (4 cucharadas) por cada galón de agua (4 litros)

Céspedes resistentes del sur: Césped Bermuda, Búfalo y Zoysia: 2 onzas líquidas (60 cc) (4 cucharadas) por cada galón de agua (4 litros)

Céspedes del sur: Césped Bahía, Centipede y San Agustín: 1 onza líquida (30 cc) (2 cucharadas) por cada galón de agua (4 litros),
Para los céspedes con tipos de hierba combinados, utilizar la tasa más baja.
Cada galón (4 litros) de solución trata hasta 500 pies cuadrados (45 m^2)

CONSEJO: Medir la longitud y ancho del área a tratar en pies (metros). Multiplicar estos dos números. El resultado es el área a tratar en pies cuadrados (m^2).

Restricciones de aplicación:

- Limitado a **2 aplicaciones por año** con un **intervalo de 30 días** entre aplicación.
- **Proporción máxima de aplicación simple** de 8,0 onzas fluidas/1.000 pies cuadrados (1,47 lbs de equivalente ácido de 2,4-D, 0,35 lb de equivalente ácido de MCPP-p y 0,16 lb de equivalente ácido de dicamba por acre con cada aplicación).

(continuación)

INDICACIONES DE USO *(continuación)*

CUÁNTO APLICAR *(continuación)*

- **Porcentaje máximo de temporada:** de 16,0 onzas fluidas./1.000 pies cuadrados (2,94 lbs de equivalente ácido de 2,4-D, 0,71 lb de equivalente ácido de MCPP-p y 0,33 lb de equivalente ácido de dicamba por acre con cada aplicación).

CÓMO APLICAR

ROCIADORES DE TANQUE ESTILO BOMBA

1. Agregar las cantidades adecuadas de agua y concentrado en el tanque del rociador.
2. Cerrar el rociador, agitar bien y presurizar.
3. Ajustar la boquilla en modo de rocío grueso.
4. Grandes áreas: aplicar uniformemente sobre el área.
   Malezas individuales: sostener la punta de la varilla a aproximadamente 12 pulgadas (30 cm) de la maleza, apuntar al centro de la maleza y rociar.
5. De vez en cuando, volver a presurizar el rociador según sea necesario para mantener un buen patrón de rociado.

ROCIADORES DE DISCO CON MANGUERA

1. Llenar el envase del rociador con concentrado suficiente para cubrir el área a tratar.
2. Cerrar el rociador y fijar el disco. DATO: los números en los rociadores de disco con manguera corresponden a las onzas de concentrado por galón de rocío.
3. Colocar el rociador en la manguera y aplicar.
4. Para evitar el contacto con el área tratada, comenzar a rociar el área más alejada del grifo y caminar hacia atrás a medida que aplica el producto.

ROCIADOR CON MANGUERA PARA TRATAR CON CONCENTRADO Y AGUA

1. Verter el concentrado en el envase del rociador hasta el nivel de onzas líquidas necesarias.
2. Agregar la cantidad correspondiente de agua hasta el nivel de galones.
3. Volver a colocar la tapa del rociador y agitar bien antes de aplicar.
4. Para evitar el contacto con el área tratada, comenzar a rociar el área más alejada del grifo y caminar hacia atrás a medida que aplica el producto.

CÓMO FUNCIONA

Este producto es un herbicida selectivo – elimina las malezas del césped en *forma selectiva* sin dañar el césped. Es una mezcla de herbicidas probados diseñados para las malezas de hoja ancha como el diente de león, trébol y capiqui.

PARA OBTENER MEJORES RESULTADOS

• Aplicar cuando las malezas son jóvenes y están en crecimiento activo. • Si el césped está seco, regar 24 horas antes de usar este producto. • El agua de lluvia o riego uno hora después de la aplicación no afectará la efectividad del producto. • Aplicar de 1 a 3 días después de segar. • Es posible que determinadas malezas difíciles de eliminar deben volver a tratarse en 30 días de forma localizada. • No aplicar sobre el césped nuevo hasta que haya alcanzado una altura mínima de 2 pulgadas (5 cm). • Los puntos descubiertos pueden sembrarse de 3 a 4 semanas después de la aplicación.

No permitir a las personas (excepto al aplicador) ni a las mascotas ingresar al área tratada durante la aplicación.
No permitir que personas o mascotas vuelvan a ingresar al área tratada hasta que el aerosol no se haya secado

IMPORTANTE – Indicaciones para almacenamiento y eliminación.

No contaminar alimentos ni forraje al almacenar o desechar el producto.

ALMACENAMIENTO DEL PESTICIDA

Almacenar en el contenedor original y en un lugar fresco y seco fuera del alcance de los niños, preferentemente en un armario bajo llave. Proteger del congelamiento

DESECHO DEL PESTICIDA Y MANIPULACIÓN DEL CONTENEDOR

Si está vacío:

- Contenedor no reusable. No volver a llenar ni utilizar este contenedor.
- Colocar el contenedor en la basura o entregarlo para reciclar si fuera posible.

Si está parcialmente lleno:

- Si es necesario eliminar el producto sin utilizar, llamar a la agencia local de residuos sólidos o al número gratuito 1-877-229-3724 para solicitar indicaciones para la eliminación.
- Nunca colocar el producto sin utilizar debajo de algún desagüe interno o externo.

6.375”

3.5” 3.5”

6/28/11

CCL LABEL YOUR GLOBAL PARTNER	Description: BOTPE 40oz. ASB_1_REV2- BOOK IN		NOTES	LEGEND
	Supplied for: Bayer	File No.: CCL4951	Total number of pages must be divisible by four (4).	Copy Image Area
	Finished Size: 8.125” (H) x 4.0” (W)	Supersedes: CCL4931		Cut/Fold Lines
	Date: 5/6/09	Supplied by: CCL Label Memphis	Measurements are in Inches	
	Application: Illustrator CS2	Prepared By: NR		