Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2015-2017 LIC. NO.

8126.127

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2012-2014 LIC. NO.

PREVENTS WEEDS up to 6 MONTHS

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2009-2011 LIC. NO.

3.75"

7.03"

7.28"

BAYER ADVANCED

Southern

SEASON LONG WEED CONTROL for LAWNS

CONCENTRATE

Kills Current Weeds

Prevents New Weeds up to 6 Months

Mata las malezas actuales
Previene las malezas nuevas hasta por 6 meses

¿HABLA ESPAÑOL? VER PANEL FRONTAL

- *Use on Bermudagrass, Centipede, St. Augustine & Zoysia Lawns*
- *Won't Harm Lawns†*

KEEP OUT OF REACH OF CHILDREN
CAUTION See back panel for additional precautionary statements

Active Ingredient:

2,4-D, dimethylamine salt	4.73%
Isoxaben	2.63%
Mecoprop-p potassium salt	1.10%
Dicamba, potassium salt	0.52%
Other Ingredients	91.02%
Total	100.00%

Treats up to 10,000 Sq Ft

NET CONTENTS 32 FL OZ (946ml)

00000000 R.0

4.0"

CCL LABEL YOUR GLOBAL PARTNER	Description: BOTPE 40oz. FRONT LABEL-_2_REV - TEMPLATE		NOTES	LEGEND
	Supplied for: Bayer	File No.: CCL4932		Copy Image Area
	Finished Size: 7.28" (H) x 4.0" (W)	Supersedes: CCL4916		Cut/Fold Lines
	Date: 5/5/09	Supplied by: CCL Label Memphis		
	Application: Illustrator CS2	Prepared By: NR	Measurements are in Inches	

7/01/09

BAYER ADVANCED

Southern

SEASON LONG WEED CONTROL for LAWNS CONCENTRATE

The days of multiple applications to kill weeds are over! Just one application of this product's proprietary 2-in-1 formula kills & prevents lawn weeds up to 6 months! It delivers Season Long Weed Control that's simple and easy to use and it will not harm your lawn when used as directed. Rainproof in one hour. Control weeds in your lawn up to 6 months!

PRECAUTIONARY STATEMENTS

CAUTION Hazards to Humans and Domestic Animals

- Harmful if absorbed through skin or if swallowed.
- Causes moderate eye irritation.
- Avoid contact with skin, eyes, or clothing.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove and wash contaminated clothing before reuse.

FIRST AID ✚

IF ON SKIN	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 – 20 minutes. • Call a poison control center or doctor for further treatment advice.
IF SWALLOWED	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF IN EYES	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for further treatment advice.

NOTE: Have the product container or label with you when calling a poison control center or doctor or going for treatment.
You may call toll-free 1-877-229-3763 for medical emergency information.
Active Ingredients: 4.73% 2,4-D, dimethylamine salt (CAS # 2008-39-1), 0.52% Dicamba, potassium salt (CAS # 10007-85-9), 1.10% Mecoprop-p, potassium salt (CAS # 66423-05-0) and 2.63% Isoxaben (CAS # 82558-50-7).

ENVIRONMENTAL HAZARDS

- This product is toxic to fish and aquatic invertebrates. Do not apply directly to water.
- Drift or runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

Produced for:
Bayer Advanced
A Business Unit of Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, NC 27709

Made in USA

EPA Reg. No. 72155-87
EPA Est. No indicated by 3rd and 4th digits of the batch number on this package.
(65) = 432-TX-1 (39) =58996-MO-1
(49) = 072155-AL-001 (13) = 071106-GA-001

Bayer (reg'd), Bayer Cross (reg'd), and Bayer Advanced™ are trademarks of Bayer.

PRESS TO RESEAL

000000R.0

7/06/09

READ THE LABEL FIRST!

Indicaciones en español después de las de inglés

OPEN
Resealable Label for Directions & Precautions

READ LABEL FIRST

QUICK FACTS

- **Protects Your Lawn from Weeds up to 6 Months**
- **Won't Harm Your Lawn†**
- **Rainproof in 1 Hour**

KILLS & PREVENTS	Kills over 200 broadleaf weeds such as Dollarweed, Chickweed, Dandelion, Henbit, Clover & Spurge Prevents New Weeds from Coming Back up to 6 Months
WHEN TO USE	Apply anytime - Spring or Fall.
WHERE TO USE	ON ESTABLISHED SOUTHERN LAWNS without harming lawn grasses †(Do not use on Carpetgrass, Floratam variety of St. Augustinegrass, Dichondra or desirable clovers.)
HOW TO USE	To prevent new weeds from growing, spray entire area you want protected. Wait 2 days after applying, then activate residual barrier by watering. See Directions for Use for further details.
COVERAGE	This bottle treats up to 10,000 sq.ft. (10,000 sq. ft. for Centipede & St. Augustine lawns; 5,000 sq. ft. for Bermudagrass and Zoysia lawns).
	For questions or comments, call toll-free 1-877-BayerAG (1-877-229-3724), or visit our website at www.BayerAdvanced.com

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt (CAS No. 2008-39-1)	4.73%
Isoxaben (CAS No. 82558-50-7)	2.63%
Mecoprop-p, potassium salt (CAS No. 66423-05-0)	1.10%
Dicamba, potassium salt (CAS No. 10007-85-9)	0.52%
OTHER INGREDIENTS	91.02%
TOTAL	100.00%

THIS PRODUCT CONTAINS:
0.337 lb 2,4-D acid equivalent per gal or 3.94%
0.225 lb Isoxaben per gal or 2.63%
0.080 lb Mecoprop-p acid equivalent per gal or 0.94%
0.038 lb Dicamba acid equivalent per gal or 0.44%

Isomer Specific by AOAC Method

EPA Reg. No. 72155-87

EPA Est. No indicated by 3rd and 4th digits of the batch number on this package.
(65) = 432-TX-1 (39) =58996-MO-1 00000000 R.0
(49) = 072155-AL-001 (13) = 071106-GA-001

FOR OUTDOOR RESIDENTIAL USE ONLY

6 FPO 5

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEFORE YOU USE	• Use this product only on established lawns. • Apply when the air is calm to avoid spray drift onto desirable plants.
HOW MUCH TO USE	• Measure area to be sprayed. Calculate 'square feet' by multiplying length times width. • For best results, use spray solution immediately. If it is necessary to store, shake vigorously before use. Do not store for more than 3 days. • **Bermudagrass, Buffalograss, Fescue, Kentucky Bluegrass, Perennial Ryegrass, Zoysiagrass 3.2 fl. oz. (6 1/2 TBS) in 2 gallons of water treats 500 sq. ft.** • **Bahiagrass, Centipedegrass, St. Augustinegrass (not for use on Floratam variety) or mixtures containing one or more of these grasses 1.6 fl. oz. (3 1/4 TBS) in 2 gallons of water treats 500 sq. ft.** • Food utensils, such as tablespoons and measuring cups, should not be used for food purposes after use with any pesticide. • The maximum application rate of this product is 6.4 fl. oz. per 1,000 sq. ft. per application per site. The maximum number of broadcast applications per treatment site is 2 per year.
HOW TO USE **Shake Bottle Well Before Use**	• **To prevent new weeds from growing, you must spray the entire area you want protected, not just the emerged weeds.** • **Wait 2 days after applying, then activate residual barrier by watering with 1/4 - 1/2" of irrigation.** **PUMP STYLE TANK SPRAYERS** *Best for spraying individual weeds & small areas.* 1. Add the appropriate amounts of water & product to sprayer tank. 2. Close sprayer, shake well and pressurize. Adjust nozzle to a coarse spray. **Individual weeds –** Hold tip of wand about 12" from weed, aim at center of weed & spray. Also spray the area underneath & around the weed to prevent new weeds from growing. **Small areas –** apply evenly over area. 3. Occasionally re-pressurize sprayer as needed to maintain a good spray pattern. **DIAL-STYLE HOSE-END SPRAYERS** *Best for spraying large areas such as the entire lawn.* 1. Fill sprayer jar with enough product for the area to be treated. DO NOT ADD WATER. 2. Close sprayer, set dial, hook to garden hose and apply. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray. 3. Pour any unused product back into this original container. **CONCENTRATE PLUS WATER HOSE-END SPRAYERS** *Best for spraying large areas such as the entire lawn.* 1. Pour product into sprayer jar to the required 'Fluid Ounce' level. 2. Add the appropriate amount of water to the 'Gallons' level. 3. Close sprayer, hook to garden hose, shake well & apply. To avoid contact with the treated area, begin by spraying area farthest from faucet walking backwards as you spray.

front

page 2

7/06/09

DIRECTIONS FOR USE *(cont'd)*

FOR BEST RESULTS

WHEN TO APPLY:
- Spray when
  - weeds are small and actively growing
  - temperatures are between 50 - 90°F
- Do not apply while grass is emerging from dormancy.
- Certain hard-to-kill weeds may be retreated in 2 weeks if necessary. Spray individual weeds using this product in a tank sprayer.
- Heavy raking will disturb the weed preventative barrier and reduce the effectiveness of this product.
- Newly seeded lawn may be sprayed after fourth mowing.
- Bare spots may be seeded 60 days after application.

MOWING:
- Do not mow for 1 to 2 days after spraying.
- Clippings from the first three mowings should be left on the treated area. Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

WATERING:
- If lawn is dry, water 1 to 2 days before using this product.
- Rain or watering 1 hour after application will not wash away effectiveness.

HOW IT WORKS

This product contains a proven blend of three weed killers to kill existing weeds PLUS a residual herbicide to keep them from coming back for up to 6 months. Weeds begin to wilt & die within hours and are completely dead in 14 – 28 days. And, it won't harm your lawn when used as directed.

KILLS (a "1" indicates *Kills & Prevents*)

Annual Yellow Sweetclover
Aster
Austrian Fieldcress
Bedstraw
Beggarticks
Betony, Florida
Bindweed, Field
Bird Vetch
Bitter Wintercress
Bittercress, Hairy[1]
Bitterweed
Black-eyed Susan
Black Medic
Black Mustard
Blackseed Plantain
Blessed Thistle
Bloodflower Milkweed
Blue Lettuce
Blue Vervain
Bracted Plantain[1]
Brassbuttons
Bristly Oxtongue
Broadleaf Dock
Broadleaf Plantain
Broomweed
Buckhorn
Buckhorn Plantain
Bulbous Buttercup
Bull Thistle
Bullnettle
Burclover
Burdock
Burning Nettle
Burweed[1]
Buttercup
Buttonweed
Canada Thistle
Carolina Geranium
Carpetweed
Catchweed Bedstraw
Catnip
Catsear
Chickweed, Common[1]
Chickweed, Mouseear
Chicory
Cinquefoil
Clover[1]
Clover, Crimson[1]
Clover, Hop[1]
Clover, Red[1]
Clover, Strawberry[1]
Clover, Sweet[1]
Clover, White[1]
Cockle
Cocklebur
Common Mullein
Creeping Charlie
Creeping Jenny
Cudweed[1]
Curly Dock
Daisy, English
Daisy, Fleabane
Daisy, Oxeye
Dandelion[1]
Dichondra
Dogbane
Dogfennel
Dollarweed[1]
Elderberry
False Dandelion[1]
False Flax
False Sunflower
Fiddleneck[1]
Florida Pusley
Frenchweed
Galinsoga
Goathead
Goldenrod
Ground Ivy
Gumweed
Hairy Fleabane[1]
Hawkweed
Healall
Heartleaf Drymary
Heath Aster
Hedge Bindweed
Hedge Mustard
Hemp
Henbit[1]
Hoary Cress
Hoary Plantain
Hoary Vervain
Horsenettle
Jimsonweed
Knawel
Knotweed

KILLS (a "1" indicates *Kills & Prevents*) *(cont'd)*

Kochia
Lambsquarters[1]
Lespedeza
Mallow[1]
Matchweed
Mexicanweed
Milk Vetch
Morningglory
Mouseear Hawkweed
Mugwort
Musk Thistle
Mustard
Narrowleaf Plantain
Narrowleaf Vetch
Nettle
Orange Hawkweed
Oriental Cocklebur
Oxalis[1]
Parsley-piert[1]
Parsnip
Pearlwort
Pennycress
Pennywort[1]
Pepperweed
Pigweed[1]
Pineywoods Bedstraw
Plains Coreopsis (Tickseed)
Plantain[1]
Poison Ivy
Poison Oak
Pokeweed
Poorjoe
Prairie Sunflower
Prickly Lettuce[1]
Prickly Sida
Prostrate Knotweed[1]
Prostrate Pigweed
Prostrate Spurge
Prostrate Vervain
Puncturevine
Purslane, Common[1]
Ragweed[1]
Red Sorrel
Redroot Pigweed
Redstem Filaree[1]
Rough Cinquefoil
Rough Fleabane[1]
Russian Pigweed
Russian Thistle[1]
Scarlet Pimpernel
Scotch Thistle
Sheep Sorrel
Shepherdspurse[1]
Slender Plantain
Smallflower Galinsoga
Smooth Dock
Smooth Pigweed
Sorrel
Sowthistle[1]
Spanishneedles
Speedwell[1]
Spiny Amaranth
Spiny Cocklebur
Spiny Sowthistle
Spotted Catsear
Spotted Spurge[1]
Spurweed
Stinging Nettle
Strawberry, India Mock
Tall Nettle
Tall Vervain
Tansy Mustard
Tanweed
Thistle
Trailing Crownvetch
Tumble Mustard
Tumble Pigweed
Velvetleaf[1]
Venice Mallow
Virginia Buttonweed
Virginia Creeper
Virginia Pepperweed[1]
Water Pennywort
Wavyleaf Bullthistle
Western Clematis
Western Salsify
White Mustard
Wild Aster
Wild Buckwheat
Wild Carrot
Wild Four-o'clock
Wild Garlic
Wild Geranium
Wild Lettuce
Wild Marigold
Wild Mustard[1]
Wild Onion[1]
Wild Parsnip
Wild Radish[1]
Wild Rape
Wild Strawberry
Wild Sweet Potato
Wild Vetch
Woodsorrel[1]
Wooly Croton
Wooly Morningglory
Wooly Plantain
Wormseed
Yarrow
Yellow Rocket
Yellowflower Pepperweed
and other broadleaf weeds

[1]*Kills & Prevents*

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.

IMPORTANT- Directions for Storage and Disposal

STORAGE CONDITIONS

Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.

DISPOSAL

If Empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.
If Partly Filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.

7/06/09

¡LEER LA ETIQUETA ANTES DE UTILIZAR EL PRODUCTO!

DATOS RÁPIDOS

• Protege su césped de las malezas hasta por 6 meses
• No daña el césped† • A prueba de lluvia en 1 hora

MATA Y PREVIENE	Mata más de 200 malezas de hoja ancha como el sombrerito de agua, capiqui, diente de león, ortiga mansa, trébol y lechetrezna Previene el regreso de nuevas malezas hasta por 6 meses
CUÁNDO USAR	Aplicar en cualquier época: primavera u otoño
DÓNDE USAR	SOBRE CÉSPEDES DEL SUR ESTABLECIDOS sin dañar la hierba †(No utilizar sobre la grama colorada, la variedad 'Floratam' del césped San Agustín, dichondra o los tréboles que desee conservar).
CÓMO USAR	Para evitar que crezcan nuevas malezas, rociar toda el área que desea proteger. Esperar 2 días después de la aplicación; luego, regar para activar la barrera residual. Consultar las Indicaciones de uso para obtener más detalles.
COBERTURA	Esta botella trata hasta 10,000 pies cuadrados (900 m2) (10,000 pies cuadrados (900 m2) para hierba Centipede y césped San Agustín; 5,000 pies cuadrados (450 m2) para céspedes Bermuda y Zoysia).
☎	Para hacer preguntas o comentarios, llamar al número gratuito 1-877-BayerAG (1-877-229-3724) o visítenos en nuestro sitio Web www.BayerAdvanced.com

SÓLO PARA USO RESIDENCIAL EN EXTERIORES

INGREDIENTES ACTIVOS:

2,4-D, sal dimetilamina (CAS Nº 2008-39-1)	4.73%	ESTE PRODUCTO CONTIENE:
Isoxaben (CAS Nº 82558-50-7)	2.63%	0.337 libras (153 g) de 2,4-D equivalente ácido por galón o 3.94%
Mecoprop-p, sal de potasio (CAS Nº 66423-05-0)	1.10%	0.225 libras (102 g) de isoxaben por galón o 2.63%
Dicamba, sal de potasio (CAS Nº 10007-85-9)	0.52%	0.080 libras (36 g) de mecoprop-p equivalente ácido por galón o 0.94%
OTROS INGREDIENTES	91.02%	0.038 libras (15 g) de dicamba equivalente ácido por galón o 0.44%
TOTAL	100.00%	Isómero específico por Métodos AOAC

INDICACIONES DE USO

Es una violación de la Ley Federal utilizar este producto de manera contraria a la indicada en la etiqueta.

ANTES DE USAR	• Aplicar este producto sólo sobre césped establecido. • Aplicar cuando el viento esté calmo para evitar que el rocío se esparza sobre las plantas que desee conservar.
CUÁNTO USAR	• Medir el área a rociar. Para calcular los "pies cuadrados" (metros cuadrados), multiplicar el largo por el ancho. • Para obtener mejores resultados, utilizar la solución de rocío de inmediato. Si es necesario almacenar, agitar vigorosamente antes de usar. No almacenar por más de 3 días. • **Césped Bermuda, hierba búfalo, cañuela, pasto azul de Kentucky; ballico perenne, césped Zoysia 3.2 onzas líquidas (100 cc) (6 1/2 cucharadas) en 2 galones (8 litros) de agua tratan 500 pies cuadrados (45 metros cuadrados).**

(continuación)

INDICACIONES DE USO *(continuación)*

CUÁNTO USAR *(continuación)*	• **Césped Bahía, hierba Centipede, césped San Agustín (no usar en la variedad Floratam) o mezclas que contengan uno o más de estos céspedes 1.6 onzas líquidas (50 cc) (3 1/4 cucharadas) en 2 galones (8 litros) de agua tratan 500 pies cuadrados (45 metros cuadrados).** • Los utensilios de cocina, como cucharas y tazas medidoras, no deben utilizarse para preparar alimentos luego de ser utilizados con insecticidas. • La tasa máxima de aplicación de este producto es de 6.4 onzas líquidas (190 cc) cada 1,000 pies cuadrados (90 metros cuadrados) de aplicación por sitio. La cantidad máxima de aplicaciones por esparcimiento por sitio de tratamiento es de 2 por año.
CÓMO USAR **Agitar bien la botella antes de usar**	• **Para evitar que crezcan nuevas malezas, debe rociar el área completa que desea proteger, no sólo las malezas que ya surgieron.** • **Esperar 2 días después de aplicar; luego, rociar con 1/4 ó 1/2" pulgada (0.6 cm ó 1.3 cm) de irrigación para activar la barrera residual.** **ROCIADORES DE TANQUE ESTILO BOMBA** *Los más adecuados para rociar malezas individuales y pequeñas áreas.* 1. Agregar las cantidades adecuadas de agua y producto en el tanque del rociador. 2. Cerrar el rociador, agitar bien y presurizar. Ajustar la boquilla en modo de rocío grueso. **Malezas individuales:** sostener la punta de la varilla a aproximadamente 12 pulgadas (30 cm) de la maleza, apuntar al centro de la maleza y rociar. Además, rociar el área debajo y alrededor de la maleza para evitar que crezcan nuevas malezas. **Pequeñas áreas:** aplicar uniformemente sobre el área. 3. De vez en cuando, volver a presurizar el rociador según sea necesario para mantener un buen patrón de rociado. **ROCIADORES DE DISCO CON MANGUERA** *Los más adecuados para rociar grandes áreas tales como el césped completo.* 1. Llenar el envase del rociador con suficiente producto para el área que se va a tratar. NO AGREGAR AGUA. 2. Cerrar el rociador y fijar el disco, colocar el rociador en la manguera y aplicar. Para evitar el contacto con el área tratada, comenzar a rociar el área más alejada del grifo y caminar hacia atrás a medida que aplica el producto. 3. Verter el producto sin usar nuevamente en este contenedor original. **ROCIADORES CON MANGUERA PARA TRATAR CON CONCENTRADO Y AGUA** *Los más adecuados para grandes áreas tales como el césped completo.* 1. Verter el producto en el envase del rociador hasta el nivel de "Onzas líquidas" necesarias. 2. Agregar la cantidad correspondiente de agua hasta el nivel de "Galones". 3. Cerrar el rociador, conectar a la manguera del jardín, agitar bien antes de aplicar. Para evitar el contacto con el área tratada, comenzar a rociar el área más alejada del grifo y caminar hacia atrás a medida que aplica el producto.
PARA OBTENER MEJORES RESULTADOS	**CUÁNDO APLICAR:** • Rociar cuando • las malezas son pequeñas y están en crecimiento activo • las temperaturas oscilan entre 50 y 90° F (10° C y 32° C) • No aplicar mientras el césped retorna del letargo. • Es posible que determinadas malezas difíciles de eliminar deban volver a tratarse en 2 semanas, si fuera necesario. Rociar las malezas individuales utilizando este producto en un rociador de tanque. • Rastrillar excesivamente alterará la barrera preventiva de malezas y reducirá la eficacia de este producto. • El césped recién sembrado puede rociarse después de segarlo por cuarta vez. • Los puntos descubiertos pueden sembrarse 60 días después de la aplicación. **SEGADO:** • No segar por 1 ó 2 días después de rociar. • Los recortes de césped de los primeros tres segados deben dejarse en el área tratada. No utilizar los recortes de las áreas tratadas como mantillo o abono orgánico alrededor de flores, plantas ornamentales, árboles o en jardines de hortalizas. **RIEGO:** • Si el césped está seco, regar 1 ó 2 días después de utilizar este producto. • La lluvia o riego una hora después de la aplicación no afectará la eficacia del producto.

(continuación)

7/06/09

INDICACIONES DE USO *(continuación)*

CÓMO FUNCIONA	Este producto contiene una mezcla probada de tres herbicidas para matar las malezas existentes ADEMÁS de un herbicida residual para evitar que regresen hasta por 6 meses. Las malezas comienzan a marchitarse y mueren en horas y están completamente muertas en 14 a 28 días. Y no daña su césped cuando es utilizado de acuerdo con las indicaciones.

MATA (un "1" significa *Mata y previene*)

Trébol amarillo
Aster
Rorippa austriaca
Galium
Aceitilla
Stachys floridana
Manto blanco
Vicia cracca
Berro de altura
Falso berro[1]
Helenium tenuifolium
Alma del poeta
Lupilina
Mostaza negra
Plántago rugelii
Cardo santo
Hierba de la cucaracha
Lechuga azul
Verbena azul
Plántago aristata[1]
Cotula
Cardo perruno
Lengua de vaca
Plántago mayor
Gutierrezia dracunculoides
Tritogonia verrucosa
Llantén menor
Hierba velluda
Cardo negro
Mala mujer
Trébol de carretilla
Bardana
Ortiga mayor
Cadillo[1]
Celidonia menor
Hierba del soldado
Calcida vera
Geranio
Verdolaga alfombra
Galium aparine
Hierba gatera
Hierba del halcón
Capiqui común[1]
Cerastium vulgatum
Endivia
Tormetilla
Trébol[1]
Trébol encarnado[1]
Trifolium agrarium[1]
Trébol rojo[1]
Trébol fresa[1]
Trébol dulce[1]
Trébol blanco[1]
Berbecho
Bardana menor
Barbasco
Pilea nummulariifolia
Lysimachia nummularia
Vira Vira[1]
Romaza
Margarita menor
Erigeron philadelphicus
Margarita mayor
Diente de león[1]
Dichondra
Verdolaga de la montana
Manzanilla cimarrona
Sombrerito de agua[1]
Baya del saúco
Falso diente de león[1]
Camelina
Falso girasol
Amsinckia menziesii var[1]
Botoncillo
Carraspique
Galinsoga
Abrojos
Vara de oro
Hiedra de tierra
Pegapega
Rama negra[1]
Flor de peña
Cola de ratón
Cinquillo
Aster ericoides
Campanilla Blanca
Erísimo
Cáñamo común
Ortiga mansa[1]
Antoñico
Llantén media
Verbena stricta
Solanum de Carolina
Belladona
Escleranto
Sanguinaria
Morenita
Cenizos[1]
Lespedeza
Malva[1]
Escobilla
Caperonia castaneifolia
Astrágalo
Campanilla
Oreja de ratón
Artemisa
Cardo pendiente
Mostaza
Plántago de hoja delgada
Veza común
Ortiga
Naranja de la vellosilla
Bardana menor
Oxalis[1]
Alquimila arvense[1]
Chirivía
Sagina procumbens
Carraspique
Obolaria virginica[1]
Lentejilla
Bledo[1]
Pineywoods Bedstraw
Coreopsos tinctoria
(pilosa de Bidens)
Plántago[1]
Hiedra venenosa
Roble venenoso
Hierba carmín
Diodia teres
Girasol silvestre
Lactuca serviola[1]
Afata hembra
Centinodia[1]
Bledo rojo
Hierba de golondrina
Verbena bracteata
Abrojos
Verdolaga[1]
Ambrosía[1]
Rumex acetosella
Yuyo colorado
Alfilerillo[1]
Potentilla norvegica
Erigeron asperugineus[1]
Axyris amaranthoides
Cardo ruso[1]
Pimpinela escarlata
Cardo escocés
Amargosa
Bolsa del pastor[1]
Plantago heterophylla
Galinsoga parviflora
Rumex altissimus
Amaranthus hybridus
Acedera
Achicoria[1]
Bidens bipinnata
Verónica[1]
Bledo espinoso
Cepa caballo
Cerraja brava
Pasto del chancho
Euphorbia maculata[1]
Soliva sessilis
Ortiga mayor
Fresa de la India
Urtica procera
Verbena lila
Pamita
Polígono anfibio
Cardo
Coronilla varia
Nabo chileno
Quelite
Malva blanca[1]
Hibisco
Diodia virginiana
Parra virgen
Lentejilla[1]
Centella asiática
Wayleaf Bulthistle
Clemátide
Salsafi
Mostaza blanca
Aster silvestre
Enredadera
(Polygonum Convolvulus)
Zanahoria silvestre
Maravilla
Ajo silvestre
Geranio silvestre
Lechuga silvestre
Caléndula silvestre
Mostaza silvestre[1]
Cebolla silvestre[1]
Chirivía silvestre
Rábano silvestre[1]
Colza silvestre
Fresa silvestre
Batata silvestre
Vicia americana
Oxalis montana[1]
Croton capitatus
Palo de rosa
Hierba del pastor
Chenopodium aristatum
Milenrama
Hierba de Santa Bárbara
Mastuerzo
y otras malezas de hoja ancha

[1]Mata y previene

No aplicar este producto de manera que entre en contacto con personas o mascotas, directamente o a través del escurrimiento. Mantener a las personas y a las mascotas alejadas del área durante la aplicación. No permitir a las personas ni a las mascotas ingresar al área tratada hasta que el rocío se haya secado.

IMPORTANTE: Indicaciones para almacenamiento y eliminación

CONDICIONES DE ALMACENAMIENTO	Almacenar en el contenedor original y en un lugar fresco y seco fuera del alcance de los niños, preferentemente en un armario bajo llave. Proteger del congelamiento
ELIMINACIÓN	**Si está vacío:** Contenedor no rellenable. No volver a utilizar ni rellenar este contenedor. Colocarlo en la basura o entregarlo para reciclar si fuera posible. **Si está parcialmente lleno:** Llamar a la agencia local de residuos sólidos o al número gratuito 1-877-229-3724 para solicitar indicaciones para la eliminación. Nunca colocar el producto sin utilizar debajo de algún desagüe interno o externo.

MANTENER FUERA DEL ALCANCE DE LOS NIÑOS

DECLARACIONES DE PRECAUCIÓN

PRECAUCIÓN — **Peligros para seres humanos y animales domésticos**

- Es perjudicial si se absorbe por la piel o se ingiere.
- Causa irritación moderada en los ojos.
- Evitar el contacto con la piel, los ojos o la ropa.
- Lavarse prolijamente con agua y jabón después de manipular este producto y antes de comer, beber, masticar goma de mascar, fumar o ir al baño.
- Quitarse y lavar la ropa contaminada antes de volver a usar.

PRIMEROS AUXILIOS ✚

EN CASO DE CONTACTO CON LA PIEL	• Quitar la ropa contaminada. • Enjuagar la piel de inmediato con abundante agua durante 15 a 20 minutos. • Llamar a un médico o a un centro de control de envenenamiento para mayores consultas sobre el tratamiento.
EN CASO DE INGESTIÓN	• Llamar inmediatamente a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento. • Hacer que la persona beba tragos de agua, si puede tragar. • No inducir el vómito a menos que lo indique un centro de control de envenenamiento o el médico. • No administrar nada por boca a una persona inconsciente.
EN CASO DE CONTACTO CON LOS OJOS	• Mantener los ojos abiertos y enjuagar en forma lenta y suave con agua durante 15 a 20 minutos. • Retirar los lentes de contacto, si los hubiera, después de los primeros 5 minutos, y luego continuar enjuagando los ojos. • Llamar a un médico o a un centro de control de envenenamiento para mayores consultas sobre el tratamiento.

NOTA: Tener el contenedor o la etiqueta del producto a mano al llamar a un médico, un centro de control de envenenamiento o al concurrir a un tratamiento.
Para solicitar información médica de emergencia, llamar al número gratuito 1-877-229-3763.
Ingredientes activos: 4.73% 2,4-D, sal dimetilamina [CAS Nº 2008-39-1], 0.52% dicamba, sal de potasio [CAS Nº 10007-85-9],1.10% mecoprop-p, sal de potasio [CAS Nº 66423-05-0] y 2.63% isoxaben [CAS Nº 82558-50-7].

PELIGROS AMBIENTALES

- Este producto es tóxico para los peces y los invertebrados acuáticos. No aplicar directamente al agua.
- El rocío y el escurrimiento pueden ser peligrosos para los organismos acuáticos presentes en las aguas adyacentes a las áreas tratadas. No contaminar el agua al eliminar el agua de lavado o enjuague del equipo.
- Para proteger el medio ambiente, no permitir que el pesticida ingrese o se escurra en las bocas de tormenta, zanjas, alcantarillas o aguas superficiales. Aplicar este producto cuando el clima esté calmo y no haya probabilidad de lluvia para las 24 horas siguientes ayudará a garantizar que el viento o la lluvia no dispersen o escurran el pesticida del área de tratamiento. Enjuagar el equipo de aplicación sobre el área tratada ayudará a evitar el escurrimiento a los cuerpos de agua o los sistemas de desagüe.

NOTA: Investigaciones y pruebas han determinado que las "Indicaciones de uso" son las adecuadas para el uso apropiado de este producto bajo las condiciones previstas. En la medida que sea acorde con la ley pertinente, el Comprador asume la responsabilidad por la falta de rendimiento o seguridad si no es utilizado de acuerdo con las indicaciones.

Garantía de devolución de dinero: Si usted no está satisfecho con este producto, gustosamente le reembolsaremos el monto original del precio de compra.

7/06/09