BAYER ADVANCED
NEW!
100% SATISFACTION GUARANTEED
3 IN 1
WEED & FEED
for SOUTHERN LAWNS
MATA Y PREVIENE LAS MALEZAS POR 6 MESES; Y FERTILIZA
1 KILLS
WEEDS INCLUDING DOLLARWEED, DANDELION & CLOVER
2 PREVENTS
BROADLEAF & GRASSY WEEDS PLUS CRABGRASS
3 FEEDS
GREENS & STRENGTHENS YOUR LAWN
UP TO 6 MONTHS
TREATS UP TO
5,000 Sq Ft
NET WT 12.5 Lbs (5.6 kg)
BEFORE
AFTER
Active Ingredients:
Dicamba acid 0.089%
Penoxsulam 0.041%
Indaziflam 0.027%
Other Ingredients 99.844%
Total 100.000%
KEEP OUT OF REACH OF CHILDREN
CAUTION See back panel for additional precautionary statements
35-0-3
FOR USE ON
SOUTHERN TURFS
BERMUDAGRASS, ST. AUGUSTINEGRASS, CENTIPEDEGRASS & ZOYSIA
FACE PANEL
CUSTOMER SEAL AREA
FOLD LINE MUST PRINT
LIMIT LINE MUST PRINT
Department of Agriculture STATE OF HAWAII
LICENSED
PERIOD 2015-2017 LIC. NO. 8126.162
LEFT GUSSET READS THIS WAY
OR READS THIS WAY
RIGHT GUSSET READS THIS WAY
OR READS THIS WAY
42" 1066mm
40.39" 1025.91mm
18" 457.2mm
1" 25.4mm
0.5" 12.7mm
0.25" 6.35mm
1.58" 40.13mm
16.42" 417.07mm
16.5" 419.1mm
0.08" 2.03mm
12.25" 311.15mm
12" 304.8mm
11.36" 288.54mm
7.5" 190.5mm
3.5" 88.9mm
0.32" 8.13mm
0.125" 3.17mm

5/29/15

BEFORE
AFTER
Active Ingredients:
Dicamba acid 0.088%
Penoxsulam 0.041%
Indaziflam 0.027%
Other Ingredients 99.844%
Total 100.000%
KEEP OUT OF REACH OF CHILDREN
CAUTION See back panel for additional precautionary statements
35-0-3
FOR USE ON
SOUTHERN TURFS
BERMUDAGRASS, ST. AUGUSTINEGRASS,
CENTIPEDEGRASS & ZOYSIA

FOLD LINE MUST PRINT

MICRODOT CENTERED IN REPEAT

CUSTOMER SEAL AREA

0.25" 6.35mm

12.7mm 0.5"

2.75" 69.85mm

41.14" 1044.96mm

18" 457.2mm

BAYER ADVANCED

3 IN 1 WEED & FEED for SOUTHERN LAWNS

quick FACTS

1. **KILLS** BROADLEAF WEEDS INCLUDING DOLLARWEED, DANDELION, CLOVER, CHICKWEED & HENBIT
2. **PREVENTS** GRASSY AND BROADLEAF WEEDS SUCH AS CRABGRASS, ANNUAL BLUEGRASS, COMMON DANDELION, GOOSEGRASS & SPURGE — SPRING/FALL UP TO 6 MONTHS
3. **FEEDS & STRENGTHENS** THE LAWN. CONTAINS SLOW RELEASE FERTILIZER FOR LONG LASTING FEEDING.

WHEN TO APPLY

Apply anytime in early spring or fall.

WHERE TO USE

Use only on established lawns of the turf types listed below:

Bermudagrass, Centipedegrass, all varieties of St. Augustinegrass including Floratam, and Zoysia

DO NOT use on Ormond variety of Bermudagrass, Fescue, Bentgrass, Carpetgrass, Kikuyugrass, Ryegrass or Dichondra lawns

COVERAGE This bag treats 5,000 sq. ft.

Questions or comments, call toll-free (877) 229-3724 www.BayerAdvanced.com

For the full Spanish translation go to www.BayerAdvanced.com

¿HABLA ESPAÑOL? VEA EL PANEL TRASERO

Tired of killing weeds in your lawn – even after you've put down your traditional weed & feed granules? Just one application of Bayer Advanced 3-in-1 Weed & Feed for Southern Lawns kills lawn weeds and prevents new ones, including Crabgrass, for up to 6 months. Plus it greens, feeds and strengthens your lawn for a thick, healthy turf. It delivers season long weed control that is simple and easy to use and it will not harm your lawn when used as directed. No need for frequent re-application. Plus, when applied in early spring, it prevents Crabgrass. And when re-applied late summer, prevents Annual Bluegrass. Up to 6 months of weed control – plus feeding – has never been easier.

****Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.***

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEFORE YOU USE

• Use this product only on established Southern lawns – new lawns must be at least 3 months old if sodded, 16 months old if sprigged & 16 months old if seeded. • Do not overlap treated areas or make repeat applications within 90 days as over application could damage the lawn. • Apply only to actively growing turf. • Do not apply to recently aerated or verticutted turf until it has recovered. • Apply when the air is calm to avoid drift onto desirable plants.

HOW TO USE

• Use rotary or drop type spreader to apply granules evenly over the lawn. To be sure you cover lawn areas uniformly, first treat the border of the lawn then fill in the center. Make parallel passes, walking at a steady pace to ensure even distribution of the granules. • Sweep, brush, or blow any accidentally applied material off driveways, sidewalks etc. back onto the lawn. • Wait 24 hours after applying then activate residual barrier by lightly watering in. • Do not make more than two applications per site in any 12 month period. • Heavy raking will disturb the weed preventative barrier and reduce the effectiveness of this product.

FOR BEST RESULTS

WHEN TO APPLY

• Apply when - weeds are small and actively growing - temperatures are between 50–90˚F. • May be applied to dry or (for best results) wet lawns. • Do not apply while turfgrass is emerging from dormancy. • Do not apply to Centipedegrass after October 1. • To prevent Crabgrass and Goosegrass apply early spring before the temperatures are regularly in the lower 80˚ F range. • To prevent Annual Bluegrass apply August - December. • Lawns may be seeded 12 months, sodded 6 months, and sprigged 12 months after applying this product.

MOWING

• If the lawn needs mowing, do it 1 to 2 days before (for best results) or 1 day after applying this product. • Clippings from the first three mowings should be left on the treated area. Do not use the clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

WATERING

• If lawn is dry, water 1 to 2 days before using this product. • Do not water for at least 24 hours after application or apply if rain is expected within 24 hours.

HOW IT WORKS

This product contains a proven blend of weed killers to kill existing weeds plus a residual herbicide to keep them from coming back for up to 6 months. This product starts working immediately to stop weed growth until weeds are dead within weeks. Plus, it feeds your lawn for quick greening and long lasting feeding.

INDICACIONES DE USO

Es una violación de la Ley Federal utilizar este producto de manera contraria a la indicada en la etiqueta.

ANTES DE USAR

• Usar este producto solo en céspedes del sur establecidos; los céspedes nuevos deben tener al menos 3 meses si se los colocó en panes, 16 meses si se colocaron retoños y 16 meses si se plantaron de semilla. • No superponer las áreas tratadas ni repetir las aplicaciones dentro de los 90 días dado que la aplicación podría dañar el césped. • Aplicar solo en grama de crecimiento activo. • No aplicar a grama aireada recientemente o que se ha sometido a poda vertical hasta que se recupere. • Aplicar cuando el viento esté calmo para evitar salpicaduras a plantas deseables.

CÓMO USAR

• Usar un rociador giratorio o por goteo para aplicar los gránulos de manera uniforme sobre el césped. Para asegurarse de cubrir las áreas del césped de manera uniforme, tratar primero el borde del césped y luego rellenar el centro. Hacer pasadas paralelas, caminar a paso lento para garantizar una distribución uniforme de los gránulos. • Barrer, rastrillar o soplar hacia el césped cualquier material que se aplique accidentalmente en caminos, aceras, etc. • Esperar 24 horas después de aplicar y luego activar la barrera residual regando ligeramente. • No hacer más de dos aplicaciones por sitio durante cualquier período de 12 meses. • Rastrillar demasiado perturbará la barrera preventiva de la maleza y reducirá la eficacia del producto.

PARA OBTENER MEJORES RESULTADOS

CUÁNDO APLICAR

• Aplicar cuando - las malezas sean pequeñas y estén en activo crecimiento - haya una temperatura entre 50–90˚F. • Puede aplicarse a céspedes secos o (para obtener mejores resultados) húmedos. • No aplicar cuando la grama esté emergiendo del período de letargo. • No aplicar sobre centipede después de 1 de octubre. • Para prevenir el garranchuelo y la pata de gallina, aplicar en primavera antes de que las temperaturas estén por debajo del rango de los 80˚F. • Para prevenir el pasto azul anual, aplicar de agosto a diciembre. • Los céspedes pueden sembrarse, colocarse en panes o en retoños después de 12 meses, 6 meses o 12 meses, respectivamente, después de aplicar el producto.

SEGADO

• Si el césped necesita ser segado, hacerlo 1 o 2 días antes (para obtener mejores resultados) o 1 día después de aplicar este producto. • Los recortes de las tres primeras segadas deben dejarse en el área tratada. No usar los recortes de las áreas tratadas como mantillo o compost alrededor de flores, plantas ornamentales, árboles o huertas.

RIEGO

• Si el césped está seco, regar 1 a 2 días antes de usar este producto. • No regar al menos durante 24 horas después de la aplicación ni aplicar si se espera que llueva dentro de las 24 horas.

CÓMO FUNCIONA

Este producto contiene una probada mezcla de herbicidas que eliminan las malezas existentes junto con un herbicida residual que previene que vuelvan a crecer por 6 meses. Este producto comienza a trabajar inmediatamente para frenar el crecimiento de las malezas hasta exterminarlas en pocas semanas. Además, fertiliza el jardín para que las plantas crezcan rápido con alimento de largo alcance.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

No aplicar este producto de manera que pueda estar en contacto con seres humanos o mascotas, ya sea en forma directa o por la corriente. Mantener a las personas y a las mascotas alejadas del área durante la aplicación. No permitir que los seres humanos o las mascotas ingresen al área tratada hasta que el producto esté asentado.

SPREADER SETTINGS

If your spreader is not listed, please call our Consumer Service Hotline toll-free (877) 229-3724 for help.

Broadcast Spreader — Drop Spreader

• Settings are approximate. The condition of your spreader, the speed at which you walk and uneven terrain may affect the rate of application. Calibrate your spreader according to manufacturer's directions.
• Tip: 1,000 sq ft = 20 ft x 50 ft
• Do not use a spreader that applies granules in narrow, concentrated or uneven bands. Be sure the spreader works properly and spreads granules evenly.

SETTING TO DELIVER 2.5lb/1000 sq. ft.

	SETTING
Broadcast Spreader	
Earthway Ev-N-Spread	10
Scotts Basic	3
Scotts Deluxe Edgeguard	3
Scotts Mini	3
Scotts Standard	8
Sta-Green	3
Drop Spreader	
Scotts Accugreen 1000	2.5
Scoots Accugreen 3000	2

KILLS

Beggarweed, Creeping; Beggarweed, Dwarf; Betony, Florida; Burdock; Buttercup, Hairy; Buttonweed, Virginia*; Clover, Hop; Clover, Red; Cocklebur, Common; Coffeeweed, (Common Chickory); Daisy, English (Lawn Daisy); Dandelion, False; Dollarweed (Pennywort); Goldenrod; Hawkweed; Heallal; Ivy, Ground (Creeping Charlie); Knotweed; Kyllinga, Green*; Lespedeza, Common; Mallow, Roundleaf; Matchweed; Nutsedge, Purple*; Nutsedge, Yellow*; Parsley-piert; Peppergrass; Pepperweed, Virginia; Plantain, Broadleaf; Smartweed; Sorrel, Sheep; Speedwell, Common; Spotted Catsear; Spurweed; Thistle; Vetch, Common; Wild Violet; Woodsorrel, Creeping

* Suppression

PREVENTS

Ageratum, Tropic; Barley, Mouse; Barnyardgrass; Bittercress; Black Nightshade, Hairy; Bluegrass, Annual; Brome, Foxtail; Buckwheat, Wild; Burweed, Lawn; California Burclover; Chamomile, False; Cheatgrass; Crabgrass, Henry; Crabgrass, Large (Hairy Crabgrass); Crabgrass, Smooth; Cudweed, Linear Leaf (Purple Cudweed); Dandelion, Cat's Ear; Doveweed; Evening Primrose, Common; Evening Primrose, Cutleaf; Flatsedge, Rice; Fleabane, Hairy; Foxtail, Giant; Foxtail, Green; Foxtail, Yellow; Filaree, Redstem; Goosegrass; Groundsel, Common; Guineagrass; Kochia; Kyllinga, Fragrant; Lovegrass, Tufted; Mallow, Little; Marestail; Mulberry Weed; Mustard, Black; Mustard, Wild; Nightshade, American Black; Nightshade, Hairy; Parthenium; Phyllanthus, Longstalked; Pigweed, Prostrate; Pigweed, Redroot; Plantain, Paleseed; Puncturevine; Redmaids; Rocket, London; Ryegrass, Italian; Ryegrass, Perennial; Sandbur; Sedge, Annual; Sesbania, Hemp; Sida, Prickly (Teaweed); Southern Brassbuttons; Sowthistle, Annual; Speedwell, Corn; Spurge, Prostrate; Sunflower, Common; Swinecress; Tassel Flower; Velvetleaf; Vetch, Purple; Wild Carrot; Willowherb; Woodsorrel (Oxalis)

KILLS AND PREVENTS

Burnweed, American; Chickweed, Common; Chickweed, Mouseear; Clover, White; Dandelion, Common; Dock, Curly; Eclipta; Henbit; Knotweed, Prostrate; Kyllingua, Annual*; Lambsquarters; Medic, Black; Plantain, Buckhorn; Purslane, Common; Pusley, Florida; Ragweed, Common; Shepardspurse; Spurge, Spotted; Thistle, Canada; Woodsorrel, Yellow

* Suppression

ENVIRONMENTAL HAZARDS

• This product is toxic to fish, aquatic invertebrates, and plants. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean water mark. Do not contaminate water when disposing of rinsate or washwater. This product may impact water through spray drift or runoff. Follow directions for use to avoid spray drift and runoff. A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams and springs will reduce the potential of this product entering water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.
• ***Ground Water Advisory:*** This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.
• ***Surface Water Advisory:*** This pesticide may impact water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application.

PELIGROS AMBIENTALES

• Este producto es tóxico para los peces, los invertebrados acuáticos y las plantas. No aplicar directamente al agua o en áreas donde haya aguas superficiales o áreas entre mareas por debajo de la marca de agua. No contaminar el agua al eliminar el agua de lavado o enjuague. Este producto puede afectar el agua a través del rocío o derrames. Seguir las instrucciones de uso para evitar rocíos y derrames. Un buen mantenimiento de la zona de contención de vegetación entre las áreas en las que se aplica este producto y la diferenciación del agua de superficie, incluidos estanques, arroyos y manantiales, reduce el potencial de penetración en el agua a causa de precipitación-derrame. El derrame de este producto se reducirá al evitar las aplicaciones cuando se pronostica lluvia dentro de las 48 horas.
• ***Recomendación sobre las napas subterráneas:*** Este pesticida contiene propiedades y características relacionadas con sustancias químicas detectadas en aguas subterráneas. Este producto químico puede filtrarse en las napas subterráneas si se utiliza en áreas donde los suelos son permeables, particularmente donde el nivel freático es poco profundo.
• ***Recomendación sobre el agua de superficie:*** Este pesticida puede impactar sobre la calidad del agua a causa de derrame de agua pluvial. Esto se cumple especialmente en suelos con drenaje deficiente y suelos con napas subterráneas poco profundas. Este producto se clasifica como con alto potencial para llegar al agua de superficie a través de derrames durante varios meses o más después de la aplicación.

PRECAUTIONARY STATEMENTS

CAUTION HAZARDS TO HUMANS & DOMESTIC ANIMALS

• Harmful if swallowed or absorbed through skin. • Causes moderate eye irritation.
• Avoid contact with eyes, skin, or clothing. • Remove and wash contaminated clothing before reuse.
• When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

After using this product:
• Rinse gloves before removing and wash hands. • If product gets on skin, wash thoroughly with soap & water.
• If product gets on clothing, remove clothing as soon as possible and launder clothing separately before reuse.
• Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

DECLARACIONES DE PRECAUCIÓN

PRECAUCIÓN RIESGOS PARA SERES HUMANOS Y ANIMALES DOMÉSTICOS

• Es perjudicial si se ingiere o absorbe por la piel. • Causa irritación moderada en los ojos. • Evitar el contacto con los ojos, la piel o la ropa. • Quitarse la ropa contaminada y lavarla antes de volver a usar. • Cuando use este producto, use camiseta de mangas largas, pantalones largos, calcetines, calzado y guantes de goma.

Después de usar este producto:
• Enjuagar los guantes antes de quitárselos y lavarse las manos. • Si el producto entra en contacto con la piel, lavarse bien con agua y jabón. • Si el producto entra en contacto con la ropa, quitarse la ropa tan pronto como sea posible y lavarla por separado antes de volver a utilizar. • Lavarse bien con agua y jabón después de manipular y antes de comer, beber, consumir goma de mascar, tabaco o usar el baño

FIRST AID

IF SWALLOWED

• Call a poison control center or doctor immediately for treatment advice.
• Have person sip a glass of water if able to swallow.
• Do not induce vomiting unless told to do so by a poison control center or doctor.
• Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING

• Take off contaminated clothing.
• Rinse skin immediately with plenty of water for 15 – 20 minutes.
• Call a poison control center or doctor for further treatment advice.

IF IN EYES

• Hold eye open and rinse slowly and gently with water for 15 – 20 minutes.
• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
• Call a poison control center or doctor for further treatment advice.

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. **You may call toll-free 1-877-229-3763 for medical emergency information.**
Active Ingredients: 0.088% Dicamba, (CAS # 1918-00-9), 0.041% Penoxsulam (CAS # 219714-96-2) and 0.027% Indaziflam (CAS # 950782-86-2).

PRIMEROS AUXILIOS

EN CASO DE INGESTIÓN • Llamar inmediatamente a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento. • Hacer que la persona beba tragos de agua, si puede tragar. • No inducir el vómito salvo que así lo indique el médico o el centro de control de envenenamiento. • No administrar nada por vía oral a una persona inconsciente.
EN CASO DE CONTACTO CON LA PIEL O LA ROPA • Quitar la ropa contaminada. • Enjuagar la piel inmediatamente con abundante agua durante 15 a 20 minutos • Llamar a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento.
EN CASO DE CONTACTO CON LOS OJOS • Mantener abiertos los ojos y enjuagar de forma lenta y suave con agua durante entre 15 y 20 minutos. • Retirar los lentes de contacto, si los hubiera, después de los primeros 5 minutos y luego continuar enjuagando los ojos. • Llamar a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento.
NOTA: Tener la etiqueta del producto a mano al llamar a un centro de control de envenenamiento. Si busca tratamiento en una sala de emergencias o en el consultorio de un médico, lleve esta etiqueta del producto para mostrársela al personal médico. **También puede llamar al número gratuito 1-877-227-3763 para solicitar información médica de emergencia.**
Ingredientes activos: 0.088% de dicamba, (CAS n.°1918-00-9), 0.041% de penoxsulam (CAS n.°219714-96-2) y 0.027% de indaziflam (CAS n.°950782-86-2).

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage and disposal

PESTICIDE STORAGE Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.
PESTICIDE DISPOSAL & CONTAINER HANDLING **If Empty:** Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.
If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.

ALMACENAMIENTO Y ELIMINACIÓN

No contaminar el agua o los alimentos al almacenar o eliminar

ALMACENAMIENTO DE PESTICIDAS Almacenar en el contenedor original en un lugar fresco y seco, fuera del alcance de los niños, preferentemente en un gabinete bajo llave. Proteger del congelamiento.
ELIMINACIÓN DEL PESTICIDAS Y MANIPULACIÓN DE ENVASES
Si está vacío: Envase no rellenable. No rellenar ni volver a utilizar este envase. Colocar el envase vacío en la basura o entregarlo para reciclar si fuera posible.
Si está parcialmente lleno: Llamar a la agencia local de residuos sólidos o al número gratuito 1-877-229-3724 para obtener indicaciones para la eliminación Nunca colocar el producto sin utilizar debajo de algún desagüe interno o externo.

6 87073 04840 7

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

EPA Reg. No. 72155-112
EPA Est. No. indicated by 3rd and 4th digits of the batch number on this package.
(26) = 82757-FL-1 (57) = 071106-GA-003

Bayer Advanced 3-in-1 Weed & Feed for Southern Lawns 35-0-3

GUARANTEED ANALYSIS
Total Nitrogen (N) 35.0%
35% Urea Nitrogen*
Soluble Potash (K_2O) 3.0%
Sulfur (S) Total 4.6%
3.5% Free Sulfur (S)
1.1% Combined Sulfur (S) **F1514**

Derived From: Urea, Polymer Coated Sulfur Coated Urea, & Potassium Sulfate
*Contains 17.5% slowly available Nitrogen from Polymer Coated Sulfur Coated Urea

Information regarding the contents and levels of metals in this product are available on the Internet at http://www.aapfco.org/metals.htm.

Check with your county or city government to determine if there are local regulations for fertilizer use.

Produced for: Bayer Advanced, A Division of Bayer CropScience LP, P.O. Box 12014, 2 T.W. Alexander Drive, Research Triangle Park, NC 27709

Bayer (reg'd), the Bayer Cross (reg'd) and Bayer Advanced™ are trademarks of Bayer

TRIM GUIDES MUST PRINT

TRIM GUIDES MUST PRINT

0.39" 9.91mm

3.5" 88.9mm

BAYER ADVANCED

3 IN 1 WEED & FEED for SOUTHERN LAWNS

1 KILLS WEEDS INCLUDING DOLLARWEED, DANDELION & CLOVER

2 PREVENTS BROADLEAF & GRASSY WEEDS PLUS CRABGRASS

3 FEEDS GREENS & STRENGTHENS YOUR LAWN

UP TO 6 MONTHS

MICRODOT CENTERED IN REPEAT

5/29/15

BAYER ADVANCED

3 IN 1 WEED & FEED for SOUTHERN LAWNS

quick FACTS

1. **KILLS** BROADLEAF WEEDS INCLUDING DOLLARWEED, DANDELION, CLOVER, CHICKWEED & HENBIT
2. **PREVENTS** GRASSY AND BROADLEAF WEEDS SUCH AS CRABGRASS, ANNUAL BLUEGRASS, COMMON DANDELION, GOOSEGRASS & SPURGE

SPRING/ FALL UP TO 6 MONTHS

Tired of killing weeds in your lawn – even after you've put down your traditional weed & feed granules? Just one application of Bayer Advanced 3-in-1 Weed & Feed for Southern Lawns kills lawn weeds and prevents new ones, including Crabgrass, for up to 6 months. Plus it greens, feeds and strengthens your lawn for a thick, healthy turf. It delivers season long weed control that is simple and easy to use and it will not harm your lawn when used as directed. No need for frequent re-application. Plus, when applied in early spring, it prevents Crabgrass. And when re-applied late summer, prevents Annual Bluegrass. Up to 6 months of weed control – plus feeding – has never been easier.

****Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.***

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEFORE YOU USE

• Use this product only on established Southern lawns – new lawns must be at least 3 months old if sodded, 16 months old if sprigged & 16 months old if seeded. • Do not overlap treated areas or make repeat applications within 90 days as over application could damage the lawn. • Apply only to actively growing turf. • Do not apply to recently aerated or verticutted turf until it has recovered. • Apply when the air is calm to avoid drift onto desirable plants.

HOW TO USE

• Use rotary or drop type spreader to apply granules evenly over the lawn. To be sure you cover lawn areas uniformly, first treat the border of the lawn then fill in the center. Make parallel passes, walking at a steady pace to ensure even distribution of the granules. • Sweep, brush, or blow any accidentally applied material off driveways, sidewalks etc. back onto the lawn. • Wait 24 hours after applying then activate residual barrier by lightly watering in. • Do not make more than two applications per site in any 12 month period. • Heavy raking will disturb the weed preventative barrier and reduce the effectiveness of this product.

FOR BEST RESULTS

WHEN TO APPLY

• Apply when - weeds are small and actively growing - temperatures are between 50–90°F.

INDICACIONES DE USO

Es una violación de la Ley Federal utilizar este producto de manera contraria a la indicada en la etiqueta.

ANTES DE USAR

• Usar este producto solo en céspedes del sur establecidos; los céspedes nuevos deben tener al menos 3 meses si se los colocó en panes, 16 meses si se colocaron retoños y 16 meses si se plantaron de semilla. • No superponer las áreas tratadas ni repetir las aplicaciones dentro de los 90 días dado que la aplicación podría dañar el césped. • Aplicar solo en grama de crecimiento activo. • No aplicar a grama aireada recientemente o que se ha sometido a poda vertical hasta que se recupere. • Aplicar cuando el viento esté calmo para evitar salpicaduras a plantas deseables.

CÓMO USAR

• Usar un rociador giratorio o por goteo para aplicar los gránulos de manera uniforme sobre el césped. Para asegurarse de cubrir las áreas del césped de manera uniforme, tratar primero el borde del césped y luego rellenar el centro. Hacer pasadas paralelas, caminar a paso lento para garantizar una distribución uniforme de los gránulos. • Barrer, rastrillar o soplar hacia el césped cualquier material que se aplique accidentalmente en caminos, aceras, etc. • Esperar 24 horas después de aplicar y luego activar la barrera residual regando ligeramente. • No hacer más de dos aplicaciones por sitio durante cualquier período de 12 meses. • Rastrillar demasiado perturbará la barrera preventiva de la maleza y reducirá la eficacia del producto.

PARA OBTENER MEJORES RESULTADOS

CUÁNDO APLICAR

• Aplicar cuando - las malezas sean pequeñas y estén en activo crecimiento - haya una temperatura entre 50–90°F. • Puede aplicarse a céspedes secos o (para obtener mejores resultados) húmedos. • No aplicar

ANNUAL BLUEGRASS,
COMMON DANDELION, GOOSEGRASS & SPURGE

3 FEEDS & STRENGTHENS
THE LAWN. CONTAINS SLOW RELEASE FERTILIZER FOR LONG LASTING FEEDING.

WHEN TO APPLY

Apply anytime in early spring or fall.

WHERE TO USE

Use only on established lawns of the turf types listed below:

Bermudagrass, Centipedegrass, all varieties of St. Augustinegrass including Floratam, and Zoysia

***DO NOT** use on Ormond variety of Bermudagrass, Fescue, Bentgrass, Carpetgrass, Kikuyugrass, Ryegrass or Dichondra lawns*

COVERAGE *This bag treats 5,000 sq. ft.*

Questions or comments, call toll-free (877) 229-3724 www.BayerAdvanced.com

For the full Spanish translation go to www.BayerAdvanced.com

¿HABLA ESPAÑOL? VER EL PANEL TRASERO

FOR BEST RESULTS

WHEN TO APPLY

• Apply when - weeds are small and actively growing - temperatures are between 50–90˚F. • May be applied to dry or (for best results) wet lawns. • Do not apply while turfgrass is emerging from dormancy. • Do not apply to Centipedegrass after October 1. • To prevent Crabgrass and Goosegrass apply early spring before the temperatures are regularly in the lower 80˚ F range. • To prevent Annual Bluegrass apply August - December. • Lawns may be seeded 12 months, sodded 6 months, and sprigged 12 months after applying this product.

MOWING

• If the lawn needs mowing, do it 1 to 2 days before (for best results) or 1 day after applying this product. • Clippings from the first three mowings should be left on the treated area. Do not use the clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

WATERING

• If lawn is dry, water 1 to 2 days before using this product. • Do not water for at least 24 hours after application or apply if rain is expected within 24 hours.

HOW IT WORKS

This product contains a proven blend of weed killers to kill existing weeds plus a residual herbicide to keep them from coming back for up to 6 months. This product starts working immediately to stop weed growth until weeds are dead within weeks. Plus, it feeds your lawn for quick greening and long lasting feeding.

PARA OBTENER MEJORES RESULTADOS

CUÁNDO APLICAR

• Aplicar cuando - las malezas sean pequeñas y estén en activo crecimiento - haya una temperatura entre 50–90˚F. • Puede aplicarse a céspedes secos o (para obtener mejores resultados) húmedos. • No aplicar cuando la grama esté emergiendo del período de letargo. • No aplicar sobre centipede después de 1 de octubre. • Para prevenir el garranchuelo y la pata de gallina, aplicar en primavera antes de que las temperaturas estén por debajo del rango de los 80˚F. • Para prevenir el pasto azul anual, aplicar de agosto a diciembre. • Los céspedes pueden sembrarse, colocarse en panes o en retoños después de 12 meses, 6 meses o 12 meses, respectivamente, después de aplicar el producto.

SEGADO

• Si el césped necesita ser segado, hacerlo 1 o 2 días antes (para obtener mejores resultados) o 1 día después de aplicar este producto. • Los recortes de las tres primeras segadas deben dejarse en el área tratada. No usar los recortes de las áreas tratadas como mantillo o compost alrededor de flores, plantas ornamentales, árboles o huertas.

RIEGO

• Si el césped está seco, regar 1 a 2 días antes de usar este producto. • No regar al menos durante 24 horas después de la aplicación ni aplicar si se espera que llueva dentro de las 24 horas.

CÓMO FUNCIONA

Este producto contiene una probada mezcla de herbicidas que eliminan las malezas existentes junto con un herbicida residual que previene que vuelvan a crecer por 6 meses. Este producto comienza a trabajar inmediatamente para frenar el crecimiento de las malezas hasta exterminarlas en pocas semanas. Además, fertiliza el jardín para que las plantas crezcan rápido con alimento de largo alcance.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

No aplicar este producto de manera que pueda estar en contacto con seres humanos o mascotas, ya sea en forma directa o por la corriente. Mantener a las personas y a las mascotas alejadas del área durante la aplicación. No permitir que los seres humanos o las mascotas ingresen al área tratada hasta que el producto esté asentado.

SPREADER SETTINGS

If your spreader is not listed, please call our Consumer Service Hotline toll-free (877) 229-3724 for help.



• Settings are approximate. The condition of your spreader, the speed at which you walk and uneven terrain may affect the rate of application. Calibrate your spreader according to manufacturer's directions.
• Tip: 1,000 sq ft = 20 ft x 50 ft
• Do not use a spreader that applies granules in narrow, concentrated or uneven bands. Be sure the spreader works properly and spreads granules evenly.

SETTING TO DELIVER 2.5lb/1000 sq. ft.

Broadcast Spreader	SETTING
Earthway Ev-N-Spread	10
Scotts Basic	3
Scotts Deluxe Edgeguard	3
Scotts Mini	3
Scotts Standard	8
Sta-Green	3
Drop Spreader	
Scotts Accugreen 1000	2.5
Scoots Accugreen 3000	2

KILLS

Beggarweed, Creeping
Beggarweed, Dwarf
Betony, Florida
Burdock
Buttercup, Hairy
Buttonweed, Virginia*
Clover, Hop
Clover, Red
Cocklebur, Common
Coffeeweed, (Common Chickory)
Daisy, English (Lawn Daisy)
Dandelion, False
Dollarweed (Pennywort)
Goldenrod
Hawkweed
Heallal
Ivy, Ground (Creeping Charlie)
Knotweed
Kyllinga, Green*
Lespedeza, Common
Mallow, Roundleaf
Matchweed
Nutsedge, Purple*
Nutsedge, Yellow*
Parsley-piert
Peppergrass
Pepperweed, Virginia
Plantain, Broadleaf
Smartweed
Sorrel, Sheep
Speedwell, Common
Spotted Catsear
Spurweed
Thistle
Vetch, Common
Wild Violet
Woodsorrel, Creeping

** Suppression*

PREVENTS

Ageratum, Tropic
Barley, Mouse
Barnyardgrass
Bittercress
Black Nightshade, Hairy
Bluegrass, Annual
Brome, Foxtail
Buckwheat, Wild
Burweed, Lawn
California Burclover
Chamomile, False
Cheatgrass
Crabgrass, Henry
Crabgrass, Large (Hairy Crabgrass)
Crabgrass, Smooth
Cudweed, Linear Leaf (Purple Cudweed)
Dandelion, Cat's Ear
Doveweed
Evening Primrose, Common
Evening Primrose, Cutleaf
Flatsedge, Rice
Fleabane, Hairy
Foxtail, Giant
Foxtail, Green
Foxtail, Yellow
Filaree, Redstem
Goosegrass
Groundsel, Common
Guineagrass
Kochia
Kyllinga, Fragrant
Lovegrass, Tufted
Mallow, Little
Marestail
Mulberry Weed
Mustard, Black
Mustard, Wild
Nightshade, American Black
Nightshade, Hairy
Parthenium
Phyllanthus, Longstalked
Pigweed, Prostrate
Pigweed, Redroot
Plantain, Paleseed
Puncturevine
Redmaids
Rocket, London
Ryegrass, Italian
Ryegrass, Perennial
Sandbur
Sedge, Annual
Sesbania, Hemp
Sida, Prickly (Teaweed)
Southern Brassbuttons
Sowthistle, Annual
Speedwell, Corn
Spurge, Prostrate
Sunflower, Common
Swinecress
Tassel Flower
Velvetleaf
Vetch, Purple
Wild Carrot
Willowherb
Woodsorrel (Oxalis)

KILLS AND PREVENTS

Burnweed, American
Chickweed, Common
Chickweed, Mouseear
Clover, White
Dandelion, Common
Dock, Curly
Eclipta
Henbit
Knotweed, Prostrate
Kyllingua, Annual*
Lambsquarters
Medic, Black
Plantain, Buckhorn
Purslane, Common
Pusley, Florida
Ragweed, Common
Shepardspurse
Spurge, Spotted
Thistle, Canada
Woodsorrel, Yellow

** Suppression*

ENVIRONMENTAL HAZARDS

- This product is toxic to fish, aquatic invertebrates, and plants. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean water mark. Do not contaminate water when disposing of rinsate or washwater. This product may impact water through spray drift or runoff. Follow directions for use to avoid spray drift and runoff. A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams and springs will reduce the potential of this product entering water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.
- ***Ground Water Advisory:*** This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.
- ***Surface Water Advisory:*** This pesticide may impact water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application.

PELIGROS AMBIENTALES

- Este producto es tóxico para los peces, los invertebrados acuáticos y las plantas. No aplicar directamente al agua o en áreas donde haya aguas superficiales o áreas entre mareas por debajo de la marca de agua. No contaminar el agua al eliminar el agua de lavado o enjuague. Este producto puede afectar el agua a través del rocío o derrames. Seguir las instrucciones de uso para evitar rocíos y derrames. Un buen mantenimiento de la zona de contención de vegetación entre las áreas en las que se aplica este producto y la diferenciación del agua de superficie, incluidos estanques, arroyos y manantiales, reduce el potencial de penetración en el agua a causa de precipitación-derrame. El derrame de este producto se reducirá al evitar las aplicaciones cuando se pronostica lluvia dentro de las 48 horas.
- ***Recomendación sobre las napas subterráneas:*** Este pesticida contiene propiedades y características relacionadas con sustancias químicas detectadas en aguas subterráneas. Este producto químico puede filtrarse en las napas subterráneas si se utiliza en áreas donde los suelos son permeables, particularmente donde el nivel freático es poco profundo.
- ***Recomendación sobre el agua de superficie:*** Este pesticida puede impactar sobre la calidad del agua a causa de derrame de agua pluvial. Esto se cumple especialmente en suelos con drenaje deficiente y suelos con napas subterráneas poco profundas. Este producto se clasifica como con alto potencial para llegar al agua de superficie a través de derrames durante varios meses o más después de la aplicación.

PRECAUTIONARY STATEMENTS

CAUTION HAZARDS TO HUMANS & DOMESTIC ANIMALS

- Harmful if swallowed or absorbed through skin. • Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing. • Remove and wash contaminated clothing before reuse.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

After using this product:

- Rinse gloves before removing and wash hands. • If product gets on skin, wash thoroughly with soap & water.
- If product gets on clothing, remove clothing as soon as possible and launder clothing separately before reuse.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

DECLARACIONES DE PRECAUCIÓN

PRECAUCIÓN RIESGOS PARA SERES HUMANOS Y ANIMALES DOMÉSTICOS

- Es perjudicial si se ingiere o absorbe por la piel. • Causa irritación moderada en los ojos. • Evitar el contacto con los ojos, la piel o la ropa. • Quitarse la ropa contaminada y lavarla antes de volver a usar. • Cuando use este producto, use camiseta de mangas largas, pantalones largos, calcetines, calzado y guantes de goma.

Después de usar este producto:

- Enjuagar los guantes antes de quitárselos y lavarse las manos. • Si el producto entra en contacto con la piel, lavarse bien con agua y jabón. • Si el producto entra en contacto con la ropa, quitarse la ropa tan pronto como sea posible y lavarla por separado antes de volver a utilizar. • Lavarse bien con agua y jabón después de manipular y antes de comer, beber, consumir goma de mascar, tabaco o usar el baño

FIRST AID

IF SWALLOWED
- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING
- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 – 20 minutes.
- Call a poison control center or doctor for further treatment advice.

IF IN EYES
- Hold eye open and rinse slowly and gently with water for 15 – 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for further treatment advice.

NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. **You may call toll-free 1-877-229-3763 for medical emergency information.**
Active Ingredients: 0.088% Dicamba, (CAS # 1918-00-9), 0.041% Penoxsulam (CAS # 219714-96-2) and 0.027% Indaziflam (CAS # 950782-86-2).

PRIMEROS AUXILIOS

EN CASO DE INGESTIÓN • Llamar inmediatamente a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento. • Hacer que la persona beba tragos de agua, si puede tragar. • No inducir el vómito salvo que así lo indique el médico o el centro de control de envenenamiento. • No administrar nada por vía oral a una persona inconsciente.

EN CASO DE CONTACTO CON LA PIEL O LA ROPA • Quitar la ropa contaminada. • Enjuagar la piel inmediatamente con abundante agua durante 15 a 20 minutos • Llamar a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento.

EN CASO DE CONTACTO CON LOS OJOS • Mantener abiertos los ojos y enjuagar de forma lenta y suave con agua durante entre 15 y 20 minutos. • Retirar los lentes de contacto, si los hubiera, después de los primeros 5 minutos y luego continuar enjuagando los ojos. • Llamar a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento.

NOTA: Tener la etiqueta del producto a mano al llamar a un centro de control de envenenamiento. Si busca tratamiento en una sala de emergencias o en el consultorio de un médico, lleve esta etiqueta del producto para mostrársela al personal médico. **También puede llamar al número gratuito 1-877-227-3763 para solicitar información médica de emergencia.**
Ingredientes activos: 0.088% de dicamba, (CAS n.°1918-00-9), 0.041% de penoxsulam (CAS n.°219714-96-2) y 0.027% de indaziflam (CAS n.°950782-86-2).

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage and disposal

PESTICIDE STORAGE Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.

PESTICIDE DISPOSAL & CONTAINER HANDLING **If Empty:** Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.
If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

ALMACENAMIENTO Y ELIMINACIÓN

No contaminar el agua o los alimentos al almacenar o eliminar

ALMACENAMIENTO DE PESTICIDAS Almacenar en el contenedor original en un lugar fresco y seco, fuera del alcance de los niños, preferentemente en un gabinete bajo llave. Proteger del congelamiento.

ELIMINACIÓN DEL PESTICIDAS Y MANIPULACIÓN DE ENVASES
Si está vacío: Envase no rellenable. No rellenar ni volver a utilizar este envase. Colocar el envase vacío en la basura o entregarlo para reciclar si fuera posible.
Si está parcialmente lleno: Llamar a la agencia local de residuos sólidos o al número gratuito 1-877-229-3724 para obtener indicaciones para la eliminación Nunca colocar el producto sin utilizar debajo de algún desagüe interno o externo.

6 87073 04840 7

EPA Reg. No. 72155-112
EPA Est. No. indicated by 3rd and 4th digits of the batch number on this package.
(26) = 82757-FL-1 (57) = 071106-GA-003

Bayer Advanced 3-in-1 Weed & Feed for Southern Lawns 35-0-3

GUARANTEED ANALYSIS
Total Nitrogen (N) 35.0%
35% Urea Nitrogen*
Soluble Potash (K_2O) 3.0%
Sulfur (S) Total 4.6%
3.5% Free Sulfur (S)
1.1% Combined Sulfur (S)

F1514

Derived From: Urea, Polymer Coated Sulfur Coated Urea, & Potassium Sulfate
*Contains 17.5% slowly available Nitrogen from Polymer Coated Sulfur Coated Urea

Information regarding the contents and levels of metals in this product are available on the Internet at http://www.aapfco.org/metals.htm.

Check with your county or city government to determine if there are local regulations for fertilizer use.

Produced for: Bayer Advanced, A Division of Bayer CropScience LP, P.O. Box 12014, 2 T.W. Alexander Drive, Research Triangle Park, NC 27709

Bayer (reg'd), the Bayer Cross (reg'd) and Bayer Advanced™ are trademarks of Bayer