

PGR IV[®]



Department of Agriculture
STATE OF HAWAII

LICENSED

LICENSED

PERIOD 2011-2013 LIC. NO.

PERIOD 2014-2016 LIC. NO.

8143.38

ACTIVE INGREDIENTS

**Indolebutyric acid	0.0028%
***Gibberellic acid	0.0030%
Other Ingredients	99.9942%
TOTAL	100.0000%

**Contains 0.8 mg indolebutyric acid/fl.oz.

***Contains 0.9 mg gibberellic acid/fl.oz.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2008-2010 LIC. NO.

KEEP OUT OF REACH OF CHILDREN CAUTION

See inside booklet for additional Precautionary Statements

For product information: 1-866-761-9397

Manufactured for:

Arysta LifeScience North America Corporation

15401 Weston Parkway, Suite 150

Cary, NC 27513

EPA Reg. No. 66330-267

EPA Est. No. 51036-GA-001

AD072602-B

101770

Net Contents: 1 Gallon



Arysta LifeScience[™]

North America

FIRST AID

IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present after the first 5 minutes, then continue rinsing eye.
HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. FOR 24- HOUR EMERGENCY MEDICAL ASSISTANCE CALL: 1-866-303-6952 FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

1. Long-sleeved shirt and long pants
2. Waterproof gloves
3. Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Protect from freezing. Store out of direct sunlight.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

EXCEPTION: If the product is soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is:

1. Coveralls over long-sleeved shirt and long pants
2. Waterproof gloves
3. Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: For one gallon plastic; Triple rinse (or equivalent). Then offer for recycling or puncture and dispose in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL CHEMIGATION INSTRUCTIONS

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduce pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quickclosing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreased to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with the pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

The pesticide supply tank should be agitated throughout the application of **PGR IV**. **PGR IV** should be applied at the end of the water application.

PGR IV should be applied in sufficient water to provide thorough and even coverage without producing excessive runoff.

SPRINKLER CHEMIGATION

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

DIRECTIONS FOR APPLICATION

When a range of rates is given, use the high rate when growing crops in highly organic soil.

PGR-IV may be applied by ground or air. Product should be mixed with enough water volume to provide even and equal distribution. (Check with your product representative).

APPLICATIONS DURING WATER DEFICIT STRESS ARE NOT BENEFICIAL.

FIELD CROPS

University trials and on farm results have shown the use of **PGR IV** can result in increased root mass, earlier fruit initiation, increased fruit retention, increased nutrient utilization, and higher yields.

COTTON CROP INFORMATION

PGR IV has a versatile range of use options on cotton. The maximum benefits of **PGR IV** are obtained by using an early planned program.

COTTON:

OPTION A –

- Step 1 - Apply 1 - 2 fl. oz./acre in seed furrow
- Step 2 - Apply 4 fl. oz./acre at pinhead square
- Step 3 - Apply 4 - 8 fl. oz./acre during active bloom period.

OPTION B –

- Step 1 - Apply 1 - 2 fl. oz./acre on a 10" - 12" band at the 2 - 5 leaf stage.
- Step 2 - Apply 4 fl. oz./acre at pinhead square.
- Step 3 - Apply 4 - 8 fl. oz./acre during active bloom period.

OPTION C –

- Step 1 - Apply 4 fl. oz./acre at pinhead square
- Step 2 - Apply 4 - 8 fl. oz./acre during active bloom period.

OPTION D –

- For fields with a history of rank growth.
- Step 1 - Apply 1 - 2 fl. oz./acre in furrow or 2 fl. oz./acre in a band at first true leaf.

NOTE: In furrow application is preferred option.

- Step 2 - Apply 2 fl. oz./acre at the 2 - 7 leaf stage.
- Step 3 - Apply 4 fl. oz./acre at pinhead square.
- Step 4 - Apply 4 - 8 fl. oz./acre during active bloom period.*

*Evaluate crop 1 to 2 weeks later for application of a growth retardant.

FIELD CORN:

OPTION A – Broadcast 8 fl. oz./acre preemergence (can be mixed with your herbicide).

OPTION B –

- STEP 1 - Apply 2 fl. oz./acre in seed furrow at planting
- STEP 2 - Band (16") 4 fl. oz./acre (set nozzles to assure even distribution) at 5 to 8 leaf stage.

MILO:

OPTION A – Broadcast 8 fl. oz./acre at 4 to 5 leaf stage.

OPTION B –

STEP 1 - Apply 2 fl. oz./acre in seed furrow at planting

STEP 2 - Band (16") 3 to 4 fl. oz./acre (set nozzles to assure even coverage) at 5 to 7 leaf stage.

SOYBEANS:**Split Application:**

STEP 1 - Apply 4 fl. oz./acre broadcast preplant incorporated with herbicide.

STEP 2 - Apply second application 4 fl. oz./acre at early bloom.

WINTER WHEAT, OATS, BARLEY, RYE:**Use options:**

If wheat is to be winter grazed: Apply 8 fl. oz./acre within two weeks postemergence.

If not grazed:

STEP 1 - Apply 4 fl. oz./acre within two weeks post emergence.

STEP 2 - Boost with 4 fl. oz./acre at early tillering.

PEANUTS:**OPTION A** –

Step 1 - Apply 1 fl. oz./acre in furrow

Step 2 - Apply 6 fl. oz./acre of **PGR IV** at initial pegging.

OPTION B –

Step 1 - Apply 4 fl. oz./acre at 15 days after emergence.

Step 2 - Apply 4 fl. oz./acre at 30 days after emergence.

Step 3 - Apply 4 fl. oz./acre at initial pegging.

NOTE: Tank mixes with 2,4-DB or other hormone type herbicides may result in unwanted additional herbicidal activity.

RICE:**OPTION A** –

Step 1 - Apply 6 fl. oz./acre at 3 to 5 leaf stage.

OPTION B –

Step 1 - Apply 4 fl. oz./acre at 3 to 5 leaf stage.

Step 2 - Apply 4 fl. oz./acre at panicle initiation.

VEGETABLE CROPS*****PEPPERS: BELL, JALAPENO, SERRANO, etc.**

STEP 1 - Band 2 fl. oz./acre 2 weeks post transplant.

STEP 2 - Broadcast 2 fl. oz./acre at early bloom.

STEP 3 - Broadcast 1 to 2 fl. oz./acre at 21 day intervals.

BEANS: GREEN, LIMA, PINTO, BLACKEYE, MUNG, ETC.

Spray 6 fl. oz./acre at first bloom (broadcast).

*****CABBAGE/BROCCOLI/CAULIFLOWER/BRUSSEL SPROUTS**

STEP 1 - Band (14") 1 to 2 fl. oz./acre 2 weeks post transplant.

STEP 2 - Boost (14" band) 1 to 2 fl. oz./acre 4 weeks post transplant.

CUCUMBERS/SQUASH

STEP 1 - Spray 3 fl. oz./acre at 2 to 4 leaf stage - broadcast.

STEP 2 - Boost with 3 fl. oz./acre at early bloom - broadcast.

STEP 3 - Boost again with 2 fl. oz./acre at 14 to 21 day intervals- broadcast.

LETTUCE/SPINACH/TURNIP/MUSTARD GREENS

Spray 2 fl. oz./acre in furrow - OR - in 14" band over the top at the three leaf stage.

Spinach and greens - boost with 2 fl. oz./acre after each cutting.

SWEET CORN

Two programs have produced good results: 8 fl. oz./acre in combination with herbicide - preemergence OR

STEP 1 - 2 fl. oz./acre in furrow at planting

STEP 2 - 4 fl. oz./acre band at 5 to 7 leaf stage (12" to 16").

*****TOMATOES: Machine Harvested**

STEP 1 - Spray 2 fl. oz./acre (14" band) 3 weeks post transplant.

STEP 2 - Spray 6 fl. oz./acre (broadcast) at early bloom.

*****TOMATOES: Hand Picked**

STEP 1 - Spray 2 fl. oz./acre (14" band) 3 weeks post transplant.

STEP 2 - Broadcast 2 fl. oz./acre 3 weeks later.

STEP 3 - Broadcast 1 fl. oz./acre in combination with foliar fertilizer every 21 days.

POTATOES - RUSSET

OPTION A - Add 4 fl. oz./acre to your fertilizer and incorporate prior to planting.

OPTION B - Broadcast 4 to 6 fl. oz./acre at 45 days after emergence.

SUGAR BEETS: As an aid to increase total sugar content

1st APPLICATION: 2 fl. oz./acre Broadcast 6 to 8 leaf stage.

2nd APPLICATION: 30 days later - 2 fl. oz./acre broadcast.

SUGAR CANE: As an aid to increase total sugar content

APPLICATION: Side Dress 8 fl. oz./acre in combination with your fertilizer.

FRUIT CROPS

CANTALoupES/WATERMELON/HONEYDEW

STEP 1 - Spray 3 fl. oz./acre broadcast when plants show first signs of running.

STEP 2 - Boost 3 fl. oz./acre broadcast 2 weeks later.

YOUNG CITRUS

STEP 1 - To reduce transplant shock and get seedlings off to a faster start - use at the rate of 1 fl. oz./2 gallons of water - dip root or spray ball prior to set - then - spray foliage lightly.

STEP 2 - Using transplant solution - 1 fl. oz./2 gallons water - spray foliage of young trees to drip or run-off 2 to 3 times per year.

***STRAWBERRIES

STEP 1 - Spray 3 fl. oz./acre 3 to 4 weeks prior to coming out of dormancy - broadcast.

STEP 2 - Spray 3 fl. oz./acre at early bloom. Broadcast.

*****TRANSPLANTS** - for quick start - dip or spray roots with solution of 1 Tablespoon of PGR-IV per gallon of water prior to transplanting.

NOTICE: PGR-IV IS NOT A FERTILIZER. USE IN COMBINATION WITH A GOOD FERTILIZER PROGRAM WHERE INDICATED.

SEED TREATMENT

For use solely as an at planting treatment. Use of seed treated with **PGR-IV** at the recommended rates precludes the need for in furrow application of **PGR-IV** at planting time. Do not allow treated seed to be used for food or feed. Apply **PGR-IV** directly to the seed using the following rate table. Use suitable treatment equipment to obtain good and even coverage. Sufficient water may be added to the **PGR-IV** to insure uniform coverage. Improper coverage will minimize product performance.

These rates are based on achieving the equivalent of one to two fluid ounces of PGR-IV per acre based on average pounds of seed commonly used per acre at planting time.

SEED TYPE

Cotton

RATE/100 lbs seed

Apply 6 - 12 fl. oz. per 100 pounds seed.

Corn

Apply 6 fl. oz. per 100 pounds of seed

(field, sweet, popcorn)

Cereal Grains

Apply 1 fl. oz. per 100 pounds of seed.

(barley, oats, rice, rye, wheat)

Milo

Apply 10 fl. oz. per 100 pounds of seed.

(sorghum)

Peanuts

Apply 1 fl. oz. per 100 pounds of seed.

COMMERCIAL SEED TREATMENT

PGR-IV may be applied as a water-based slurry with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.

NOTE: Federal law requires that bags containing treated seed shall be labeled with the following information:

"This seed has been treated with Arysta **PGR-IV**. Do not use treated seed for food, feed or oil purposes."

Store away from feeds and foodstuffs.

The purchaser of this product is responsible for insuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21 CFR, Part 2.25. Any dye added must be cleared for use under 40 CFR, Part 180.1001.

CONDITIONS OF SALE

It is impossible to eliminate all risks associated with the use of this product. Such risks may arise from weather conditions, soil factors, off-target movement, unconventional farming techniques, the presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of Arysta LifeScience North America Corporation ("Arysta"), and can cause crop injury, injury to non-target crops or plants, ineffectiveness of the product, or other unintended consequences. All such risks shall be assumed by the user or buyer.

Arysta warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to Arysta, and is subject to the inherent risks described above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARYSTA DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARYSTA, MANUFACTURER, AND SELLER DISCLAIM AND SHALL NOT BE LIABLE FOR ANY SPECIAL, INCIDENTAL, INDIRECT, OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE, HANDLING, APPLICATION, STORAGE, OR DISPOSAL OF THIS PRODUCT OR FOR DAMAGES IN THE NATURE OF PENALTIES, AND THE USER AND BUYER WAIVE ANY RIGHT THAT THEY MAY HAVE TO SUCH DAMAGES. NO AGENT, REPRESENTATIVE OR EMPLOYEE OF ARYSTA IS AUTHORIZED TO MAKE ANY WARRANTY, GUARANTEE OR REPRESENTATION BEYOND THOSE CONTAINED HEREIN OR TO MODIFY THE WARRANTIES CONTAINED HEREIN. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE TOTAL LIABILITY OF ARYSTA, MANUFACTURER, AND SELLER, SHALL BE LIMITED TO THE PURCHASE PRICE PAID, OR AT ARYSTA'S ELECTION, THE REPLACEMENT OF THE PRODUCT.

PGR-IV is a registered trademark of Arysta LifeScience North America Corporation.

NOTES

NOTES

PGR IV®

ACTIVE INGREDIENTS

**Indolebutyric acid	0.0028%
***Gibberellic acid	0.0030%
Other Ingredients	99.9942%
TOTAL	100.0000%

**Contains 0.8 mg indolebutyric acid/fl.oz.

***Contains 0.9 mg gibberellic acid/fl.oz.

KEEP OUT OF REACH OF CHILDREN CAUTION

See inside booklet for additional Precautionary Statements and Directions For Use

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: For one gallon plastic; Triple rinse (or equivalent). Then offer for recycling or puncture and dispose in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

HOT LINE NUMBERS

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

FOR 24- HOUR EMERGENCY MEDICAL ASSISTANCE CALL: 1-866-303-6952

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

PGR-IV is a registered trademark of Arysta LifeScience North America Corporation

For product information: 1-866-761-9397

Manufactured for:

Arysta LifeScience North America Corporation

15401 Weston Parkway, Suite 150

Cary, NC 27513

EPA Reg. No. 66330-267

AD072602-B 101770

Net Contents: 1 Gallon

EPA Est. No. 51036-GA-001



Arysta LifeScience™

North America