

STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO.

8143.65

Herbicide

GROUP 2 HERBICIDE

Water Dispersible Granule
For Use on Wheat, Barley, and Fallow

ACTIVE INGREDIENTS:

Thifensulfuron-methyl

Methyl 3-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl) amino]carbonyl]amino]sulfonyl]-2-thiophenecarboxylate 25.0%

Tribenuron-methyl

Methyl 2-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)methylamino]carbonyl]amino]sulfonyl]benzoate 25.0%

OTHER INGREDIENTS: 50.0%

TOTAL: 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

See booklet for First Aid, additional Precautionary Statements and Directions for Use.

Produced for:

ARYSTA LIFESCIENCE NORTH AMERICA, LLC

15401 Weston Parkway, Suite 150

Cary, North Carolina 27513

103356—031413

EPA Registration No. 66330-418

EPA Establishment No. 69016-MS-001

AD021413

NET CONTENTS: 32 OUNCES



Arysta LifeScience

FIRST AID

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15–20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE: Call PROSAR at
1-866-303-6952 or 1-651-632-8946 if calling from outside the U.S.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC at
1-800-424-9300 or 1-703-527-3887 if calling from outside of the U.S.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options follow the instructions for Category A on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, made of any waterproof material such as natural rubber.
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Ensure that all operation employees accurately measure pesticides.
- Mix only enough product for the job at hand.
- Avoid overfilling of spray tank.
- Do not discharge excess material on the soil at a single spot in the field, grove, or mixing/loading station.
- Dilute and agitate excess solution and apply at labeled rates or uses.
- Avoid storage of pesticides near well sites.
- When triple-rinsing the pesticide container, be sure to add the rinsate to the spray mix.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

This product is for use on wheat, barley and fallow in many states. Check with your state extension or Dept. of Agriculture before use, to be certain this product is registered in your state. Arysta LifeScience North America, LLC will not be responsible for losses or damages resulting from the use of this product in any manner not in accordance with instructions on this label.

USE INFORMATION

This product is a water dispersible granule that is used for selective postemergence weed control in wheat (including durum), barley, and fallow. The best control is obtained when this product is applied to young, actively growing weeds. The specified use rate will depend on weed spectrum and size of weed at time of application. The degree and duration of control may depend on the following:

- weed spectrum and infestation intensity
- weed size at application
- environmental conditions at and following treatment

This product is noncorrosive, nonflammable, nonvolatile, and does not freeze. This product should be mixed, and completely dissolved in water and applied as a uniform broadcast spray.

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

This product is absorbed primarily through the foliage of plants, rapidly inhibiting the growth of susceptible weeds. One to three weeks after application to weeds, leaves of susceptible plants appear chlorotic, and the growing point subsequently dies.

This product provides the best control in vigorously growing crops that shade competitive weeds. Weed control in areas of thin

crop stand or seeding skips may not be as satisfactory. However, a crop canopy that is too dense at application can intercept spray and reduce weed control.

The herbicidal action of this product may be affected in crops stressed from adverse environmental conditions (such as extreme temperatures or moisture), abnormal soil conditions, cultural practices, or variations in crop variety. In warm, moist conditions, the expression of herbicide symptoms is accelerated; in cold, dry conditions, expression of herbicide symptoms is delayed. In addition, weeds hardened-off by drought stress are less susceptible to this product.

APPLICATION INFORMATION

USE RATE

Apply this product at a rate of 0.4 to 1 ounce per acre. When applying 0.4 to 0.6 ounce per acre, this product must be used in a tank-mix combination with other registered herbicides.

WHEAT (INCLUDING DURUM) AND BARLEY

Apply 0.4 to 1 ounce of this product per acre to wheat (including durum) or barley. The total amount of this product cannot exceed 1 ounce per acre per crop season.

FALLOW

Apply 0.4 to 1 ounce of this product per acre to fallow. The total amount of this product cannot exceed 1 ounce per acre per crop season. This product should be applied in combination with other suitable registered fallow herbicides such as glyphosate plus 2,4-D (ester formulations work best) or glyphosate plus dicamba.

When this product is applied at a rate of 0.4 to 0.6 ounce per acre, this product must be used in a tank mix combination with other registered fallow herbicides.

PRE-PLANT BURNDOWN

Apply 0.4 to 1 ounce of this product per acre as a burndown treatment prior to, or shortly after planting (prior to emergence). The total amount of this product cannot exceed 1 ounce per acre per crop season.

Apply up to 0.6 ounces per acre of this product as a pre-plant burndown to Cotton. Allow at least 14 days from time of application to time of planting cotton.

POST HARVEST

Apply this product at 0.4 to 1.0 ounce per acre to crop stubble after harvest. Use the 1.0 ounce per acre rate when weed infestation is heavy and predominantly consists of those weeds listed under the **WEEDS PARTIALLY CONTROLLED** section of this label or when application timing and environmental conditions are marginal. (See the **APPLICATION TIMING** section of this label for restriction

on planting intervals). This product should be applied in combination with other suitable registered burndown herbicides (See the **TANK MIXTURES** section of this label for additional information).

Sequential treatments of this product may also be made provided the total amount of this product applied during one fallow/pre plant cropland season does not exceed 1.0 ounce per acre.

APPLICATION TIMING

WHEAT (INCLUDING DURUM) AND BARLEY

Make applications after the crop is in the 2-leaf stage, but before the flag leaf is visible.

FALLOW

This product may be used as a fallow treatment, in the spring, summer or fall when the majority of weeds have emerged and are actively growing.

PRE-PLANT BURNDOWN

Apply this product as a burndown treatment to wheat (including durum) and barley to control emerged weeds prior to, or shortly after planting (prior to emergence). Make applications when the majority of weeds have emerged and are actively growing. Wheat and Barley may be replanted anytime after the application of this product.

Allow at least 14 days between application of this product and planting of cotton. Allow at least 60 days between application of this product and planting of sugar beets, winter rape and canola. Allow at least 45 days between application of this product and planting of any other crop (such as corn, rice, grain sorghum or soybeans).

POST HARVEST

This product may be used as a burndown treatment to crop stubble when the majority of weeds have emerged and are actively growing. (See the **CROP ROTATION** section of this label for additional information.)

Since this product has very little or no soil activity, it controls only those weeds that have germinated; therefore, apply this product when all or most of the weeds have germinated. Annual broadleaf weeds should be past the cotyledon stage, actively growing, and less than 4" tall or wide. Rainfall immediately after treatment can wash this product off of weed foliage, resulting in reduced weed control. Several hours of dry weather are needed to allow this product to be sufficiently absorbed by weed foliage.

CROP ROTATION

Wheat (including durum) and barley may be replanted anytime after the application of this product.

Cotton can be planted 14 days after the application of this product.

Sugar beets, Winter Rape and Canola can be planted at 60 days after the application of this product. Any other crop may be planted 45 days after the application of this product.

WEEDS CONTROLLED

This product effectively controls the following weeds when used according to label directions:

Annual knawel	Common sunflower	London rocket	Slimleaf lambsquarters
Annual sowthistle	Corn chamomile	Marshelder	Smallflower buttercup
Black mustard	Corn gromwell*	Mayweed chamomile	Smallseed falseflax
Blue/Purple mustard	Corn spurry	Miners lettuce	Stinking chickweed
Broadleaf dock	Cow cockle	Narrowleaf lambsquarters	Stinking mayweed/Dogfennel
Bur buttercup	Cress (mouse ear)	Nightflowering catchfly	Sunflower
Bushy wallflower/ Treacle mustard	Curly dock	Pennsylvania smartweed	Swinecress
Canada Thistle*	False chamomile	Pineappleweed	Tansymustard
Clasping pepperweed	Field chickweed	Prickly lettuce*	Tarweed fiddleneck
Coast fiddleneck	Field pennycress	Prostrate knotweed	Tumble/Jim Hill mustard
Common buckwheat	Filaree (redstem, Texas)	Prostrate pigweed	Volunteer canola
Common chickweed	Flixweed	Redmaids	Volunteer lentils
Common cocklebur*	Green smartweed	Redroot pigweed	Volunteer peas
Common groundsel	Henbit	Russian thistle*	Wild buckwheat*
Common lambsquarters	Kochia*	Scentless chamomile/ mayweed	Wild chamomile
Common ragweed*	Ladysthumb	Shepherd's-purse	Wild mustard
	Lanceleaf sage*		

WEEDS PARTIALLY CONTROLLED**
This product partially controls the following weeds when used according to label directions:
Catchweed bedstraw
Mallow (common, little)
Marestail
WEEDS CONTROLLED
Nightshade (cutleaf, hairy)
*See <i>SPECIFIC WEED PROBLEMS</i> for more information.
**Partial control: A visual reduction of weed population as well as a significant loss of vigor. For better results, use the highest rate of this product per acre and include a tank mix partner such as 2,4-D, MCPA, Buctril®/Maestro® 2EC or Banvel®/Clarity® or Diablo®/Clash™ (refer to <i>TANK MIXTURES</i>).

SPECIFIC WEED PROBLEMS

Canada Thistle: For control in wheat and barley, use 0.8 ounce per acre plus surfactant when all thistles are 4" to 8" with 2" to 6" of new growth. Make the application in the spring. Control will be improved by using this product in combination with 2,4-D or dicamba (refer to ***TANK MIXTURES***).

Common cocklebur, Common ragweed, Lanceleaf sage: In wheat and barley, apply this product at 0.4 to 0.8 ounce per acre in combination with 2,4-D at rates from ¼ to ¾ pound active ingredient (ester formulations work best) when weeds are small and actively growing. When using ¼ pound active ingredient of 2,4-D, be sure to add surfactant at the rate of ¼ to ½ quart per 100 gallons of spray solution (0.06 to 0.125% v/v—use the higher rate under stress conditions).

Corn gromwell, Wild buckwheat: For control in wheat and barley, use 0.8 ounce of this product per acre plus surfactant.

Kochia, Russian Thistle, Prickly lettuce: Naturally occurring resistant biotypes of these weeds are known to occur. For best results, use this product in a tank mix with Starane®/Comet®, Starane + Sword®, Starane + Salvo®, dicamba (such as Banvel®/Clarity or Diablo®/Clash) and 2,4-D; or Bromoxynil (such as Buctril®/Maestro 2EC) and 2,4-D (¾ to 1 pint Buctril®/Maestro 2EC + ¼ to ¾ pound active ingredient 2,4-D ester). This product should be applied in the spring when weeds are 2" to 4" tall or 2" to 4" across and are actively growing. Refer to the ***TANK MIXTURES*** section of this label for additional details.

SPRAY ADJUVANTS

Always include a spray adjuvant with applications of this product. In addition to a spray adjuvant, an ammonium nitrogen fertilizer may be used.

Consult your Ag dealer or applicator, local Arysta LifeScience North America, LLC fact sheets, technical bulletins, and service policies prior to using an adjuvant system. If another herbicide is tank mixed with this product, select adjuvants authorized for use with both products. Products must contain only EPA-exempt ingredients (40 CFR 1001).

NONIONIC SURFACTANT (NIS)

- Apply 0.06 to 0.50% volume/volume (½ to 4 pints per 100 gallon of spray solution).
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12. See **TANK MIXTURES** section of this label for additional information.

PETROLEUM CROP OIL CONCENTRATE (COC) OR MODIFIED SEED OIL (MSO)

- Apply at 1% volume/volume (1 gallon per 100 gallon of spray solution) or 2% volume/volume under arid conditions.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

SPECIAL ADJUVANT TYPES

- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions.
- In addition to the adjuvants specified above, other adjuvant types may be used if they provide the same functionality and have been evaluated and approved by Arysta LifeScience North America, LLC product management. Consult separate Arysta LifeScience North America, LLC technical bulletins for detailed information before using adjuvant types not specified on this label.

AMMONIUM NITROGEN FERTILIZER

- Use 2 qt/acre of a high-quality urea ammonium nitrate (UAN), such as 28%N or 32%N, or 2 lb/acre of a spraygrade ammonium sulfate (AMS). Use 4 qt/acre UAN or 4 lb/acre AMS under arid conditions.

GROUND APPLICATION

For optimum spray distribution and thorough coverage, use flat-fan or low-volume flood nozzles.

For flat-fan nozzles, use a spray volume of at least 5 gallon per acre (GPA).

For flood nozzles on 30" spacings, use at least 10 GPA, flood nozzles no larger than TK10 (or the equivalent), and a pressure of at least 30 psi. For 40" nozzle spacings, use at least 13 GPA; for 60" spacings use at least 20 GPA. It is essential to overlap the nozzles 100% for all spacings.

RA Raindrop® nozzles are not recommended for this product applications, as weed control performance may be reduced. Use screens that are 50-mesh or larger.

AERIAL APPLICATION

Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage.

- Use 2 to 5 GPA
- Use at least 3 GPA in Idaho, Oregon, or Utah.

Do not apply this product by air in the state of New York.

See the ***SPRAY DRIFT MANAGEMENT*** section of this label.

CHEMIGATION

Do not apply this product through any irrigation system.

PRODUCT MEASUREMENT

This product is measured using this product volumetric measuring cylinder. The degree of accuracy of this cylinder varies by +/- 7.5%. For more precise measurement, use scales calibrated in ounces.

TANK MIXTURES

This product may be tank mixed with other suitable registered herbicides to control weeds listed as suppressed, weeds resistant to this product or weeds not listed under ***WEEDS CONTROLLED***. Read and follow all manufacturer's label instructions for the companion herbicide. If those instructions conflict with this label, do not tank mix the herbicide with this product.

This product can also be mixed with registered fungicides, insecticides, or liquid fertilizer for use on wheat, barley, or fallow.

WITH 2,4-D (AMINE OR ESTER) OR MCPA (AMINE OR ESTER)

This product may be tank mixed with the amine or ester formulations of 2,4-D or MCPA herbicides for use on wheat and barley. For best results in the Red River Valley and adjacent areas of North Dakota and Minnesota, add the ester formulations of 2,4-D or MCPA herbicides to the tank at $\frac{3}{8}$ pound active ingredient (such as $\frac{3}{4}$ pint of a 4 lb/gal product, or $\frac{1}{2}$ pint of a 6 lb/gal product). No additional surfactant is needed with this mixture.

For best results in other areas, add the ester formulations of 2,4-D or MCPA herbicides to the tank at $\frac{1}{4}$ to $\frac{3}{8}$ pound active ingredient (such as $\frac{1}{2}$ to $\frac{3}{4}$ pt of a 4 lb/gal product, or $\frac{1}{4}$ to $\frac{1}{2}$ pt of a 6 lb/gal product). Surfactant may be added to the mixture at $\frac{1}{2}$ to 1 quart per 100 gallons of spray solution (0.125 to 0.25% v/v); however, adding surfactant may increase the potential for crop injury, especially at the higher phenoxy rates.

Higher rates of 2,4-D or MCPA may be used, but do not exceed the highest rate allowed by those respective labels. Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using these tank mixtures.

WITH DICAMBA (SUCH AS BANVEL/CLARITY, DIABLO/CLASH)

This product may be tank mixed with $\frac{1}{16}$ to $\frac{1}{8}$ pound active ingredient dicamba (such as 2 to 4 fluid ounces Banvel/Diablo, or 2 to 4 fluid ounces Clarity/Clash). Use higher rates when weed infestation is heavy. Surfactant may be added to the mixture at $\frac{1}{2}$ to 1 quart per 100 gallons of spray solution (0.125 to 0.25% v/v); however, adding surfactant may increase the potential for crop injury. Refer to the specific dicamba label for application timing and restrictions. Tank mixes of this product plus dicamba may result in reduced control of some broadleaf weeds.

WITH 2,4-D (AMINE OR ESTER) AND BANVEL/CLARITY, DIABLO/CLASH)

This product may be applied in a 3-way tank mix with formulations of dicamba and 2,4-D. Make application of this product + $\frac{1}{16}$ to $\frac{1}{8}$ pound active ingredient dicamba (such as 2 to 4 ounces Banvel/Diablo, or 2 to 4 ounces Clarity/Clash) + $\frac{1}{4}$ to $\frac{3}{8}$ pound active ingredient of 2,4-D ester or amine per acre. Use higher rates when weed infestation is heavy. Surfactant may be added to the mixture at $\frac{1}{2}$ to 1 quart per 100 gallons of spray solution (0.125 to 0.25% v/v); however, adding surfactant may increase the potential for crop injury. Consult the specific 2,4-D label, dicamba label, or local recommendations for more information and restrictions.

Apply this 3-way combination to winter wheat after the crop is tillering and prior to jointing (first node). In Spring Wheat (including Durum) apply after the crop is tillering and before it exceeds the 5-leaf stage.

In Spring Barley, apply after the crop is tillering and before it exceeds the 4-leaf stage.

WITH BROMOXNYL (SUCH AS BUCTRIL/MAESTRO 2EC/BRONATE®/MAESTRO MA)

This product may be tank mixed with bromoxnyl containing herbicides registered for use on wheat, barley, or fallow. For best results, add bromoxnyl containing herbicides to the tank at $\frac{3}{16}$ to $\frac{3}{8}$ pound active ingredient per acre (such as Bronate/Maestro MA/Buctril/Maestro 2 EC at $\frac{3}{4}$ to 1 $\frac{1}{2}$ pint per acre).

Read and follow all label instructions on timing, precautions, and warnings for these herbicides before using these tank mixtures. Follow the most restrictive labeling. Tank mixes of this product plus Buctril/Maestro 2EC may result in reduced control of Canada thistle.

WITH STARANE/COMET, STARANE+SWORD, STARANE+SALVO

For improved control of Kochia (2 to 4" tall), Russian thistle, mustard species, and wild buckwheat this product may be tank mixed with $\frac{1}{2}$ to 1 $\frac{1}{2}$ pints per acre of Starane/Comet, $\frac{2}{3}$ to 2 $\frac{2}{3}$ pints per acre of Starane+Salvo or $\frac{3}{4}$ to 2 $\frac{3}{4}$ pints per acre of Starane+Sword. Additional 2,4-D or MCPA can be added based on local recommendations (refer to 2,4-D and MCPA labels for maximum amount that can be applied to the crop). Refer to the Starane/Comet, Starane+Salvo or Starane+Sword label for information regarding use restrictions, labeled crops, rotational cropping recommendations, sprayer cleanup, use precautions and other information. The most restrictive provisions on any label will apply. Do not use the tank mix if any restrictions on the labels conflict with instructions on this product label.

WITH HOELON® HERBICIDE

This product may be used in combination with Hoelon 3EC and Buctril/Maestro 2EC herbicides in accordance with the Hoelon 3EC label. For best results, use the three-way tank mix of this product at 0.4 ounce per acre plus Hoelon 3 EC at 2½ pint per acre plus Buctril/Maestro 2EC at 1½ pints per acre. Apply only to winter wheat. This tank mix should only be used under good soil conditions when wild oat is in the 1 to 4 leaf stage. If conditions are not ideal for the performance of Hoelon 3EC, wild oat control may be reduced. Be sure to follow all warnings and cautions on the Hoelon 3EC and Buctril/Maestro 2EC labels.

WITH ASSERT® HERBICIDE

This product can be tank mixed with Assert. When tank mixing this product with Assert, always include another broadleaf weed herbicide with a different mode of action (for example: 2,4-D ester, MCPA ester, Buctril/Maestro 2EC, or Bronate/Maestro MA). Tank-mixed applications of this product plus Assert may cause temporary crop discoloration, stunting, or injury when heavy rainfall occurs shortly after application.

WITH OTHER GRASS CONTROL PRODUCTS

Tank mixtures of this product and grass control products may result in poor grass control. Arysta LifeScience North America, LLC recommends that you first consult your state experiment station, university, or extension agent, Agricultural dealer, or Arysta LifeScience North America, LLC representative as to the potential for antagonism before using the mixture. If no information is available, limit the initial use of this product and the grass product to a small area.

Do not tank mix with Achieve Herbicide.

WITH INSECTICIDES OR FUNGICIDES

This product may be tank mixed or used sequentially with insecticides (or fungicides) registered for use on cereal grains. However, under certain conditions (drought stress, or if the crop is in the 2- to 4- leaf stage), tank mixes or sequential applications of this product with organophosphate insecticides (such as parathion) may produce temporary crop yellowing or, in severe cases, crop injury. Test these mixtures in a small area before treating large areas. However, review all insecticide and fungicide labels for restrictions.

Do not use this product plus Malathion, as crop injury will result.

WITH LIQUID NITROGEN SOLUTION FERTILIZER

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing this product in fertilizer solution. Do not add this product directly to liquid nitrogen fertilizer; the granules will not dissolve. This product must be thoroughly mixed with clean water before it is added to liquid nitrogen fertilizer. If granules remain when the mixture is poured out, add more clean water and mix until all granules have disappeared. Ensure that the agitator is running when this product premix is added. Use of this mixture may result in temporary crop yellowing and stunting.

If using low rates of liquid nitrogen fertilizer in the spray solution (less than 50% of the spray solution volume), the addition of

surfactant is necessary. Add surfactant at ¼ to 1 quart per 100 gallon of spray solution (0.06 to 0.25% v/v) based on local recommendations. When using high rates of liquid nitrogen fertilizer solution in the spray solution, adding surfactant increases the risk of crop injury. Consult your agricultural dealer, consultant, fieldman, or Arysta LifeScience North America, LLC representative for a specific recommendation before adding an adjuvant to these tank mixtures.

If 2,4-D or MCPA is included with this product and fertilizer mixture, ester formulations tend to be more compatible (see manufacturer's label). Additional surfactant is not needed when using this product in tank mix with 2,4-D ester or MCPA ester and liquid nitrogen fertilizer solutions.

Do not use low rates of liquid nitrogen fertilizer solution as a substitute for a surfactant.

Do not use with liquid fertilizer solutions with a pH less than 3.0.

TANK MIXTURES IN FALLOW

This product may be used as a fallow treatment, and may be tank mixed with other herbicides that are registered for use in fallow. Read and follow all manufacturer's label instructions for the companion herbicide. If those instructions conflict with this label, do not tank mix the herbicide with this product.

TANK MIXTURES IN PRE-PLANT BURNDOWN

This product may be used as a pre-plant burndown treatment alone or tank mixed with other herbicides that are registered for use as a pre-plant burndown product, such as glyphosate plus 2,4-D (ester formulations work best) or glyphosate plus dicamba. Read and follow all manufacturer's label instructions for the companion herbicide. If those instructions conflict with this label, follow the most restrictive labeling (such as planting interval after application), or do not tank mix the herbicide with this product.

TANK MIXTURES IN POST HARVEST APPLICATIONS

This product may be used as a post harvest treatment to crop stubble, and may be tank mixed with other herbicides that are registered for use in fallow.

MIXING INSTRUCTIONS

Do not use with spray additives that alter the pH of the spray solution below pH 5.0 or above pH 9.0 as rapid product degradation can occur. Spray solutions of pH 6.0 to 8.0 allow for optimum stability of this product.

1. Fill the tank ¼ to ½ full of water.
2. While agitating, add the required amount of this product.
3. Continue agitation until this product is fully dissolved, at least 5 minutes.
4. Once this product is fully dissolved, maintain agitation and continue filling tank with water.

5. As the tank is filling, add tank mix partners and then add the required volume of spray adjuvant. Always add spray adjuvant last. Antifoaming agents may be used. Do not use with spray additives that alter the pH of the spray solution below pH 6.0 as rapid product degradation can occur. Spray solutions of pH 7.0 and higher allow for optimum stability of this product.
6. Dispersed tank mix partners can settle if the tank mixture is not continually agitated. If settling occurs, thoroughly re-agitate before using.
7. Apply this product spray mixture within 24 hours of mixing to avoid product degradation.
8. If this product and a tank mix partner are to be applied in multiple loads, fully dissolve this product in clean water prior to adding to the tank.

GRAZING

Do not graze livestock in treated areas. In addition, do not feed forage or hay from treated areas to livestock (straw harvested after grain harvest may be used for bedding and/or feed).

SPRAY EQUIPMENT

For specific application equipment, refer to the manufacturer's recommendations for additional information on GPA, pressure, speed, nozzle types and arrangements, nozzle heights above the target canopy, etc.

Be sure to calibrate air or ground equipment properly before application. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern with minimum drift. Use higher spray volumes to obtain better coverage when crop canopy is dense. Avoid swath overlapping, and shut off spray booms while starting, turning, slowing, or stopping to avoid injury to the crop.

Do not make applications using equipment and/or spray volumes or during weather conditions that might cause spray to drift onto nontarget sites. For additional information on spray drift refer to **SPRAY DRIFT MANAGEMENT** section of label.

Continuous agitation may be required to keep this product and tank-mix partners in solution or suspension. Refer to tank-mix partner labels for additional information.

SPRAYER CLEANUP

The spray equipment must be cleaned before this product is sprayed. Follow the cleanup procedures specified on the labels of the previously applied products. If no directions are provided, follow the six steps outlined in **AFTER SPRAYING** in this label.

AT THE END OF THE DAY

It is recommended that during periods when multiple loads of this product are applied, at the end of each day of spraying, the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses flushed. This will prevent the buildup of dried pesticide deposits which can accumulate in the application equipment.

AFTER SPRAYING AND BEFORE SPRAYING CROPS OTHER THAN WHEAT AND BARLEY

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of this product as follows:

1. Drain tank; thoroughly rinse spray tanks, boom and hoses with clean water. Loosen and physically remove any visible deposits.
 2. Fill the tank with clean water and 1 gal of household ammonia* (contains 3% active) for every 100 gal of water. Flush the hoses, boom and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 minutes. Flush the hoses, boom and nozzles again with the cleaning solution, and then drain the tank.
 3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
 4. Repeat step 2.
 5. Rinse the tank, boom, and hoses with clean water.
 6. If only ammonia is used as a cleaner, the rinsate solution may be applied back to the crop(s) recommended on this label. Do not exceed the maximum labeled use rate. If other cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.
- * Equivalent amounts of an alternate strength ammonia solution can be used in the cleanout procedure. Carefully read and follow the individual cleaner instructions. Consult your Ag dealer for a listing of approved cleaners.

Notes:

1. CAUTION: Do not use chlorine bleach with ammonia as dangerous gases will form. Do not clean equipment in an enclosed area.
2. Steam-cleaning aerial spray tanks is recommended prior to performing the above cleanout procedure to facilitate the removal of any caked deposits.
3. When this product is tank mixed with other pesticides, all cleanout procedures should be examined and the most rigorous procedure should be followed.
4. In addition to this cleanout procedure, all precleanout guidelines on subsequently applied products should be followed as per the individual labels.
5. Where routine spraying practices include shared equipment frequently being switched between applications of this product and applications of other pesticides to sensitive crops during the same spray season, it is recommended that a sprayer be dedicated to this product to further reduce the chance of crop injury.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

IMPORTANCE OF DROPLET SIZE

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

The most effective way to reduce drift potential is to apply large droplets (>150 to 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS! See **WIND**, **TEMPERATURE AND HUMIDITY**, and **TEMPERATURE INVERSIONS** sections of this label.

Controlling Droplet Size—GENERAL TECHNIQUES

Volume—Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure—Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.

Nozzle Type—Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Number of Nozzles—Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.

Nozzle Orientation—Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.

Nozzle Type—Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.

Boom Length—The boom length should not exceed $\frac{3}{4}$ of the wing or rotor length—longer booms increase drift potential.

Application Height—Application more than 10 feet above the canopy increases the potential for spray drift.

BOOM HEIGHT

Setting the boom at the lowest referenced height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given wind speed. **AVOID GUSTY AND WINDLESS CONDITIONS.**

NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature And Humidity

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

Temperature Inversions

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog. However, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring.

Note: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the **SPRAY EQUIPMENT** section of this label to determine if use of an air assist sprayer is recommended.

RESISTANCE

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become

dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

RESTRICTIONS AND PRECAUTIONS

Injury to or loss of adjacent sensitive crops, desirable trees or vegetation may result from failure to observe the following:

- Do not apply, drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants.
- Take all necessary precautions to avoid all direct or indirect contact (such as spray drift) with non-target plants or areas.
- Carefully observe all sprayer cleanup instructions both prior to and after using this product, as spray tank residue may damage crops other than wheat or barley.
- Do not apply this product by air in the state of New York.
- Do not apply this product through any irrigation system.
- Do not exceed the maximum application rate of 1 ounce per acre per crop season.

Wheat and barley may differ in their response to various herbicides. Arysta LifeScience North America, LLC recommends that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use of this product to a small area.

Under certain conditions, such as heavy rainfall, prolonged cold weather (daily high temperature less than 50°F), or wide fluctuations

in day/night temperatures prior to or soon after this product's application, temporary discoloration and/or crop injury may occur. To reduce the potential of crop injury, tank mix this product with 2,4-D (ester formulations perform best-see **TANK MIXTURES** section of this label) and apply after the crop is in the tillering stage of growth. This product should not be applied to wheat and barley that is stressed by severe weather conditions, drought (including low levels of subsoil moisture), low fertility, water saturated soil, disease, or insect damage, as crop injury may result. Risk of injury is greatest when crop is in the 2- to 5-leaf stage. Severe winter stress, drought, disease, or insect damage following application also may result in crop injury.

Do not apply to wheat or barley crops underseeded with another crop.

Dry, dusty field conditions may result in reduced control in wheel track areas.

Also observe the following:

Do not graze treated fields or feed treated forage or hay. Harvested straw may be used for bedding and/or feed.

Do not harvest wheat or barley sooner than 45 days after the last application of this product.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store product in original container only. Store in a cool, dry place.

Pesticide Disposal: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal:

For Plastic Containers: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire or other emergency contact CHEMTREC 1-800-424-9300.

Warranty and Disclaimer Statement

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Such risks may arise from weather conditions, soil factors, off-target movement, unconventional farming techniques, the presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of Arysta LifeScience North America, LLC ("Arysta"), and can cause crop injury, injury to non-target crops or plants, ineffectiveness of the product, or other unintended consequences. All such risks shall be assumed by the user or buyer.

Arysta warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to Arysta, and is subject to the inherent risks described above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARYSTA DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARYSTA, MANUFACTURER, AND SELLER DISCLAIM AND SHALL NOT BE LIABLE FOR ANY SPECIAL, INCIDENTAL, INDIRECT, OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE, HANDLING, APPLICATION, STORAGE, OR DISPOSAL OF THIS PRODUCT OR FOR DAMAGES IN THE NATURE OF PENALTIES, AND THE USER AND BUYER WAIVE ANY RIGHT THAT THEY MAY HAVE TO SUCH DAMAGES. NO AGENT, REPRESENTATIVE OR EMPLOYEE OF ARYSTA IS AUTHORIZED TO MAKE ANY WARRANTY, GUARANTEE OR REPRESENTATION BEYOND THOSE CONTAINED HEREIN OR TO MODIFY THE WARRANTIES CONTAINED HEREIN.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE TOTAL LIABILITY OF ARYSTA, MANUFACTURER, AND SELLER, SHALL BE LIMITED TO THE PURCHASE PRICE PAID, OR AT ARYSTA'S ELECTION, THE REPLACEMENT OF THE PRODUCT.

Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation.

Audit and the Audit logo are registered trademarks of Arysta LifeScience North America, LLC.

Banvel and Clarity are registered trademarks of BASF Corporation.

Buctril, Bronate and Hoelon are registered trademarks of Bayer CropScience.

Starane is a registered trademark of Dow AgroSciences, LLC.

Salvo and Sword are registered trademarks of UAP Loveland Products Inc.

RA Raindrop is a registered trademark of Delavan.

Assert, Maestro, Comet and Diablo are registered trademarks and Clash is a trademark of Nufarm Americas Inc.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

Audit 1:1

Herbicide

GROUP 2 HERBICIDE

Water Dispersible Granule
For Use on Wheat, Barley, and Fallow

ACTIVE INGREDIENTS:

Thifensulfuron-methyl

Methyl 3-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl) amino]carbonyl] amino]sulfonyl]-2-thiophenecarboxylate 25.0%

Tribenuron-methyl

Methyl 2-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)methylamino] carbonyl]amino]sulfonyl]benzoate 25.0%

OTHER INGREDIENTS: 50.0%

TOTAL: 100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

If in eyes:	Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
--------------------	---

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE: Call PROSAR at 1-866-303-6952 or 1-651-632-8946 if calling from outside the U.S.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC at 1-800-424-9300 or 1-703-527-3887 if calling from outside of the U.S.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options follow the instructions for Category A on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, made of any waterproof material such as natural rubber.
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store product in original container only. Store in a cool, dry place.

Pesticide Disposal: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal:

For Plastic Containers: Nonrefillable container. Do not reuse or refill this container. Triple rinse container(or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

See booklet for additional Precautionary Statements, Storage and Disposal and Directions for Use.

Produced by:
ARYSTA LIFE SCIENCE NORTH AMERICA, LLC

15401 Weston Parkway, Suite 150
Cary, North Carolina 27513
103356–031413

NET CONTENTS: 32 OUNCES



Arysta LifeScience

EPA Reg. No. 66330-418
EPA Est. No. 69016-MS-001
AD021413