



LICENSED

PERIOD 2014-2016 LIC. NO.

8153.4



LICENSED

PERIOD 2011-2013 LIC. NO.



LICENSED

PERIOD 2008-2010 LIC. NO.

BACTERICIDAL ACTIVITY (BACTERIA) in the presence of organic soil (5% blood serum) against *Escherichia coli* (E. coli) ATCC 11229, *Escherichia coli* 0157:H7 ATCC 43895, *Salmonella choleraesuis* ATCC 10708, *Pseudomonas aeruginosa* ATCC 15442, Vancomycin Resistant *Enterococcus Faecalis* (VRE) ATCC 51575, *Staphylococcus aureus* ATCC 6538 and Methicillin Resistant *Staphylococcus aureus* (MRSA) ATCC 33593 at 10 minutes contact time. Sanizide Plus® Wipe is bactericidal in the presence of organic soil (5% blood serum) against Vancomycin Intermediate Resistant *Staphylococcus aureus* (VISA), Clinical Isolate No. 99286, at 10 minutes contact time.

TUBERCULOCIDAL ACTIVITY in the presence of organic soil (5% blood serum) against *Mycobacterium bovis* BCG (Tuberculosis) at 20°C at 10 minutes contact time.

***VIRUCIDAL ACTIVITY** in the presence of organic soil (5% blood serum) against Human Immunodeficiency Virus Type1, Strain HTLV-IIIB (HIV-1) (associated with AIDS) at 2 minutes contact time and against Norovirus and Feline Calicivirus (FCV) at 10 minutes contact time.

Sanizide Plus® Wipe is a non-woven, disposable wipe, pre-saturated with a quaternary disinfectant cleaner.

Sanizide Plus® Wipe is intended for use in: hospitals and healthcare settings: ambulances, anesthesia, CAT labs, CCU, Central Supply, Clinics, Dental Offices, Dialysis Clinics, Doctor's offices, Emergency Rooms, E.R., Home health care, Hospices, ICU, Laboratory, Laundry rooms, Neonatal Intensive Care Units (NICU), Newborn nursery, Nursing homes, Operating rooms, Pediatrics, Pediatric Intensive Care Units (PICU), Physical therapy, Physician's offices, Ophthalmic offices, Radiology, Recovery Room, Rehabilitation, Respiratory therapy, Surgery, Transport vehicles, X-Ray

Sanizide Plus® Wipe will clean and disinfect hard, non-porous, surfaces of: bed railings, cabinets, carts, chairs, counters, exam tables, gurneys, isolettes, IV poles, stethoscopes, stretchers, tables, telephones, toilet seats, toilets, bathtubs, bathroom fixtures, shower stalls, sinks, tiles, vanity tops, hospital beds, traction devices, MRI, CAT, scales, paddles, wheelchairs, ultrasound transducers and probes.

Sanizide Plus® Wipe will clean and disinfect hard, non-porous surfaces of: grocery cart handles, grocery carts, shopping carts, grocery cart child seats.

Sanizide Plus® Wipe will clean and disinfect hard, non-porous surfaces of: changing tables, cribs, diaper pails, trashcans, garbage cans.

Sanizide Plus® Wipe will clean and disinfect hard, non-porous surfaces of: cabinets, chairs, computers, countertops, desks, desktops, doorknobs, fax machines, garbage cans, headsets, keyboards, railings, seats, sinks, tables, tabletops, telephones, trashcans.

Sanizide Plus® Wipe will clean and disinfect hard, non-porous surfaces of: gym equipment, exercise machines, wrestling mats, gymnastic equipment.

Use Sanizide Plus® Wipe to clean and disinfect non-porous personal protective safety equipment, protective headgear, hard hats, half mask respirators, full face breathing apparatus, gas masks, goggles, spectacles, face shields, silicone rubber or PVC hearing protectors and vinyl covered ear muffs. Rinse all equipment that comes in prolonged contact with skin before reuse with clean warm water (about 120°F), and allow to air dry. (Caution: Cleaning at the recommended 120°F temperature will avoid overheating and distortion of the personal safety equipment that would necessitate replacement.)

Sanizide Plus® Wipe is formulated to disinfect hard, non-porous, inanimate environmental surfaces such as floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, non-polycarbonate plastic surfaces, bathrooms, shower stalls, bathtubs, and cabinets.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE – Store in an area inaccessible to children.

CONTAINER DISPOSAL – Do not reuse empty canister. Rinse thoroughly before discarding in trash.

160ct.

**Pre-moistened
Towelettes
6.75" x6"**

Manufactured by **SAFETEC OF AMERICA, Inc.**
Buffalo, NY 14215 800-456-7077 www.safetec.com

Sanizide Plus

GERMICIDAL WIPES

CLEANS | DISINFECTS | DEODORIZES

- **TUBERCULOCIDAL**
- **BACTERICIDAL**
- **VIRUCIDAL***
- **FUNGICIDAL**

Sanizide Plus® Germicidal Wipes are effective against the following organisms:

- *Escherichia coli* (E. coli)
- Methicillin Resistant *Staphylococcus aureus* (MRSA)
- Vancomycin Intermediate Resistant *Enterococcus Faecalis* (VRE)
- Vancomycin Intermediate Resistant *Staphylococcus aureus* (VISA)
- *Mycobacterium bovis* BCG (tuberculosis)
- Human Immunodeficiency Virus Type 1*

ACTIVE INGREDIENTS

Alkyl (60% C14, 30% C16, 5% C12, 5% C18) dimethyl benzyl ammonium chlorides.....	0.105%
Alkyl (68% C12, 32% C14) dimethyl ethylbenzyl ammonium chlorides	0.105%
INERT INGREDIENTS.....	99.790%
Total	100.000%

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panel for additional precautionary statements

EPA REG. NO. 1839-174-067161
EPA EST. NO 067161-NY-001



DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is not for use on critical and semi-critical medical device surfaces.

CLEANING: Use one Sanizide Plus® wipe to completely remove all gross debris from surfaces to be treated.

DISINFECTING: To disinfect hard, non-porous surfaces, thoroughly wet the surface to be treated. Treated surface must remain visibly wet for disinfection: To disinfect against *Escherichia coli* (E. coli), *Escherichia coli* 0157:H7, *Pseudomonas aeruginosa*, *Salmonella choleraesuis* and Vancomycin Resistant *Enterococcus* (VRE), allow the treated surface to remain wet for 10 minutes. To disinfect against *Mycobacterium bovis* BCG (Tuberculosis), allow the treated surface to remain wet for 10 minutes. To disinfect against *Staphylococcus aureus* and Methicillin Resistant *Staphylococcus aureus* (MRSA), allow the treated surface to remain wet for 10 minutes. To disinfect against Vancomycin intermediate Resistant *Staphylococcus aureus* (VISA), allow the treated surface to remain wet for 10 minutes. A potable water rinse is required for surfaces in direct contact with food. For plastic and painted surfaces, spot test on an inconspicuous area before use. For heavily soiled surfaces, clean before following disinfecting instructions.

FOR ULTRASOUND TRANSDUCERS AND PROBES: Use only on external surfaces of ultrasound transducers and probes. With a clean washcloth, clean gel from external surface of the ultrasound transducer or probe to be disinfected. Then thoroughly wet probe with wipe(s). Treated surface must remain wet for 10 minutes. Use additional wipe(s) if needed to assure continuous 10 minute wet contact time. Let air dry.

***VIRUCIDAL ACTIVITY:** Sanizide Plus® Wipe is an effective virucide on hard, non-porous surfaces against: Human Immunodeficiency Virus Type 1 (HIV-1) when the treated surface is allowed to remain wet for 2 minutes, Norovirus, Feline Calicivirus when the treated surface is allowed to remain wet for 10 minutes. Use additional wipe(s) if needed to assure that the treated surface remains continuously wet for the prescribed time.

KILLS HIV-1 ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, or eye coverings.

CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. Sanizide Plus® Wipe can be used for this purpose.

CONTACT TIME: Thoroughly wet surface with a new Sanizide Plus® Wipe. Allow surface to air dry. Although efficacy at a 2 minute contact time has been shown to be adequate against HIV-1, use the recommended contact time for the disinfection of other organisms listed on this label.

DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

PRECAUTIONARY STATEMENTS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **IF IN EYES:** Hold eyes open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

Not for personal cleansing. **THIS IS NOT A BABY WIPE!** Do not flush in toilet.

Reorder no. 34823