



ferti·lome[®]

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2014-2016** LIC. NO.

8249.126

Weed-Out[®] Lawn Weed Killer

Ready To Spray

Contains **TRIMEC[®]** Herbicide

- Kills troublesome broadleaf weeds:

Dandelion, Chickweed, Clover, Henbit & many others

- Quart Covers Up To 8,000 Sq. Ft.

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

**SEE BACK PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS
AND FIRST AID**

NET CONTENTS ONE QUART (32 FL. OZS.)

ACTIVE INGREDIENTS:

Dimethylamine salt of (+)-(R)-2-
(2-methyl-4-chlorophenoxy) propionic acid..... 3.33%
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid..... 1.91%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid) ... 0.81%

OTHER INGREDIENTS:..... 93.95%
TOTAL:..... 100.00%

See back panel for substatement of ingredient statement.

FIRST AID

If In Eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.
If On Skin Or On Clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a Poison Control Center or doctor for treatment advice.
If Swallowed	<ul style="list-style-type: none"> • Call a Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a Poison Control Center or doctor. • Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment. You may also contact 1-800-255-3924 for emergency medical treatment advice.

In the event of a medical or chemical emergency contact  North America 1-800- 255-3924

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

Weed-Out® is a registered trademark of Voluntary Purchasing Groups, Inc.

Manufactured By:



BONHAM, TEXAS 75418
 EPA Reg. No. 2217-695-7401
 EPA Est. No. 7401-TX-001
 Visit Us At: www.fertilome.com



Product Questions? 855-270-4776

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear long-sleeved shirt, long pants, socks, shoes and gloves. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Use only lawn-type sprayers. Do not exceed specified dosages for any area. This product may be injurious to St. Augustinegrass, bentgrass and centipedegrass under conditions of high temperature and moisture. Under these conditions or in cases of accidental overdose some slight temporary discoloration of the St. Augustinegrass may occur. Cultivars vary in tolerance to this product. Do not apply this product to 'Floritam' St. Augustinegrass.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of run-off or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

WHERE TO USE

Ferti-lome® Weed Out Lawn Weed Killer Ready-To-Spray is a combination of three proven herbicides – 2,4-D, Mecoprop-p and Dicamba. This formulation will control a wide range of broadleaf weeds in residential and ornamental lawns. Established turf of northern (cool season) and southern (warm season) grasses may be treated. The northern (cool season) turfgrasses include Kentucky bluegrass, perennial ryegrass and the fescues. The southern (warm season)

turfgrasses include St. Augustinegrass, centipedegrass, bentgrass, bahiagrass and bermudagrass.

WHEN TO USE

For best results spraying should be done when broadleaf weeds are young and actively growing. Older weeds will be harder to kill.

1. Spray at any time during the growing season when weeds are actively growing, although spray works faster with temperatures above 70°F. To avoid damage do not apply to St. Augustinegrass or centipedegrass while emerging from winter dormancy. Damage may also occur on these grasses if temperatures drop below 45°F within 72 hours of application. Avoid spraying St. Augustinegrass when temperatures are above 90°F and under conditions of high humidity. Under these conditions or in cases of accidental overdose, some slight temporary discoloration of the St. Augustinegrass may occur.
2. Irrigate the lawn 2 or 3 days prior to spraying to ensure that weeds are growing and able to absorb the herbicide.
3. Mow 1 to 2 days before treating.
4. On new lawns, wait until the grass has reached a height of 2 inches before making a broadcast application. Wait 3 weeks before seeding grass, planting sprigs or planting stolons.
5. Do not water for 24 hours after treatment.
6. Do not mow for 48 hours after treatment.

Poor weed control may result if spray is applied during drought or just before rain.

Limitations on broadcast applications to residential turfgrass:

The maximum application rate is 5.3 fl.oz. of product per 1,000 sq.ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 10.6 fl.oz. of product per 1,000 sq.ft.

HOW TO USE

No mixing of this product with water is required. For the southern (warm season) grasses, apply the entire contents of this container (32 fluid ounces) on 8,000 square feet of turf [4 fl. ozs. per 1,000 sq. ft.]. For the northern (cool season) grasses, apply the entire contents of the container (32 fluid ounces) on 6,000 square feet of turf [5.3 fl. ozs. per 1,000 sq. ft.]. The application instructions are presented as follows:

HOW TO USE THE READY-TO-SPRAY SYSTEM

Connect

1. Shake well before using.
2. Connect sprayer to hose.
3. Turn on water.

Spray

1. To begin spraying, point nozzle in the direction you want to spray.
2. Bend small plastic tab back and Turn knob clockwise to ON position.
3. Spray evenly to the area to be treated. Refer to the sight gauge clear view strip (graduated scale) on the side of the container to determine the amount of product sprayed.

Finish

1. To stop spraying, turn knob counter clockwise to OFF position.
2. Turn off water.
3. Relieve water pressure by [bending plastic tab back and] turning knob to ON position until water slows to a drip. Then turn knob back to OFF position.
4. Disconnect sprayer from hose.

BROADLEAF WEEDS

Annual yellow sweetclover	Carolina geranium	Goldenrod	Parsley-piert
Aster	Carpetweed	Ground ivy	Parsnip
Austrian fieldcress	Catchweed bedstraw	Gumweed	Pearlwort
Bedstraw	Catnip	Hairy fleabane	Pennycress
Beggaricks	Catsear	Hawkweed	Pennywort
Betony, Florida	Chickweed, common	Healall	Peppergrass
Bindweed, field	Chickweed, mouseear	Heartleaf drymary	Pepperweed
Bird vetch	Chicory	Heathaster	Pigweed
Bitter wintercress	Cinquefoil	Hedge bindweed	Pineywoods bedstraw
Bittercress, hairy	Clover, crimson	Hedge mustard	Plains coreopsis (tickseed)
Bitterweed	Clover, hop	Hemp	Plantain
Black-eyed Susan	Clover, red	Henbit	Poison ivy
Black medic	Clover, strawberry	Hoary cress	Poison oak
Black mustard	Clover, sweet	Hoary plantain	Pokeweed
Blackseed plantain	Clover, white	Hoary vervain	Poorjoe
Blessed thistle	Cockle	Horsenettle	Prairie sunflower
Bloodflower milkweed	Cocklebur	Jimsonweed	Prickly lettuce
Blue lettuce	Common mullein	Knawel	Prickly sida
Blue vervain	Creeping jenny	Knotweed	Prostrate knotweed
Bracted plantain	Cudweed	Kochia	Prostrate pigweed
Brassbuttons	Curly dock	Lambsquarters	Prostrate spurge
Bristly oxtongue	Daisy, English	Lespedeza	Prostrate vervain
Broadleaf dock	Daisy fleabane	Mallow	Puncturevine
Broadleaf plantain	Daisy, oxeye	Matchweed	Purslane, common
Broomweed	Dandelion	Mexicanweed	Ragweed
Buckhorn	Dichondra	Milk vetch	Red sorrel
Buckhorn plantain	Dogbane	Morningglory	Redroot pigweed
Bulbous buttercup	Dogfennel	Mouseear hawkweed	Redstem filaree
Bull thistle	Dollarweed	Mugwort	Rough cinquefoil
Bullnettle	False dandelion	Musk thistle	Rough fleabane
Burclover	False flax	Mustard	Russian pigweed
Burdock	False sunflower	Narrowleaf plantain	Russian thistle
Burning nettle	Fiddleneck	Narrowleaf vetch	Scarlet pimpernel
Burweed	Florida pusley	Nettle	Scotch thistle
Buttercup	Frenchweed	Orange hawkweed	Sheep sorrel
Buttonweed	Galinsoga	Oriental cocklebur	Sheepspurse
Canada thistle	Goathead	Oxalis	Slender plantain

BROADLEAF WEEDS (continued)

Smallflower galinsoga	Tall nettle	Western clematis	Wild rape
Smooth dock	Tall vervain	Western salsify	Wild strawberry
Smooth pigweed	Tansy ragwort	White mustard	Wild sweet potato
Sorrel	Tansy mustard	Wild aster	Wild vetch
Sowthistle	Tanweed	Wild buckwheat	Woodsorrel
Spanishneedles	Thistle	Wild carrot	Woolly croton
Speedwell	Trailing crownvetch	Wild four-o'clock	Woolly morningglory
Spiny amaranth	Tumble mustard	Wild garlic	Woolly plantain
Spiny cocklebur	Tumble pigweed	Wild geranium	Wormseed
Spiny sowthistle	Velvetleaf	Wild lettuce	Yarrow
Spotted catsear	Venice mallow	Wild marigold	Yellow rocket
Spotted spurge	Virginia buttonweed	Wild mustard	Yellowflower pepperweed
Spurweed	Virginia creeper	Wild onion	
Stinging nettle	Virginia pepperweed	Wild parsnip	
Strawberry, India mock	Wavyleaf bullthistle	Wild radish	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

THIS PRODUCT CONTAINS:

0.232 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 2.75%

0.134 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 1.59%

0.057 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.68%

Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

Manufactured By:



230 FM 87 • BONHAM, TEXAS 75418

EPA Reg. No. 2217-695-7401

EPA Est. No. 7401-TX-01

Visit Us At: www.fertilome.com



7 32221 10513 3