



ferti·lome®



Department of Agriculture  
STATE OF HAWAII

**LICENSED**

PERIOD 2011-2013 LIC. NO.




Department of Agriculture  
STATE OF HAWAII

**LICENSED**

PERIOD 2014-2016 LIC. NO.

**8249.28**



# Weed-Out Broadleaf Weed Control

**CONTAINS TRIMEC® HERBICIDE**

- **Controls: Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and Many Other Broadleaf Weeds.**
- **For Use On Bahia, Bermuda, Centipede, Zoysia, Kentucky Bluegrass, Perennial Rye, Fescue and Bentgrass Lawns.**

<b>ACTIVE INGREDIENTS:</b>		<b>THIS PRODUCT CONTAINS:</b>	
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid.....	1.85%	27.4 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton or 1.37%.	
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid....	0.40%	6.6 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.33%.	
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid).....	0.18%	3.0 lbs. 3,6-dichloro-o-anisic acid equivalent per ton or 0.15%.	
<b>OTHER INGREDIENTS:</b> .....	<b>97.77%</b>	Isomer Specific by AOAC Methods.	
<b>TOTAL:</b> .....	<b>100.00%</b>	Contains the single isomer form of Mecoprop-p.	
		TRIMEC® is a registered trademark of PBI/Gordon Corporation.	

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL  
NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

**Net Weight 10 LBS. (4.54 KG)**

**Covers Up To  
5,000 Sq. Ft.**

Weed-Out Broadleaf Weed Control

**10925**

# Weed-Out Broadleaf Weed Control

**CONTAINS TRIMEC® HERBICIDE**

- **Controls: Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and Many Other Broadleaf Weeds.**
- **For Use On Bahia, Bermuda, Centipede, Zoysia, Kentucky Bluegrass, Perennial Rye, Fescue and Bentgrass Lawns.**

**ACTIVE INGREDIENTS:**

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid.....	1.65%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid....	0.40%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid).....	0.18%

**OTHER INGREDIENTS:**..... 97.77%

**TOTAL:**.....100.00%

**THIS PRODUCT CONTAINS:**

27.4 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton or 1.37%.  
6.6 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.33%.  
3.0 lbs. 3,6-dichloro-o-anisic acid equivalent per ton or 0.15%.  
Isomer Specific by AOAC Methods.  
Contains the single Isomer form of Mecoprop-p.  
TRIMEC® is a registered trademark of PBI/Gordon Corporation.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL  
NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

**Net Weight 10 LBS. (4.54 KG)**

**Covers Up To  
5,000 Sq. Ft.**



**READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.

**FIRST AID**

<b>If In Eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a Poison Control Center or doctor for treatment advice.</li> </ul>
<b>If On Skin Or Clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>• Call a Poison Control Center or doctor for treatment advice.</li> </ul>
<b>If Swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a Poison Control Center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a Poison Control Center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>

**Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment. In the event of a medical or chemical emergency, contact ChemTel™ North America 1-800-255-3924 or Worldwide Intl. + 01- 813-248-0585.**



**USER SAFETY RECOMMENDATIONS**

**User should:**  
 -Wash hands before eating, drinking, chewing gum or tobacco, or using the toilet.  
 -Remove clothing/PPE immediately if pesticide gets inside. Wash thoroughly and put on clean clothing.  
 -Remove clothing/PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in contamination of drinking water or groundwater.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

**STORAGE AND DISPOSAL**

**STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.  
**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.  
**DISPOSAL: If Empty:** Do not reuse bag. Place in trash or offer for recycling if available. **If Partly Filled:** Call your Local Solid Waste Agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**USE PRECAUTIONS**

Do not apply this product to St. Augustinegrass, Carpetgrass, Dichondra, Ornamentals, or Lawns containing desirable Clovers. Avoid spreading granules onto Landscape, Plantings, Flowers, Vegetables or Ornamentals. Do not spread over exposed roots of Trees and Ornamentals. Do not exceed specified dosages for any area; be particularly careful within drip-line of Tree and other Ornamental Species. Treated areas may be seeded 3 to 4 weeks after application. Application of this product to newly seeded areas should be delayed until the third mowing.

**WHERE TO USE**

**ferti-lome® WEED-OUT® BROADLEAF WEED CONTROL** is a **Trimec®** complex, a combination of three proven Weed Killers – 2, 4-D, Mecoprop and Dicamba. Together, they control a wide range of Lawn Weeds, including Dandelion, Chickweed, Knotweed, Plantains, Henbit, Spurge and many others found in Home Lawns. Established turf of Kentucky Bluegrass, Bentgrass, Bahiagrass, Centipedegrass, Fescues, Bermudagrass, Zoysiagrass, and Perennial Ryegrass may be treated.

**HOW MUCH TO APPLY**

## HOW MUCH TO APPLY

**For Kentucky Bluegrass, Bentgrass, Bahiagrass, Centipedegrass, Fescues, Perennial Ryegrass, Bermudagrass, and Zoysiagrass:** Apply 2.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of Weeds or for adverse seasonal conditions. The second application of 2.0 pounds of product per 1,000 square feet should be used 30 days after the first application. The maximum application rate to turf is 1.2 pounds 2,4-D acid equivalent per acre per application per site.

The maximum number of broadcast applications per treatment site is 2 per year.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when Weeds are young and actively growing, preferably in the morning when dew is on the Grass. If Grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget Plants.
4. Do not wash from Weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

## APPLICATION EQUIPMENT AND INSTRUCTIONS

This product can be applied with a drop or broadcast (rotary) spreader. For best results, apply this product evenly and uniformly over the turfgrass (lawn) area avoiding streaking, skips or overlaps.

Before each application, determine the correct amount (lbs.) of product to place in the spreader hopper for the area to be treated. For example, if the label states to apply 4 lbs. per 1000 sq. ft. and your area is 2000 sq. ft, then place 8 lbs. of product in the hopper and set the spreader setting at 4 1/4 (per table below) if using an Accugreen drop spreader.

A more uniform application can be made by spreading one-half of the recommended amount of this product over the turfgrass area and then applying the remaining one-half of this product at a right angle to the previous application direction.

**NOTE: The spreader settings listed below are approximate. The speed at which you walk and the working condition of the spreader will affect the rate of application.**

SPREADER	SPREADER SETTINGS	
	Lbs. of product per 1,000 sq. ft.	
Scotts/Republic Accugreen (Drop)	4 1/4	7
Scotts/Republic Speedy Green (Broadcast)	3 3/4	6
Ferti-lome/Earthway Ev-N-Spred (Broadcast)	14	16

## THE FOLLOWING BROADLEAF WEEDS ARE CONTROLLED BY THIS PRODUCT:

Annual Yellow Sweetclover	Burclover	Daisy, Fleabane	Knawel	Poison Ivy	Sorrel	White Mustard
Austrian Fieldcress	Burdock	Daisy, Oxeye	Knotweed	Poison Oak	Sowthistle	Wild Aster
Bedstraw	Burning Nettle	Dandelion	Kochia	Pokeweed	Spanishneedles	Wild Buckwheat
Beggarticks	Burweed	Dichondra	Lambsquarters	Poorjoe	Speedwell	Wild Carrot
Betony, Florida	Buttercup	Dogfennel	Lespedeza	Prairie Sunflower	Spiny Amaranth	Wild Four-O'Clock
Bindweed, Field	Buttonweed	Dollarweed	Mallow	Prickly Lettuce	Spiny Cocklebur	Wild Garlic
Bird Vetch	Canada Thistle	False Dandelion	Matchweed	Prickly Sida	Spiny Sowthistle	Wild Geranium
Bitter Wintercress	Carolina Geranium	Falseflax	Mexicanweed	Prostrate Knotweed	Spotted Catsear	Wild Lettuce
Bittercress, Hairy	Carpetweed	False Sunflower	Milk Vetch	Prostrate Pigweed	Spotted Spurge	Wild Marigold
Bitterweed	Catchweed Bedstraw	Fiddleneck	Morningglory	Prostrate Spurge	Spurweed	Wild Mustard
Black-eyed Susan	Catnip	Florida Pusley	Mouseear Hawkweed	Prostrate Vervain	Stinging Nettle	Wild Onion
Black Medic	Catsear	Frenchweed	Mugwort	Puncturevine	Strawberry, India Mock	Wild Parsnip
Black Mustard	Chickweed, Common	Galinsoga	Musk Thistle	Purslane, Common	Tall Nettle	Wild Radish
Blackseed Plantain	Chickweed, Mouseear	Goldenrod	Mustard	Ragweed	Tall Vervain	Wild Rape
Blessed Thistle	Chicory	Ground Ivy	Narrowleaf Plantain	Red Sorrel	Tansy Ragwort	Wild Strawberry
Bloodflower Milkweed	Cinquefoil	Gumweed	Narrowleaf Vetch	Redroot Pigweed	Tansy Mustard	Wild Sweet Potato
Blue Lettuce	Clover, Crimson	Hairy Fleabane	Nettle	Redstem Filaree	Tanweed	Wild Vetch
Blue Vervain	Clover, Hop	Hawkweed	Orange Hawkweed	Rough Cinquefoil	Thistle	Woodsorrel
Bracted Plantain	Clover, Red	Healall	Oriental Cocklebur	Rough Fleabane	Trailing Crownvetch	Woolly Croton
Bristly Oxtongue	Clover, Strawberry	Heartleaf Drymar	Oxalis	Russian Pigweed	Tumble Mustard	Woolly Morningglory
Broadleaf Dock	Clover, Sweet	Heath Aster	Parsley-piert	Russian Thistle	Tumble Pigweed	Woolly Plantain
Broadleaf Plantain	Clover, White	Hedge Bindweed	Parsnip	Scarlet Pimpernel	Velvetleaf	Wormseed
Broomweed	Cockle	Hedge Mustard	Pearlwort	Scotch Thistle	Venice Mallow	Yarrow
Buckhorn	Cocklebur	Henbit	Pennycress	Sheep Sorrel	Virginia Buttonweed	Yellow Rocket
Buckhorn Plantain	Common Mullein	Hoary Cress	Pepperweed	Sheperdspurse	Virginia Creeper	Yellowflower Pepperweed
Bulbous Buttercup	Creeping Jenny	Hoary Plantain	Pigweed	Slender Plantain	Virginia Pepperweed	and other Broadleaf Weeds
Bull Thistle	Cudweed	Hoary Vervain	Pineywoods Bedstraw	Smallflower Galinsoga	Wavyleaf Bulthistle	
Bullnettle	Curly Dock	Horsenettle	Plains Coreopsis	Smooth Dock	Western Clematis	
	Daisy, English	Jimsonweed	Plantain	Smooth Pigweed	Western Salsify	

Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

Manufactured By:



230 FM 87 • BONHAM, TEXAS 75418

EPA Reg. No.: 2217-657-7401

EPA Est. No.: 7401-TX-1

Visit Us At: [www.fertilome.com](http://www.fertilome.com)

RV072310

