

LICENSED

PERIOD **2011-2013** LIC. NO.

PERIOD **2014-2016** LIC. NO.

8249.2

PERIOD **2008-2010** LIC. NO.

**KEEP OUT OF REACH OF CHILDREN.
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

WARNING: Causes substantial, but temporary eye injury. Do not get in eyes. Wear goggles or safety glasses. Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not allow people or pets to contact treated areas until sprays are dry.

FIRST AID

IF IN EYES: Hold eyelids open and flush eyes with a steady, gentle stream of water. Get medical attention.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Move victim to fresh air.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: If ingested, induce emesis or lavage stomach. Treat symptomatically.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment wash water.

Notice: Buyers assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith. Voluntary Purchasing Groups, Inc. warrants this product to conform to the chemical description on the label and for the purposes stated on this label. The limit of any liability incurred shall be the purchase price paid by the User or Buyer.

▲ PEEL FROM CORNER OF BOOKLET

32M-6-112M



ferti·lome[®]



\$11⁹⁸

Liquid Systemic Fungicide





- Prevents And Stops Major Diseases On Roses, Flowers, Lawns, Trees And Shrubs.
- Absorbed Into The Plant – Can't Be Washed Off By Rain Or Sprinklers After Spray Has Dried.
- Works Up To 4 Weeks On Certain Diseases.
- Encourages Increased Turf Rooting.
- Improves Color In Woody Shrubs.

ACTIVE INGREDIENT:	
Propiconazole: (CAS No. 60207-90-1)	1.55%
INERT INGREDIENTS	98.45%
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN.

WARNING

See Additional Precautionary Statements On Back Panel.
NET CONTENTS 8 Fl. Ozs.

LAWNS

- 1) Apply after mowing or allow sprayed area to completely dry before mowing.
- 2) Measure the length and width of area to be treated before application. Length x width = square feet. For example, an area 25 ft. long and 20 ft. wide is 500 sq. ft.

ESTABLISHED LAWNS

DISEASE	HOW MUCH TO APPLY	COMMENTS
Brown Patch	5 fl. oz. in 1 gallon of water per 250 sq. ft. of lawn	If needed, repeat treatment every 10 days. First application should be in May or June before disease is present.
Dollar Spot, Powdery Mildew, Rust, Anthracnose	3 fl. oz. in 1 gallon of water per 250 sq. ft. of lawn	Apply every 14 days.
Red Thread, Dichondra Rust on Dichondra Lawns	5 fl. oz. in 1 gallon of water per 250 sq. ft. of lawn	If needed, repeat every 21 days when conditions are favorable for disease to occur.
Stripe Smut	3 fl. oz. in 1 gallon of water per 250 sq. ft. of lawn	Apply in Fall after grass is dormant or in early Spring before grass starts to grow.

NEW SEEDINGS

HOW MUCH TO APPLY	COMMENTS
5 fl. oz. in 2 gallons of water per 500 sq. ft. of lawn	Apply at the 2 to 3-leaf stage of growth for faster root development and top growth.

4

TREES AND SHRUBS

- General Directions:** 1) Spray just until product runs off leaves.
2) Allow the spray to dry before watering plants.

PLANT	DISEASE	HOW MUCH TO APPLY	COMMENTS
Crab Apple <i>*Note: Do not use fruit from treated Crab Apples for food purposes.</i>	Scab, Rust	½ Tablespoon (¼ fl. oz.) per gallon of water	Apply when buds just begin to break open. For Scab, apply every 14 days. For Rust, make 3 applications, beginning when buds just begin to break open. Apply at 7-day intervals.
	Powdery Mildew	1½ Tablespoons (¾ fl. oz.) per gallon of water	Apply every 14 days.
Crape Myrtle	Powdery Mildew	3 Tablespoons (1½ fl. oz.) per gallon of water	Apply every 21 days.
Dogwood	Anthracnose	1 Tablespoon (½ fl. oz.) per gallon of water	Apply every 14–28 days.
Douglas Fir, Shasta Fir	Needle Rust	3 Tablespoons (1½ fl. oz.) per gallon of water	For Douglas Fir, apply every 21 days. For Shasta Fir, apply in May–June before disease occurs.
Jeffrey Pine	Tip Blight	3 Tablespoons (1½ fl. oz.) per gallon of water	Make the first application in mid–late Winter (Nov.–Jan.) Apply every 2 months up to 3 times.
Oak, Red Tip Photinia	Leaf Spot	3 Tablespoons (1½ fl. oz.) per gallon of water	Apply every 14 days.

5

NON-BEARING FRUIT AND NUT TREES

- General Directions:** 1) Non-bearing Fruit and Nut Trees are those trees that will not produce fruit for at least one year after application of this product.
2) Apply product to the point just when it begins to run off of the leaves.

PLANT	DISEASE	HOW MUCH TO APPLY	COMMENTS
Apple	Scab	½ Tablespoon (¼ fl. oz.) per gallon of water	Start applications when leaves start growing in the Spring and spray every 14 days.
	Rust	½ Tablespoon (¼ fl. oz.) per gallon of water	Make 3 applications beginning when buds just begin to break open. Apply at 7-day intervals.
	Powdery Mildew	1 Tablespoon (½ fl. oz.) per gallon of water	Start applications in early Spring and spray every 14 days.
Cherry	Powdery Mildew, Leaf Spot	1 Tablespoon (½ fl. oz.) per gallon of water	Start applications in early Spring and spray every 21 days.
Citrus	Greasy Spot	4 Tablespoons (2 fl. oz.) per gallon of water	Apply June–August every 21 days.

6

NON-BEARING FRUIT AND NUT TREES (CONTINUED)

- General Directions:** 1) Non-bearing Fruit and Nut Trees are those trees that will not produce fruit for at least one year after application of this product.
2) Apply product to the point just when it begins to run off of the leaves.

PLANT	DISEASE	HOW MUCH TO APPLY	COMMENTS
Pecan	Scab, Powdery Mildew, Downy Spot, Zonate Leafspot, Fungal Leaf Scorch, Brown Spot	2 Tablespoons (1 fl. oz.) per gallon of water	Start applications at bud break and make 3 applications at 14-day intervals.
Walnut	Anthracnose	2 Tablespoons (1 fl. oz.) per gallon of water	Apply every every 14 days.

7

Distributed By:
VOLUNTARY PURCHASING GROUPS, INC.
 P.O. BOX 460
 BONHAM, TEXAS 75418
 EPA Reg. No. 100-773-7401
 EPA Est. No. 7401-TX-1
 U.S. Patent No.: 4,079,062
 NCPPL 773A-LIC 0200



DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

STOP! Read all instructions inside before using. Failure to follow the Directions for Use and Precautions on this label may result in poor disease control or injury to desirable plants.

STOP! DO NOT ALLOW PEOPLE OR PETS TO CONTACT TREATED AREAS UNTIL SPRAY HAS DRIED.

STORAGE AND DISPOSAL

Storage: Store in a cool, dry place, away from heat or open flame.
Disposal: Do not reuse container. Securely wrap container in several layers of newspaper and discard in trash. For minor spills, follow all precautions on this label and clean up immediately. In the event of an emergency, call 1-800-424-9300, day or night.

1) WHAT IS THIS PRODUCT FOR?

This water-based Systemic Fungicide effectively prevents and stops major Diseases on Roses, Flowers, Lawns, Trees & Shrubs.

2) HOW DO YOU USE IT?

USE TIPS:

- This product is highly effective when it is used to prevent Diseases or applied at the first sign of Disease. Consider treating early in the season. This product will not eliminate Disease already present.
- When used as directed, this product is safe to plants and controls the Diseases listed. Plants not listed may be injured by this product and Diseases not listed may not be controlled.
- Do not apply this product to African Violets, Begonias, Boston Ferns, or Geraniums.
- If you need more information about Lawn and Ornamental Diseases, contact your local County Extension Service.

3) HOW TO MIX:

Shake well before using. Apply only with a pump-up pressure sprayer, concentrate plus water-style, or a dial-style hose-end sprayer. Use of hose-end sprayers which have only an on/off switch or those which are continuously filled with water during the application is not recommended.

NOTE: ½ fl. oz. = 1 Tablespoon; 1 fl. oz. = 2 Tablespoons

Food utensils, such as measuring cups and spoons, must not be used for food purposes after use in measuring pesticides.

PUMP-UP SPRAYER:

1. Fill with required amount of water.
2. Add the amount of concentrate specified in the charts that follow.
3. Shake well before application.

CONCENTRATE PLUS WATER-STYLE SPRAYER:

1. From the following chart, determine the amount of concentrate required.
2. Pour concentrate into sprayer jar to the required "fl. oz." level.
3. Then add the required amount of water to the "gals." level.
4. Replace sprayer top on jar and shake well before spraying.

DIAL-STYLE HOSE-END SPRAYER:

1. Set sprayer on the "fl. oz./gallon" level specified in the following chart.
2. Use only enough product for the area to be treated.
3. Close sprayer, attach to hose and apply product to lawn.
 (For example: to treat a 1,000 sq. ft. lawn for powdery mildew, pour 12 fl. oz. into sprayer jar. Set sprayer dial at 3 fl. oz. per gallon.)

4) HOW TO APPLY: ROSES AND FLOWERS

General Directions: 1) Apply product to the point just when it begins to run off of the leaves.
 2) Allow spray to completely dry before watering.

PLANT	DISEASE	HOW MUCH TO APPLY	COMMENTS
Roses	Powdery Mildew	1 Tablespoon (½ fl. oz.) per gallon of water	Thorough coverage needed for best results. Begin application at bud break. Treat every 10 days.
	Black Spot, Rust	2 Tablespoons (1 fl. oz.) per gallon of water	Treat every 10 days. Note: May not provide 100% control of Black Spot under severe conditions.
Calendula	Powdery Mildew	1 Tablespoon (½ fl. oz.) per gallon of water	Apply every 30 days.
Chrysanthemums	Ray Blight	2 Tablespoons (1 fl. oz.) per gallon of water	Apply every 7 days.
Delphinium	Powdery Mildew	1 Tablespoon (½ fl. oz.) per gallon of water	Apply every 14 days.
Snapdragons	Rust	2 Tablespoons (1 fl. oz.) per gallon of water	Apply every 10 days.
Sweet William	Leafspot	1 Tablespoon (½ fl. oz.) per gallon of water	Apply every 14 days.