



ferti·lome®



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO.

8249.30

DISC. Weed-Out®
Lawn Weed Killer

Contains TRIMEC® Herbicide

- **Kills troublesome broadleaf weeds:
Dandelion, Chickweed, Clover, Henbit & many others**

**KEEP OUT OF REACH
OF CHILDREN**

CAUTION

**SEE BACK PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS
AND FIRST AID**

Net Contents One Quart (32 FL. OZS.)

ACTIVE INGREDIENTS:

Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid.....	5.45%
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid.....	5.88%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid).....	1.21%
OTHER INGREDIENTS:.....	87.46%
TOTAL:.....	100.00%

**See back panel for substatement of
ingredient statement.**

ferti•lome® WeedOut Lawn Weed Killer

20M-39-0612 FH

FIRST AID

If In Eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a Poison Control Center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none">• Call a Poison Control Center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a Poison Control Center or doctor.• Do not give anything by mouth to an unconscious person.
If On Skin Or Clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a Poison Control Center or doctor for treatment advice.
If Inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a Poison Control Center or doctor for further treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment.

In the event of a medical or chemical emergency contact  **North America 1-800-255-3924 or Worldwide Intl. + 01-813-248-0585**

Manufactured By:



230 FM 87
BONHAM, TEXAS 75418
EPA Reg. No. 2217-597-7401
EPA Est. No. 7401-TX-001
Visit Us At: www.fertilome.com



PEEL FROM CORNER OF BOOKLET



STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear long-sleeved shirt, long pants, socks, shoes and gloves. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drain systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift.

- 
- Keep people and pets out of the area during application.
 - Do not allow people or pets to enter the treated area until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container and place in locked storage area when not using.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down and indoor or outdoor drain.

WHEN TO USE

Spray at any time when Weeds are growing. For best results, apply in Spring when Weeds are small and growing rapidly. Apply in Fall to remove late germinating Weeds.

HOW TO USE

Hose-End Sprayer: Use 1 gallon of water for every 250 square feet to be sprayed. For concentrate sprayers, add only the required amount of **ferti-lome® Weed Out Lawn Weed Killer** listed in the rate chart depending on type of Grass and number of square feet to be treated. Do not add water to concentrate sprayers. For dilution sprayers, add the required amount of **ferti-lome® Weed Out Lawn Weed Killer** and fill sprayer with water to proper level. Spray uniformly.

Pressure-Sprayer: Use 1/2 gallon of water for every 250 square feet to be sprayed. Add the required amount of **ferti-lome® Weed Out Lawn Weed Killer** listed in the rate chart for the type of Grass to be treated and fill

sprayer with water to the proper level. Adjust spray nozzle to deliver coarse spray. Spray uniformly.

Spot-Treatment: If hardy Weeds persist in Sensitive Grasses such as St. Augustine, Centipedegrass, Bermudagrass, Buffalograss or Bentgrass, spot treat individual Weeds using 1 teaspoon of **ferti-lome® Weed Out Lawn Weed Killer** in 1 quart of water (equivalent to 4 teaspoons per gallon of water.) Minimize contact of Grass.

RATES OF ferti-lome® Weed Out Lawn Weed Killer FOR VARIOUS LAWN GRASSES					
Gallons Of Water Spray		Area to Treat (Sq. Ft.)	St. Augustine- grass, Centipedegrass (fl. oz.)	Bermudagrass, Buffalograss Bentgrass (fl. oz)	Bluegrass Zoysia- grass, Fescue (fl. oz)
Hose End Sprayer	Pressure Sprayer				
1	1/2	250	1/2	2/3	2/3 to 1
4	2	1,000	2	2 2/3	2 2/3 to 4
8	4	2,000	4	5 1/3	5 1/3 to 8
20	10	5,000	10	13 1/3	13 1/3 to 20
32	16	8,000	16	21 1/3	21 1/3 to 32

EQUAL MEASURES

- 1 fluid ounce (fl. oz.) = 2 tablespoons (Tbs.) or 6 teaspoons (tsp.)
- 2 fluid ounces = 4 tablespoons 4 fluid ounces = 1/2 cup
- 8 fluid ounces = 1 cup 16 fluid ounces = 1 pint (pt.)
- 32 fluid ounces = 1 quart (qt.)

HOW TO USE IN THE LAWN WEED KILLER APPLICATOR

1. Connect sprayer nozzle to garden hose; remove safety cap from container and insert sprayer unit, securing tightly.
2. Turn on water; place finger over hole on right side of nozzle and product will automatically be dispersed.

3. Spray evenly and uniformly over area requiring application. (Check application guide to determine the correct area to be covered per ounce.)

APPLICATION GUIDE			
Fluid Ounces	Recommended Coverage Area Per Ounce		
	Sensitive Grasses Such As Centipedegrass And St. Augustinegrass (sq. ft.)	Bermudagrass, Bentgrass, And Buffalograss (sq. ft.)	Bluegrass, Fescue, And Zoysiagrass (sq. ft.)
1	500	375	250
2	1,000	750	500
3	1,500	1,125	750
4	2,000	1,500	1,000
5	2,500	1,875	1,250
6	3,000	2,250	1,500
7	3,500	2,525	1,750
8	4,000	3,000	2,000
9	4,500	3,375	2,250
10	5,000	3,750	2,500
11	5,500	4,125	2,750
12	6,000	4,500	3,000
13	6,500	4,875	3,250
14	7,000	5,250	3,500
15	7,500	5,625	3,750
16	8,000	6,000	4,000

Use Precautions: Do not apply this product through any type of irrigation system. Use only lawn sprayer that delivers coarse spray to reduce drift. Avoid spray drift to Vegetables, Flowers, Ornamental Plants. Do not spray roots of Trees and Ornamentals. Do not exceed specified dosages for any area; be particularly careful within dripline of Trees and other Ornamental species. Do not spray Carpetgrass, Dichondra, or Lawn where desirable Clovers are present. Do not apply this product to "Floritam" St. Augustine Grass in Florida. Do not apply to newly seeded Lawn Grasses until well established. Do not spray when air temperatures exceed 85°F. Seed can be



sown 3 to 4 weeks after application. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides. Failure to follow above precautions may result in injury.

Other Use Precautions:

- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays and oceans. Do not apply when windy. To prevent run-off, do not over water the treat area(s) to the point of run-off or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

PRODUCT INFORMATION

Ferti-lome® Weed Out Lawn Weed Killer controls troublesome common Lawn Weeds, including Poison Ivy, in both Northern and Southern Grasses.

- BROADLEAF WEEDS -

Annual Yellow Sweetclover	Bull Thistle	Cudweed
Aster	Bullnettle	Curly Dock
Austrian Fieldcress	Burclover	Daisy, English
Bedstraw	Burdock	Daisy Fleabane
Beggarticks	Burning Nettle	Daisy, Oxeye
Betony, Florida	Burweed	Dandelion
Bindweed, Field	Buttercup	Dichondra
Bird Vetch	Buttonweed	Dogfennel
Bitter Wintercress	Canada Thistle	Dollarweed
Bittercress, Hairy	Carolina Geranium	False Dandelion
Bitterweed	Carpetweed	Falseflax
Black-Eyed Susan	Catchweed Bedstraw	False Sunflower
Black Medic	Catnip	Fiddleneck
Black Mustard	Catsear	Florida Pusley
Blackseed Plantain	Chickweed, Common	Frenchweed
Blessed Thistle	Chickweed, Mouseear	Galinsoga
Bloodflower Milkweed	Chicory	Goldenrod
Blue Lettuce	Cinquefoil	Ground Ivy
Blue Vervain	Clover, Crimson	Gumweed
Bracted Plantain	Clover, Hop	Hairy Fleabane
Bristly Oxtongue	Clover, Red	Hawkweed
Broadleaf Dock	Clover, Strawberry	Healall
Broadleaf Plantain	Clover, Sweet	Heartleaf Drymary
Broomweed	Clover, White	Heathaster
Buckhorn	Cockle	Hedge Bindweek
Buckhorn Plantain	Cocklebur	Hedge Mustard
Bulbous Buttercup	Common Mullein	Henbit
	Creeping Jenny	Hoary Cress

- BROADLEAF WEEDS - *(Continued)*

Hoary Plantain	Pennywort	Scotch Thistle
Hoary Vervain	Pepperweed	Sheep Sorrel
Horsenettle	Pigweed	Sherherdspurse
Jimsonweed	Pineywoods Bedstraw	Slender Plantain
Knawel	Plains Coreopsis	Smallflower Galinsoga
Knotweed	Plantain	Smooth Dock
Kochia	Poison Ivy	Smooth Pigweed
Lambsquarters	Poison Oak	Sorrel
Lespedeza	Pokeweed	Sowthistle
Mallow	Poorjoe	Spanishneedles
Matchweed	Prairie Sunflower	Speedwell
Mexicanweed	Prickly Lettuce	Spiny Amaranth
Milk Vetch	Prickly Sida	Spiny Cocklebur
Morningglory	Prostrate Knotweed	Spiny Sowthistle
Mouseear Hawkweed	Prostrate Pigweed	Spotted Catsear
Mugwort	Prostrate Spurge	Spotted Spurge
Musk Thistle	Prostrate Vervain	Spurweed
Mustard	Puncturevine	Stinging Nettle
Narrowleaf Plantain	Purslane, Common	Strawberry, India
Narrowleaf Vetch	Ragweed	Mock
Nettle	Red Sorrel	Tall Nettle
Orange Hawkweed	Redroot Pigweed	Tall Vervain
Oriental Cocklebur	Redstem Filaree	Tansy Ragwort
Oxalis	Rough Cinquefoil	Tansy Mustard
Parsley-Piert	Rough Fleabane	Tanweed
Parsnip	Russian Pigweed	Thistle
Pearlwort	Russian Thistle	Trailing Crownvetch
Pennycress	Scarlet Pimpernel	Tumble Mustard

- BROADLEAF WEEDS - (Continued)

Tumble Pigweed	Wild Four-O'Clock	Woodsorrel
Velvetleaf	Wild Garlic	Woolly Croton
Venice Mallow	Wild Geranium	Woolly Morningglory
Virginia Buttonweed	Wild Lettuce	Woolly Plantain
Virginia Creeper	Wild Marigold	Wormseed
Virginia Pepperweed	Wild Mustard	Yarrow
Wavyleaf Bullthistle	Wild Onion	Yellow Rocket
Western Clematis	Wild Parsnip	Yellowflower
Western Salsify	Wild Radish	Pepperweed
White Mustard	Wild Rape	and other Broadleaf
Wild Aster	Wild Strawberry	Weeds
Wild Buckwheat	Wild Sweet Potato	
Wild Carrot	Wild Vetch	

LIMITED WARRANTY AND DISCLAIMER

The Manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This Limited Warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the Manufacturer or the Seller. Any damages arising from a breach of the Manufacturer's Warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss

or profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the Seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

This page left blank intentionally.

THIS PRODUCT CONTAINS:

0.39 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 4.50%.

0.42 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 4.88%.

0.09 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 1.00%.

Isomer Specific by AOAC Methods.

Contains the single isomer form of Mecoprop-p.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

Manufactured By:



230 FM 87 • BONHAM, TX 75418

EPA Reg. No. 2217-597-7401

EPA Est. No. 7401-TX-001

Visit Us At: www.fertilome.com



Get the free mobil app at
<http://gettag.mobi>