



ferti·lome®



BEFORE

AFTER

WEED FREE ZONE

Ready To Use

- Rainfast in as little as 3 hours
- Visible results in just 24 hours
- Proven cool-weather performance.

**KEEP OUT OF REACH
OF CHILDREN**

CAUTION

SEE BACK PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

TRIMEC® is a registered trademark
of PBI/Gordon Corporation

ACTIVE INGREDIENTS:

MCPA, 2-ethylhexyl ester	0.3370%
Mecoprop-p acid	0.0660%
Dicamba acid	0.0180%
Carfentrazone-ethyl	0.0023%
INERT INGREDIENTS:	99.5767%
TOTAL:	100.0000%

THIS PRODUCT CONTAINS:

2-methyl-4-chlorophenoxyacetic acid
equivalent, 0.216%

Net Contents One Quart (32 FL. OZS.)



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2013-2015 LIC. NO. 8249.55

FIRST AID

If In Eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after the first five minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If On Skin or Clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a Poison Control Center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none"> • Call a Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a Poison Control Center or doctor. • Do not give anything to an unconscious person.

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

In the event of a medical or chemical emergency, contact  **ChemTel**
North America 1-800-255-3924 or Worldwide Intl. + 01-813-248-0585

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Harmful if absorbed through the skin. Avoid contact with eyes, skin, or clothing. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

This pesticide may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment wash water or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding application when rainfall is forecasted to occur within 48 hours.



230 FM 87 • BONHAM, TEXAS 75418
EPA Reg. No. 2217-865-7401
EPA Est. No. 7401-TX-1
Visit Us At: www.fertilome.com
10M-1-0512 RTUH
(RV011510)



PEEL FROM CORNER OF BOOKLET





READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Precautions and Restrictions:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Not for use on turf being grown for sale or other commercial use on sod, or for commercial seed production, or for research purposes.

PRODUCT FACTS:

ferti-lome® WEED FREE ZONE Ready-To-Use contains four active ingredients including carfentrazone-ethyl that broaden the spectrum of Weed Control.

ferti-lome® WEED FREE ZONE Ready-To-Use offers these advantages:

- Excellent postemergent activity with proven performance for Broadleaf Weed control in turfgrass.
- Superior cool weather performance.
- Rapid and effective Weed control for common and troublesome Weed Species in Turfgrass, e.g., Spurge, Pennywort (Dollarweed), Dandelion, and White Clover.
- Injury to Weeds can be noticed within hours of application and Plant death can occur within 7 - 14 days.

WHERE TO USE

On Lawns: Use to spot treat Broadleaf Weeds commonly found in Cool Season and Warm Season Grasses such as:

Kentucky Bluegrass
Perennial Ryegrass
Tall Fescue
Red or Fine Leaf Fescues
Mixtures of Cool Season Species
Common Bermudagrass

Hybrid Bermudagrass
St. Augustinegrass
(excluding 'Floratam')
Zoysiagrass

Prohibitions:

- Do not apply this product to Bentgrass, Carpetgrass, Dichondra, Legumes and Lawns where desirable clovers are present.
- Do not apply to 'Floratam' St. Augustinegrass grown in Florida.
- Do not allow spray to contact Vegetables, Flowers or Ornamentals as injury may result.
- Do not apply more than 1.9 gallons of product per 1,000 sq. ft. per year (1.5 lb. a. i. MCPA per acre per year).
- Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

APPLICATION SCHEDULES (WHEN TO APPLY)

For best results, treat Weeds in the Spring and the Fall when Weeds are actively growing. However, this product is effective any time during the Growing Season.

- Do not apply when air temperatures are above 85°F as Lawn injury can occur.
- Hard-to-kill Weeds may require a repeat application in 3 weeks.
- Do not water Lawn within 12 hours of application. If rainfall occurs within 12 hours after application, retreatment may be necessary.
- Do not apply to newly seeded Lawns until after the second mowing.

- Delay application to newly sodded, plugged, or sprigged Lawns until new Grass has been mowed four times.
- Treated areas may be reseeded 2 weeks after application.
- Applications to Dormant Bermudagrass, Zoysiagrass, and St. Augustinegrass may be used to control actively growing Broadleaf Weeds.

HOW TO USE

ferti-lome® WEED FREE ZONE Ready-To-Use is applied directly from the container fitted with the trigger sprayer. Shake the container well. Direct the spray to the target Weeds to provide the uniform coverage of the upper leaf surfaces. Apply to the point of runoff only. Excessive spraying can injure Lawns.

BROADLEAF WEEDS CONTROLLED

ferti-lome® WEED FREE ZONE Ready-To-Use will control or suppress the following Broadleaf Weeds:

BROADLEAF WEEDS

Aster, White Heath & White Prairie	Compassplant
Bedstraw	Curly Dock
Beggarweed, Creeping	Dandelion
Bindweed	Dayflower
Black Medic	Deadnettle
Broadleaf Plantain	Dock
Buckhorn Plantain	Dogfennel
Bull Thistle	English Daisy
Burclover	False Dandelion (*Spotted Catsear
Burdock, Common	& Common Catsear)Field
Buttercup, Creeping	Bindweed
Carpetweed	(*Morningglory
Chickweed, Common	& Creeping Jenny)
Chicory	Field Oxeye-Daisy
Cinquefoil	(*Creeping Oxeye)
Clover	Filaree, Whitestem & Redstem

*Synonyms

BROADLEAF WEEDS (Continued)

Florida Pusley	Prostrate Knotweed
Ground Ivy	(* Knotweed)
Groundsel	Puncturevine
Hawkweed	Purple Cudweed
Healall	Purslane
Henbit	Ragweed
Innocence (Blue-Eyed Mary)	Redweed
Lambsquarters	Red Sorrel (*Sheep Sorrel)
Lawn Burweed	Shepherdspurse
Lespedeza, Common	Spotted Spurge
Mallow, Common	Spurge, Prostrate
Matchweed	Thistle
Mouseear Chickweed	Veronica (*Corn Speedwell)
Old World Diamond Flower	Virginia Buttonweed
Oxalis	White Clover (*Dutch Clover
(*Yellow Woodsorrel & Creeping Woodsorrel)	Honeysuckle Clover, White Trefoil, & Purplewort)
Parsley-Piert	Wild Carrot
Pennsylvania Smartweed	Wild Garlic
Pennywort (*Dollarweed)	Wild Geranium
Pepperweed	Wild Lettuce
Pigweed	Wild Mustard
Pineappleweed	Wild Onion
Plantain	Wild Strawberry
Poison Ivy	Wild Violet
Poison Oak	Yarrow
	Yellow Rocket

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container in an area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

The Manufacturer warrants only that the chemical composition of this product conforms to the Ingredient Statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions For Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This Limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the Manufacturer or the Seller. Any damages arising from a breach of the Manufacturer's Warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss or profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the Seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.



**Weed Free Zone
Ready-To-Use**

Manufactured By:



230 FM 87

BONHAM, TEXAS 75418

EPA Reg. No. 2217-865-7401

EPA Est. No. 7401-TX-1

Visit Us At: www.fertilome.com

(RV011510)

