





ferti·lome®




Dollarweed




Florida Pusley




Foxtail




Dandelion



Crabgrass



Chickweed



Clover



Henbit



Goosegrass



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2014-2016 LIC. NO.

8249.70

A-Vert Plus Lawn Food

LAWN WEED PREVENTER AND FERTILIZER

• Covers Up To 2,000 Sq. Ft.

• One Easy Application Stops Weeds Before They Start!

• A Fertilizer Containing Gallery* And Team*,
Preemergence Herbicides For Control Of Annual Grasses And
Broadleaf Weeds In Established Home Lawns.

ACTIVE INGREDIENT:

Isoxaben: N-(3-(1-ethyl-1-methylpropyl)-5-isoxazolyl)-2,6-dimethoxybenzamide and isomers	0.29%
Trifluralin: α,α,α-trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine	0.27%
Benfen: N-butyl-N-ethyl-α,α,α-trifluoro-2,6-dinitro-p-toluidine	0.53%
OTHER INGREDIENTS:	98.91%
TOTAL:	100.00%

Contains 0.1308 pounds of active ingredient per 12 pound bag.

*TRADEMARK of Dow AgroSciences LLC

KEEP OUT OF REACH OF CHILDREN

CAUTION

Refer To Back Panel For Precautionary Statements And Directions For Use Including Storage And Disposal.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label. If terms are unacceptable, return at once unopened.

Agricultural Chemical:

Do Not Ship or Store with Food, Feeds, Drugs or Clothing.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.

Net Weight 12 LBS. (5.44 KG)

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or inhaled, Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals, Avoid breathing dust.

USER SAFETY RECOMMENDATIONS

Users should: • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

FIRST AID

If Swallowed: • Call a poison control center or doctor immediately for treatment advice. • If person is unconscious, do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.

If Inhaled: • Move person to fresh air. • Call a poison control center or doctor for further treatment advice. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In the event of a medical or chemical emergency contact **ChemTel**  North America 1-800-255-3924 or Worldwide Int. + 01-813-248-6535

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose as waste.

Container Reuse: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying **ferti-lome® A-VERT PLUS LAWN FOOD**.

Application Restrictions: Do not apply **ferti-lome® A-VERT PLUS LAWN FOOD** in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Entry Restrictions: Do not allow people or pets to enter the treated areas until dusts have settled. If watering-in is required after the application, do not enter or allow others to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

HOW THIS PRODUCT WORKS

ferti-lome® A-VERT PLUS LAWN FOOD is a granular fertilizer containing Gallery® and Team™, pre-emergence herbicides for application to established home lawns for control of certain broadleaf weeds and annual grasses. **ferti-lome® A-VERT PLUS LAWN FOOD** also feeds the lawn with a fertilizer designed to provide prolonged, even graining without excess top growth.

ferti-lome® A-VERT PLUS LAWN FOOD will not control existing weeds. Existing weeds should be controlled with a foliarly applied post-emergence herbicide either before or after application of **ferti-lome® A-VERT PLUS LAWN FOOD**.

GENERAL USE PRECAUTIONS AND RESTRICTIONS

- Do not apply to dichondra.
- Do not apply to newly sprigged or hydromulched areas of Bermudagrass, St. Augustinegrass, centipede grass or zoysiagrass until turfs are well established.
- Any cultural practices that disturb the soil, such as aeration or verticuting, should be done prior to application of **ferti-lome® A-VERT PLUS LAWN FOOD**.
- Do not make more than 2 applications of **ferti-lome® A-VERT PLUS LAWN FOOD** per year.
- Apply **ferti-lome® A-VERT PLUS LAWN FOOD** only to established turf. Apply to newly seeded turf (including overseeded turf) only after seedlings are established and well rooted (generally after 3 mowings).
- Reseeding:** Do not reseed treated areas for at least 8 weeks following application of **ferti-lome® A-VERT PLUS LAWN FOOD**.
- Do not apply to non-turf areas such as flower beds or gardens. Take precautions to prevent inadvertent application from rotary-type applicators to such areas. The herbicides in **ferti-lome® A-VERT PLUS LAWN FOOD** prevent seed emergence

- and may injure bedding plants.
- ferti-lome® A-VERT PLUS LAWN FOOD** may injure turf that is weakened due to severe winter weather or is not well established.
- Do not apply **ferti-lome® A-VERT PLUS LAWN FOOD** in the spring and summer to tall fescue turfs as well as the previous fall.
- Do not apply in the spring to Bermudagrass that has been overseeded with winter annual grasses since thinning may occur.
- Do not overdose or overlap during application in such a way as to apply in excess of labeled rates as turf injury may occur.
- Apply **ferti-lome® A-VERT PLUS LAWN FOOD** only to the grass species listed on this label. The user assumes responsibility for any plant damage or other liability resulting from use of **ferti-lome® A-VERT PLUS LAWN FOOD** on turf species not recommended on this label.
- Maximum Single Application Rate:** 7.2 lbs. of **ferti-lome® A-VERT PLUS LAWN FOOD** per 1,000 sq. ft.
- Maximum Yearly Rate:** 14.4 lb of **ferti-lome® A-VERT PLUS LAWN FOOD** per 1,000 sq. ft.
- Minimum Re-Treatment Interval:** Cool season turf – 56 to 70 days; warm season turf – 70 to 84 days.

THIS PRODUCT IS RECOMMENDED FOR USE ON THE FOLLOWING ESTABLISHED TURFGRASS SPECIES:

ESTABLISHED COOL SEASON GRASSES				ESTABLISHED WARM SEASON GRASSES		
Bentgrass, Colonial Bentgrass, Creeping	Bluegrass, Kentucky Fescue, Chewings	Fescue, Creeping Red Fescue, Sheep	Fescue, Tall Ryegrass, Perennial	Bahiagrass Bermudagrass	Centipede grass Fescue, Tall	St. Augustinegrass Zoysiagrass

WEEDS CONTROLLED

WEEDS CONTROLLED WHEN APPLIED PRIOR TO WEED GERMINATION						
Annual Bluegrass Barnyardgrass Barnyardgrass Buttercup Carpetweed	Celery, Wild Chickweed, Common Chickweed, Mouseear Crabgrass (Large Crabgrass)	(Smooth Crabgrass) Field Sandbur (Field Sandspur) Footspursley Foxtails	(Bottlegrass) (Bristlegrass) (Giant Foxtail) (Green Foxtail) (Piggrass)	(Robust Foxtail) Geranium, Carolina Goosegrass Herbit Knobweed, Prostrate	Mallow, Little Parsley - Piert Pennywort, Lawn Pineappleweed Speedwell, Thymeleaf	Spurge, Prostrate Spurge, Spotted Spurry, Corn Woodsorrel, Creeping Woodsorrel, Yellow

WEEDS CONTROLLED BY A REPEAT APPLICATION 8 WEEKS AFTER THE FIRST APPLICATION WHEN APPLIED PRIOR TO WEED GERMINATION

Clover spp' Clover, White'	Dandelion' Groundsel, common	Medic, Black Plantain, Broadleaf	Plantain, Buckhorn Plantain, slender
-------------------------------	---------------------------------	-------------------------------------	---

From Seed Only

HOW TO APPLY

Apply **ferti-lome® A-VERT PLUS LAWN FOOD** using a drop or rotary-type spreader designed to apply granular fertilizer. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution. Optimum weed control performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours. A single rainfall or sprinkler irrigation of 0.5 inch or more is required to activate **ferti-lome® A-VERT PLUS LAWN FOOD**.

achieved by spreading half the required amount of product over the area and then applying the remaining half in swaths at right angles to the first. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution. Optimum weed control performance will be obtained when an application is followed by sprinkler irrigation or rainfall within a few hours. A single rainfall or sprinkler irrigation of 0.5 inch or more is required to activate **ferti-lome® A-VERT PLUS LAWN FOOD**.

APPLICATION EQUIPMENT AND INSTRUCTIONS

SPREADER	Spreader Settings		This product can be applied with a drop or broadcast (rotary) spreader. For best results, apply this product evenly and uniformly over the turfgrass (lawn) area avoiding streaking, skips or overlaps.
	Lbs. of product per 1,000 sq. ft.		
	5.75 LBS.	7.2 LBS.	
Scotts/Republic Accugreen (Drop)	5-3/4	7-1/4	Before each application, determine the correct amount (lbs.) of product to place in the spreader hopper for the area to be treated. For example, if the label states to apply 5.75 lbs. per 1,000 sq. ft. and your area to treat is 2,000 sq. ft. then place 11.5 lbs. of product in the hopper and set the spreader setting at 5-3/4 (per table below) if using an Accugreen drop spreader.
Scotts/Republic Speedy Green (Broadcast)	5-1/2	7	
Ferti-lome/Earthway Ev-N-Spred (Broadcast)	18	20	A more uniform application can be made by spreading one-half of the recommended amount of this product over the turfgrass area and then applying the remaining one-half of this product at a right angle to the previous application direction.

NOTE: The spreader settings listed below are approximate. The speed at which you walk and the working condition of the spreader will affect the rate of application.

HOW MUCH TO APPLY

Apply **ferti-lome® A-VERT PLUS LAWN FOOD** to cool season turfgrass at a rate of 5.75 lb. per 1,000 sq. ft. and to warm season turfgrass at a rate of 7.2 lbs. per 1,000 sq. ft.

WHEN TO APPLY

ferti-lome® A-VERT PLUS LAWN FOOD can be applied at any time of the spring or fall. The type of weeds controlled will vary depending upon time of application so other weed control products may be needed to provide a total annual weed control program.

Early Spring (before Forsythia blooms): If applied in early spring, **ferti-lome® A-VERT PLUS LAWN FOOD** will prevent crabgrasses and other annual grasses (including Poa annual) and broadleaf weeds, such as dandelion, clover, plantain, and spurge, that normally germinate in the spring and summer. When used in this manner, **ferti-lome® A-VERT PLUS LAWN FOOD** may be substituted for a multi-step weed and feed program. For your area, consult the local extension service for optimum timing for control of crabgrass.

Mid-Late Spring (after Forsythia blooms, before Memorial Day): **ferti-lome® A-VERT PLUS LAWN FOOD** will prevent annual grasses (including Poa annual) and broadleaf weeds such as dandelion, clover, plantain, and spurge, that have yet to germinate in the spring and summer, ensuring season-long weed control.

Early fall (before average night temperatures fall below 60°F): **ferti-lome® A-VERT PLUS LAWN FOOD** will prevent establishment of winter annual grasses (including Poa annual) and broadleaf weeds, such as chickweed, herb, and speedwell, that normally germinate in the fall and winter and broadleaf weeds that germinate the following spring season. A supplemental application of a crabgrass pre-emergence herbicide may be needed in the spring for spring and summer weed control.

Late fall: **ferti-lome® A-VERT PLUS LAWN FOOD** will prevent establishment of broadleaf weeds, such as chickweed, herb, and speedwell, that may germinate over winter and broadleaf weeds that germinate the following spring season. A supplemental application of a crabgrass pre-emergence herbicide may be needed in the spring for spring and summer weed control.

TERMS AND CONDITIONS OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise,

use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SELLER MAKES NO OTHER EXPRESS OR IMPLIED

WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.),

abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer.

LIMITATION OF REMEDIES

To the extent permitted by law, the exclusive remedy for losses or damages resulting from handling or use of **ferti-lome® A-VERT PLUS LAWN FOOD** (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

Seller shall not be liable for losses or damages resulting from handling or use of **ferti-lome® A-VERT PLUS LAWN FOOD** unless Seller is promptly notified of such loss or damage in writing. To the extent consistent with applicable law, in no case shall Seller be liable for consequential or incidental damages or losses.

- Refund of purchase price paid by buyer or user for product bought, or
- Replacement of amount of product used.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

18-0-12 GUARANTEED ANALYSIS

Total Nitrogen (N)	18%
3.3% Ammoniacal Nitrogen	
14.7% Urea Nitrogen	
Soluble Potash (K ₂ O)	12%
Boron (B)	0.02%
Copper (Cu)	0.05%
0.05% Water Soluble Copper (Cu)	
Iron (Fe)	0.10%
0.10% Water Soluble Iron (Fe)	
Manganese (Mn)	0.05%
0.05% Water Soluble Manganese (Mn)	
Zinc (Zn)	0.05%
0.05% Water Soluble Zinc (Zn)	

Derived From: Urea, Ammonium Sulfate, Muriate of Potash, Sodium Borate, Copper Sulfate, Ferrous Sulfate, Manganese Sulfate, Zinc Sulfate.

Manufactured By:



230 FM 87
BONHAM, TEXAS 75418
EPA Reg. No. 62719-192-7401 • EPA Est. No. 7401-TX-1
Visit Us At: www.fertilome.com

Information regarding the contents and levels of metals in this product is available on the internet at: <http://aaipco.org/metals.htm>



15M-1-111-1