

Note to Customer: Weiman Products adds a revision code to all artwork items for tracking purposes. As the customer & controller of your own artwork, we ask that you either denote a specific area for this revision code and/or ask Weiman for the specific revision code that is to be added & place on artwork.



LICENSED

PERIOD 2013-2015 LIC. NO. **8262.5**

18.5 inches Width

5 inches Height

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply this undiluted product by sponge, rag, cloth, mop, spray or immersion to wet all surfaces thoroughly.

When used as directed this product effectively kills the following microorganisms at room temperature with a: 10-second sanitization** contact time against *Enterobacter aerogenes**, *Staphylococcus aureus****, and 2-minute disinfectant contact time against *Acinetobacter baumannii*, Avian Influenza A (H₃N₂) Reassortant Virus*, *Escherichia coli* (ESBL) (*E. coli*), Hepatitis B Virus* (HBV), Hepatitis C Virus* (HCV), Herpes Simplex Virus* (HSV-2), Human Immunodeficiency Virus* (HIV-1) (associated with AIDS), Influenza A Virus*, *Klebsiella pneumoniae*-Carbapenem-resistant (KPC), Methicillin Resistant *Staphylococcus aureus* (MRSA), *Mycobacterium bovis* BCG (tuberculosis) (TB), *Pseudomonas aeruginosa*, Rotavirus*, *Salmonella enterica*, *Staphylococcus aureus*, Swine Influenza A Virus* (H₁N₁) (formerly known as 'swine flu'), *Trichophyton mentagrophytes* (Ringworm) and Vancomycin-Resistant *Enterococcus faecalis* (VRE). For disinfection of Avian Influenza A, use this product as a spray only for surfaces that are not conducive to treatment by immersion or excess liquid.

Use this product to clean, disinfect and sanitize** soft and hard non-food contact surfaces, non-porous inanimate surfaces such as those made of acrylic, metals (aluminum, brass, stainless steel, chrome), glass, plastics (polycarbonate, polypropylene, polystyrene and polyvinylchloride), clear plastics, Plexiglas®, vinyl and leather surfaces as well as laminated and painted surfaces.

This product is effective in dental laboratories, dental offices, oral health care facilities, orthodontic offices and other similar establishments.

May be used on items such as dental impressions***, dental trays, dental chair surfaces, dental equipment surfaces, patient care fixtures, computer keyboards, computer screens, light lens covers, suction units, toothbrushes** and other similar items. (**Rinse with potable water before use.)

SURFACES

For surface cleaning: Prior to surface disinfection or sanitization**, pre-clean surfaces, making sure to remove all gross filth and heavy soil, by saturating surfaces with this product then immediately wipe surface clean using a fresh paper or cloth towel. Repeat, if necessary, until surfaces are visibly clean. Discard paper towels or launder cloth towels before reusing.

For cleaning walls and floors: In a clean bucket, dilute 2 to 4 oz. of this product per gallon of warm, clean, potable water. Thoroughly damp-mop using a clean mop, cloth towel, or sponge. Allow to air dry.

For surface sanitization: Apply this product to clean surfaces by sponge, cloth, mop, spray or immersion to wet all surfaces thoroughly. Allow to remain visibly wet for 10 seconds at room temperature, then wipe surfaces using a clean mop, paper or cloth towel, or rinse surface using potable water or wipe surface dry or allow to air dry. Discard or launder spent materials before reuse.

For surface disinfection: Apply this product to pre-cleaned surfaces by sponge, cloth, mop, spray or immersion to wet all surfaces thoroughly. Allow to remain wet for 2 minutes, then wipe dry using a clean mop, paper or cloth towel, or rinse surface using potable water and wipe surface dry or allow to air dry. Discard or launder spent materials before reuse.

INSTRUMENTS / DEVICES

For use as a pre-cleaning instrument/device soak: Fill appropriately sized container with a sufficient amount of this product to allow for complete submersion of items. Place items into solution and allow to soak for 2 minutes. Be certain to flush lumens of hollow devices with this product. Remove items and rinse. For critical and semi-critical items, follow with the appropriate cleaning and disinfection or sterilization process. Change solution as needed, when the solution becomes diluted or visibly soiled.

For use as a pre-cleaning spray: Place item into a suitable container. Thoroughly spray item with this product until drenched, thoroughly rinse. Follow with an appropriate terminal cleaning and sterilization/disinfection process.

For use as a non-critical medical item disinfection immersion solution: Thoroughly pre-clean items being sure to remove all gross filth and heavy soil, then rinse and dry. Allowing for complete submersion of item(s), place pre-cleaned item(s) into an appropriate container that has been filled with this product. Flush lumens of hollow devices with this product. Allow item(s) to remain submerged for 2 minutes. Remove, rinse with clean, potable water and dry. Change solution daily or when it becomes visibly soiled or discolored.

For cleaning and disinfecting dental impressions: Apply this product until thoroughly drenched, then rinse with potable water and wipe or air dry. Reapply product to thoroughly wet the surfaces and allow to remain wet for 2 minutes. Rinse with potable water and wipe or air dry.

For use as a protective eye wear/loupes cleaner/disinfectant/sanitizer:** Thoroughly wet entire surface and wipe clean using a soft cloth. Reapply and allow to remain wet for 10 seconds for sanitization** or 2 minutes for disinfection and re-wipe clean using a fresh cloth. This product should not be used on contact lenses.

Opti-Cide³

Virucidal*
Fungicidal
Bactericidal
Tuberculocidal

Hospital & Dental
Hard and Soft Surface Sanitizer & Disinfectant / Cleaner**
Ready to Use

KEEP OUT OF REACH OF CHILDREN
CAUTION

See side panels for First Aid and additional Precautionary Statements.

ACTIVE INGREDIENTS:

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈)	
dimethyl benzyl ammonium chloride	0.154%
n-Alkyl (68% C ₁₂ , 32% C ₁₄)	
dimethyl ethylbenzyl ammonium chloride	0.154%
Isopropanol	21.000%
OTHER INGREDIENTS	78.692%
TOTAL	100.000%

EPA REG. NO 70144-1-51003 EPA EST. 1130-IL-1



Made in the U.S.A.
Manufactured for:
BIOTROL™

Net Contents:
1 Gallon (3.78 L)
Order No. D0CP04-128

HBV, HCV AND HIV-1 PRECAUTIONS

This product kills **HBV, HCV and HIV-1** on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in health care or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects can be associated with the potential for transmission of HBV, HCV and/or the human immunodeficiency virus Type-1 (HIV-1) associated with AIDS. **Special instructions for cleaning and decontamination against HBV, HCV and HIV-1 of surfaces/objects soiled with blood/body fluids:** **Personal Protection:** Specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable latex gloves, gowns, masks and eye coverings. **Cleaning Procedure:** Blood and body fluids must be thoroughly cleaned from surfaces/objects before disinfection. **Contact Time:** Allow the surface to remain wet for 2 minutes. **Infectious Materials Disposal:** Cleaning materials used that may contain blood/body fluids must be autoclaved and/or disposed of in accordance with local regulations for infectious materials disposal.

This product is not to be used as a terminal sterilant or high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes, but which does not ordinarily penetrate the blood barrier or otherwise enter sterile areas of the body. This product may be used to pre-clean or decontaminate critical and semi-critical medical devices prior to sterilization or high-level disinfection.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container in a cool, well-ventilated area inaccessible to children and pets. **Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. **Container Disposal:** Non-refillable container. Do not reuse or refill this container. Triple rinse as follows: Fill container 1/4 full with water and reclose the container. Agitate vigorously and dispose of rinsate in accordance with Pesticide Disposal Instructions. Repeat two more times. Offer for recycling if available. Puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

FIRST AID

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-255-3924 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: May be harmful if inhaled or absorbed through skin. May cause moderate eye irritation. Avoid breathing vapors and contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

Manufactured in the USA for:
Biotrol
13705 Shoreline Court East
Earth City, MO 63045
(800) 822-8550

10700189C
D0CP04128-81482C-0613



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply this undiluted product by sponge, rag, cloth, mop, spray or immersion to wet all surfaces thoroughly.

When used as directed this product effectively kills the following microorganisms at room temperature with a: 10-second sanitization** contact time against *Enterobacter aerogenes*** , *Staphylococcus aureus*** ; and 2-minute disinfectant contact time against *Acinetobacter baumannii*, Avian Influenza A (H₃N₂) Reassortant Virus*, *Escherichia coli* (ESBL) (*E. coli*), Hepatitis B Virus* (HBV), Hepatitis C Virus* (HCV), Herpes Simplex Virus* (HSV-2), Human Immunodeficiency Virus* (HIV-1) (associated with AIDS), Influenza A Virus*, *Klebsiella pneumoniae*-Carbapenem-resistant (KPC), Methicillin Resistant *Staphylococcus aureus* (MRSA), *Mycobacterium bovis* BCG (tuberculosis) (TB), *Pseudomonas aeruginosa*, Rotavirus*, *Salmonella enterica*, *Staphylococcus aureus*, Swine Influenza A Virus* (H₁N₁) (formerly known as 'swine flu'), *Trichophyton mentagrophytes* (Ringworm) and Vancomycin-Resistant *Enterococcus faecalis* (VRE). For disinfection of Avian Influenza A, use this product as a spray only for surfaces that are not conducive to treatment by immersion or excess liquid.

Use this product to clean, disinfect and sanitize** soft and hard non-food contact surfaces, non-porous inanimate surfaces such as those made of acrylic, metals (aluminum, brass, stainless steel, chrome), glass, plastics (polycarbonate, polypropylene, polystyrene and polyvinylchloride), clear plastics, Plexiglas®, vinyl and leather surfaces as well as laminated and painted surfaces.

This product is effective in dental laboratories, dental offices, oral health care facilities, orthodontic offices and other similar establishments.

May be used on items such as dental impressions***, dental trays, dental chair surfaces, dental equipment surfaces, patient care fixtures, computer keyboards, computer screens, light lens covers, suction units, toothbrushes*** and other similar items. (***Rinse with potable water before use.)

SURFACES

For surface cleaning: Prior to surface disinfection or sanitization**, pre-clean surfaces, making sure to remove all gross filth and heavy soil, by saturating surfaces with this product then immediately wipe surface clean using a fresh paper or cloth towel. Repeat, if necessary, until surfaces are visibly clean. Discard paper towels or launder cloth towels before reusing.

For cleaning walls and floors: In a clean bucket, dilute 2 to 4 oz. of this product per gallon of warm, clean, potable water. Thoroughly damp-mop using a clean mop, cloth towel, or sponge. Allow to air dry.

For surface sanitization: Apply this product to clean surfaces by sponge, cloth, mop, spray or immersion to wet all surfaces thoroughly. Allow to remain visibly wet for 10 seconds at room temperature, then wipe surfaces using a clean mop, paper or cloth towel, or rinse surface using potable water or wipe surface dry or allow to air dry. Discard or launder spent materials before reuse.

For surface disinfection: Apply this product to pre-cleaned surfaces by sponge, cloth, mop, spray or immersion to wet all surfaces thoroughly. Allow to remain wet for 2 minutes, then wipe dry using a clean mop, paper or cloth towel, or rinse surface using potable water and wipe surface dry or allow to air dry. Discard or launder spent materials before reuse.

INSTRUMENTS / DEVICES

For use as a pre-cleaning instrument/device soak: Fill appropriately sized container with a sufficient amount of this product to allow for complete submersion of items. Place items into solution and allow to soak for 2 minutes. Be certain to flush lumens of hollow devices with this product. Remove items and rinse. For critical and semi-critical items, follow with the appropriate cleaning and disinfection or sterilization process. Change solution as needed, when the solution becomes diluted or visibly soiled.

For use as a pre-cleaning spray: Place item into a suitable container. Thoroughly spray item with this product until drenched, thoroughly rinse. Follow with an appropriate terminal cleaning and sterilization/disinfection process.

For use as a non-critical medical item disinfection immersion solution: Thoroughly pre-clean items being sure to remove all gross filth and heavy soil, then rinse and dry. Allowing for complete submersion of item(s), place pre-cleaned item(s) into an appropriate container that has been filled with this product. Flush lumens of hollow devices with this product. Allow item(s) to remain submerged for 2 minutes. Remove, rinse with clean, potable water and dry. Change solution daily or when it becomes visibly soiled or discolored.

For cleaning and disinfecting dental impressions: Apply this product until thoroughly drenched, then rinse with potable water and wipe or air dry. Reapply product to thoroughly wet the surfaces and allow to remain wet for 2 minutes. Rinse with potable water and wipe or air dry.

For use as a protective eye wear/loupes cleaner/disinfectant/sanitizer:** Thoroughly wet entire surface and wipe clean using a soft cloth. Reapply and allow to remain wet for 10 seconds for sanitization** or 2 minutes for disinfection and re-wipe clean using a fresh cloth. This product should not be used on contact lenses.

Opti-Cide³®

Virucidal*

Fungicidal

Bactericidal

Tuberculocidal

Hospital & Dental

Hard and Soft Surface Sanitizer & Disinfectant / Cleaner**

Ready to Use

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See side panels for First Aid and
additional Precautionary Statements.

ACTIVE INGREDIENTS:

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈)	
dimethyl benzyl ammonium chloride.....	0.154%
n-Alkyl (68% C ₁₂ , 32% C ₁₄)	
dimethyl ethylbenzyl ammonium chloride	0.154%
Isopropanol	21.000%

OTHER INGREDIENTS 78.692%

TOTAL 100.000%

EPA REG. NO 70144-1-51003

EPA EST. 1130-IL-1



Made in the U.S.A.
Manufactured for:

BIOTROL™

Net Contents:

1 Gallon (3.78 L)

Order No. DOCP04-128

HBV, HCV AND HIV-1 PRECAUTIONS

This product kills **HBV, HCV and HIV-1** on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in health care or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects can be associated with the potential for transmission of HBV, HCV and/or the human immunodeficiency virus Type-1 (HIV-1) associated with AIDS. **Special instructions for cleaning and decontamination against HBV, HCV and HIV-1 of surfaces/objects soiled with blood/body fluids:** **Personal Protection:** Specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable latex gloves, gowns, masks and eye coverings. **Cleaning Procedure:** Blood and body fluids must be thoroughly cleaned from surfaces/objects before disinfection. **Contact Time:** Allow the surface to remain wet for 2 minutes. **Infectious Materials Disposal:** Cleaning materials used that may contain blood/body fluids must be autoclaved and/or disposed of in accordance with local regulations for infectious materials disposal.

This product is not to be used as a terminal sterilant or high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes, but which does not ordinarily penetrate the blood barrier or otherwise enter sterile areas of the body. This product may be used to pre-clean or decontaminate critical and semi-critical medical devices prior to sterilization or high-level disinfection.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container in a cool, well-ventilated area inaccessible to children and pets. **Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. **Container Disposal:** Non-refillable container. Do not reuse or refill this container. Triple rinse as follows: Fill container 1/4 full with water and reclose the container. Agitate vigorously and dispose of rinsate in accordance with Pesticide Disposal instructions. Repeat two more times. Offer for recycling if available. Puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

FIRST AID

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-255-3924 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: May be harmful if inhaled or absorbed through skin. May cause moderate eye irritation. Avoid breathing vapors and contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

Manufactured in the USA for:

Biotrol
13705 Shoreline Court East
Earth City, MO 63045
(800) 822-8550

10700189C
DOCP04128-81482C-0613

