

# DISC.

Department of Agriculture  
STATE OF HAWAII

Department of Agriculture  
STATE OF HAWAII

LICENSED

LICENSED

PERIOD 2011-2013 LIC. NO.

PERIOD 2014-2016 LIC. NO.

# Pasada® 1.6F

8275.20

## FLOWABLE INSECTICIDE

For control of certain insects infesting various crops.

**GROUP 4A INSECTICIDE**

ACTIVE INGREDIENT	% BY WT.
Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]- <i>N</i> -nitro-2-imidazolidinimine	17.4%
<b>INERT INGREDIENTS:</b>	<b>82.6%</b>
<b>TOTAL</b>	<b>100.0%</b>

Contains 1.6 pounds of active ingredient per gallon.

**SHAKE WELL BEFORE USING**

**STOP - Read the label before use**  
**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

FIRST AID	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>

EPA Reg. No. 264-763-66222

FIRST AID	
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For medical emergencies involving this product, contact Prosar at 1-877-250-9291.</p>	
<p><b>NOTE TO PHYSICIAN:</b> No specific antidote is available. Treat the patient symptomatically.</p>	

For additional precautionary, handling, and use statements, see inside of this booklet.



Manufactured for:  
**Makhteshim Agan**  
of North America, Inc.  
4515 Falls of Neuse Road  
Suite 300  
Raleigh, NC 27609

EPA 081307/  
Notif 092109/Rev B  
US04321921C

**Net Contents: 1 Gallon**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC), or Viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENGINEERING CONTROLS STATEMENTS**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates.

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**OBSERVE THE FOLLOWING PRECAUTIONS WHEN MIXING AND APPLYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, ESTUARIES, AND COMMERCIAL FISH FARM PONDS.**

## SPRAY DRIFT MANAGEMENT

The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

### Mixing and Loading Requirements

To avoid potential contamination of groundwater, the use of a properly designed and maintained containment pad for mixing and loading of any pesticide into application equipment is recommended. If containment pad is not used, maintain a minimum distance of 25 feet between mixing and loading areas and potential surface to groundwater conduits such as field sumps, uncased well heads, sinkholes, or field drains.

### For Aerial Applications

The spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used and must not exceed 75% of the wing span or rotor diameter.

### Importance of Droplet Size

An important factor influencing drift is droplet size. Small droplets (<150 - 200 microns) drift to a greater extent than large droplets. Within typical equipment specifications, applications should be made to deliver the largest droplet spectrum that provides sufficient control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the airstream as much as possible, and by avoiding excessive spray boom pressure.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

### Wind Speed Restrictions

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors including droplet size, canopy, and equipment specifications determine drift potential at any given wind speed. Do not apply when winds are greater than 15 mph and avoid gusty and windless conditions. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

### Restrictions During Temperature Inversions

Do not make aerial or ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.

### Airblast (Air-Assist) Specific Recommendations for Tree Crops and Vineyards

Airblast sprayers carry droplets into the canopy of trees/vines via a radially or laterally directed airstream. The following specific drift management practices should be followed:

- Adjust deflectors and aiming devices so that spray is only directed into the canopy.
- Block off upward-pointed nozzles when there is no overhanging canopy.
- Use only enough air volume to penetrate the canopy and provide good coverage.
- Do not allow the spray to go beyond the edge of the cultivated area (i.e., turn off sprayer when turning at end rows).
- Only spray inward toward the orchard or vineyard for applications to the outside rows.

### **No-Spray Zone Requirements for Foliar Applications**

Do not apply by ground within 25 feet or by air within 150 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.

### **Runoff Management**

Do not cultivate within 10 feet of the aquatic areas to allow growth of a vegetative filter strip. When used on erodible soils, Best Management Practices for minimizing runoff should be employed. Consult your local Natural Resources Conservation Service for recommendations in your use area.

### **Endangered Species Notice**

Under the Endangered Species Act, it is a Federal offense to use any pesticide in a manner that results in the death of a member of an endangered species. Consult your local county bulletin, County Extension Agent, or Pesticide State Lead Agency for information concerning endangered species in your area.

### **Resistance Management**

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area.

Pasada 1.6F contains a Group 4A insecticide. Insect biotypes with acquired or inherent tolerance to Group 4A products may eventually dominate the insect population if Group 4A products are used repeatedly as the predominant method of control for targeted species. This may eventually result in partial or total loss of control of those species by Pasada 1.6F and to other Group 4A products.

The active ingredient in Pasada 1.6F is a member of the neonicotinoid chemical class. Avoid using a block of more than three consecutive applications of Pasada 1.6F and/or other Group 4A products having the same or similar mode of action. Following a neonicotinoid block of treatments, Makhteshim Agan of North America, Inc. strongly encourages the rotation to a block of applications with effective products of a different mode before using additional applications of neonicotinoid products. Using a block rotation or windowed approach along with other IPM practices is considered an effective use strategy for preventing or delaying an insect pest's ability to develop resistance to this class of chemistry.

Foliar applications of Pasada 1.6F or other Group 4A products from the neonicotinoid chemical class should not be used on crops previously treated with a long-residual, soil-applied product from the neonicotinoid chemical class. Other Group 4A neonicotinoid products used as foliar treatments include: Actara, Assail, CALYPSO, Centric, Clutch, Couraze, Gallant, Impulse, Intruder, LEVERAGE, Nuprid, Provado, TRIMAX PRO, and Venom.

Other Group 4A neonicotinoid products used as soil/seed treatment include: ADMIRE PRO, Advise, Alias, Belay, Couraze, Cruiser, GAUCHO, Macho, Macho Max, Nuprid, Platinum, Venom, and Widow.

Contact your local extension specialist, certified crop advisor, and/or product manufacturer for additional insect resistance management recommendations. Also, for more information on Insect Resistance Management (IRM), visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://irac-online.org/>.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC), or Viton
- Shoes plus socks

#### APPLICATION DIRECTIONS

##### **Do Not Apply Pasada 1.6F In Enclosed Structures Such As Greenhouses Or Planthouses.**

Pasada 1.6F should be applied as a directed or broadcast foliar spray. Thorough coverage of foliage is necessary without runoff for optimum insecticidal efficacy. Use adequate spray volumes, properly calibrated application equipment, and spray adjuvant if necessary to obtain thorough coverage. Failure to provide adequate coverage and retention of Pasada 1.6F on leaves and fruit may result in loss of insect control or delay in onset of activity. Pasada 1.6F may be applied with properly calibrated ground or aerial application equipment. Minimum recommended spray volumes, unless otherwise specified on crop-specific **Recommended Application** sections, are 10 gallons/acre by ground application and 5 gallons/acre through aerial equipment. Pasada 1.6F may also be applied by overhead chemigation (see additional **CHEMIGATION DIRECTIONS FOR USE** section below) if allowed in crop-specific **Recommended Application** section.

Pasada 1.6F use on crops grown for production of true seed intended for private or commercial planting is generally not recommended but may be allowed under state-specific supplemental labeling. As with any insecticide, care should be taken to minimize exposure of Pasada 1.6F to honey bees and other pollinators. Use of Pasada 1.6F on crops requiring bee pollination should be avoided during bloom and a minimum of 10 days prior to bloom. Additional information on Pasada 1.6F uses for these crops and other questions may be obtained from the Cooperative Extension Service, PCAs, consultants, or local Makhteshim Agan of North America, Inc. representatives.

Do not apply more than 0.5 lb. active ingredient per acre per crop season regardless of formulation or method of application, unless specified within a crop-specific **Recommended Application** section for a given crop.

### **MIXING INSTRUCTIONS**

To prepare the application mixture, add a portion of the required amount of water to the spray tank, and with agitation, add Pasada 1.6F. Complete filling tank with balance of water needed. Maintain sufficient agitation during both mixing and application. Pasada 1.6F may also be used with other pesticides and/or fertilizer solutions. Please see **Compatibility Note** below. When tank mixtures of Pasada 1.6F and other pesticides are involved, prepare the tank mixture as recommended above and follow suggested **Mixing Order** below.

#### **Mixing Order**

When pesticide mixtures are needed, add wettable powders or wettable granules first, Pasada 1.6F or other flowables second, and emulsifiable concentrates last. Ensure good agitation as each component is added. Do not add an additional component until the previous is thoroughly mixed. If a fertilizer solution is added, a fertilizer pesticide compatibility agent may be needed. Maintain constant agitation during both mixing and application to ensure uniformity of spray mixture.

#### **Compatibility Note**

Test compatibility of the intended tank mixture before adding Pasada 1.6F to the spray or mix tank. Add proportionate amounts of each ingredient in the appropriate order to a pint or quart jar, cap, shake for 5 minutes, and let set for 5 minutes. Poor mixing or formation of precipitates that do not readily re-disperse indicates an incompatible mixture that should not be used. For further information, contact your local Makhteshim Agan of North America, Inc. representative.

### **CHEMIGATION DIRECTIONS FOR USE**

Refer to general **DIRECTIONS FOR USE** section before proceeding with chemigation application.

#### **Types of Irrigation Systems**

Chemigation applications of Pasada 1.6F may be made to crops through overhead sprinkler chemigation systems if specified in crop-specific **Recommended Application** sections. Do not apply Pasada 1.6F through any other type of irrigation system.

#### **Water Volume**

Pasada 1.6F chemigation applications should be made as concentrated as possible. Retention of Pasada 1.6F on target site of insect infestation is necessary for optimum activity. Chemigation of Pasada 1.6F in water volumes exceeding 0.10 inches/acre are not recommended.

#### **Uniform Water Distribution and System Calibration**

The irrigation system must provide uniform distribution of treated water. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

#### **Chemigation Monitoring**

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.

## **Drift**

Do not apply when the wind speed favors drift beyond the area intended for treatment.

## **Required System Safety Devices**

The system must contain a functional check valve, vacuum-relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional, interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

## **Using Water from Public Water Systems**

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (RPZ), or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

**ROTATIONAL CROPS\***

Treated areas may be replanted with any crop specified on an imidacloprid label or any crop for which a tolerance exists for the active ingredient as soon as practical following the last application. For crops not listed on an imidacloprid label or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval should be observed.

**Immediate Plant-back:**

All crops on this label plus the following crops not on this label: barley, canola, corn (field, sweet, and pop), rapeseed, sorghum, soybean, sugar beet, and wheat.

**30-Day Plant-back:**

Cereals (including buckwheat, millet, oats, rice, rye and triticale), safflower

**10-Month Plant-back:**

Onion and bulb vegetables

**12-Month Plant-back:**

All other crops

\*Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.

**FIELD CROPS****Recommended Applications - Pasada 1.6F**

Apply specified rate per acre as a broadcast or directed foliar spray to an infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Pasada 1.6F may not knock down established and heavy insect populations. Two applications may be required to achieve control. Scout fields and re-treat if needed. Pasada 1.6F may be tank mixed with other insecticides as recommended for knockdown of pests or for improved control of other pests.

**COTTON**

Pests Controlled		Rate Fluid Ounces/Acre
Cotton aphid Cotton fleahopper Bandedwinged whitefly Plant bugs (excludes <i>Lygus hesperus</i> ) Green stink bug Southern green stink bug Bollworm/Budworm (ovicidal effect)		2.5 - 5.0
Pests Suppressed		
Lygus bug ( <i>Lygus hesperus</i> ) Whiteflies (other than bandedwinged whitefly)		3.8 – 5.0
<b>Notes and Restrictions</b>		
Pre-Harvest Interval (PHI): <b>14 days</b>		
Minimum interval between applications: <b>7 days</b>		
Maximum Pasada 1.6F allowed per crop season: <b>25.0 fluid ounces/acre</b> (0.31 lb. a.i./a)		
Do not graze treated fields after any application of Pasada 1.6F.		
Pasada 1.6F may be applied through properly calibrated ground, aerial, or chemigation application equipment.		
Tank Mix Recommendations		
Pests Controlled (in addition to pests listed above)	Pasada 1.6F Rate Fluid Ounces/Acre	Bidrin® 8* Rate Fluid Ounces/Acre
<b>For early season control of:</b> Thrips	2.5 – 3.8	1.6 – 3.2
<b>For mid- to late-season control of:</b> Plant bugs Stink bugs (including Brown stink bug) Grasshoppers Saltmarsh caterpillar Cotton leafperforator	2.5 – 3.8	4.0 – 8.0
<b>Notes and Restrictions (in addition to Notes and Restrictions listed above)</b>		
*Refer to the Bidrin® 8 product label for specific use recommendations; observe all restrictions and precautions that appear on the label.		

**PEANUT <sup>1/</sup>**

Pests Controlled	Rate Fluid Ounces/Acre
Aphids Leafhoppers Whiteflies	3.5
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>14 days</b> Minimum interval between applications: <b>5 days</b> Maximum Pasada 1.6F allowed per crop season: <b>10.5 fluid ounces/acre</b> (0.13 lb. a.i./acre)	
<sup>1/</sup> Use not permitted in California unless otherwise directed by supplemental labeling.	

**POTATO**

Pests Controlled	Rate Fluid Ounces/Acre
Aphids Colorado potato beetle Flea beetles Leafhoppers Psyllids	3.8
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>7 days</b> Minimum Interval between applications: <b>7 days</b> Maximum Pasada 1.6F allowed per crop season: <b>16.0 fluid ounces/acre</b> (0.2 lb. a.i./a)	

**TOBACCO**

Pests Controlled	Rate Fluid Ounces/Acre
Aphids	2.0 – 4.0
Flea beetles Japanese beetle	4.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>14 days</b> Minimum interval between applications: <b>7 days</b> Maximum Pasada 1.6F allowed per crop season: <b>22.4 fluid ounces/acre</b> (0.28 lb. a.i./a)	

**VEGETABLE and SMALL FRUIT CROPS**  
**Recommended Applications - Pasada 1.6F**

Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Pasada 1.6F may not knock down established and heavy insect populations. Two applications may be required to achieve control. Scout fields and re-treat if needed. Pasada 1.6F may be tank mixed with other insecticides as recommended for knockdown of pests or for improved control of other pests.

Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website ([www.epa.gov](http://www.epa.gov)) for latest crop groups.

**FRUITING VEGETABLES<sup>1</sup>**

**Crops of Crop Group 8 Plus Okra including:** Eggplant, Ground cherry, Okra, Pepper (including bell, chili, cooking, pimento, and sweet), Tomato, Pepinos, Tomatillo

Pests Controlled	Rate Fluid Ounces/Acre
Aphids Colorado potato beetle Leafhoppers Whiteflies	3.8 – 6.2
Pepper weevil	6.2

**Notes and Restrictions**

Pre-Harvest Interval (PHI): **0 day**

Minimum interval between applications: **5 days**

Maximum Pasada 1.6F allowed per crop season: **19.2 fluid ounces/acre** (0.24 lb. a.i./a)

**Applications**

For pepper weevil, apply specified dosage of Pasada 1.6F by ground equipment only, timing applications prior to a damaging population becoming established. Good coverage of foliage and fruit is necessary for optimum control. Applications of Pasada 1.6F must be incorporated into a full-season program, where alternations of effective products from multiple classes of chemistry and different modes of action are utilized in a blocked or windowed approach.

For additional information, please contact your Makhteshim Agan of North America, Inc. representative, Extension Specialist, or crop advisor.

When targeting adult whiteflies, use higher rates.

<sup>1</sup>Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.

## GLOBE ARTICHOKE

<b>Pests Controlled</b>	<b>Rate Fluid Ounces/Acre</b>
Aphids Leafhoppers	4.0 –10.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>7 days</b> Minimum interval between applications: <b>14 days</b> Maximum Pasada 1.6F allowed per crop season: <b>40.0 fluid ounces/acre</b> (0.5 lb. a.i./a)	

## HERBS

**Crops of Crop Subgroup 19A including:** Angelica, Balm (lemon balm), Basil (fresh and dried), Borage, Burnet, Camomile, Catnip, Chervil (dried), Chinese chive, Chive, Clary, Coriander (cilantro or Chinese parsley leaves), Costmary, Culantro (leaf), Curry (leaf), Dill weed, Horehound, Hyssop, Lavender, Lemongrass, Lovage (leaf), Marigold, Marjoram, Nasturtium, Parsley (dried), Pennyroyal, Rosemary, Rue, Sage, Savory (summer and winter), Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood

<b>Pests Controlled</b>	<b>Rate Fluid Ounces/Acre</b>
Aphids Flea beetles Leafhoppers Whiteflies	3.5
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>7 days</b> Minimum interval between applications: <b>5 days</b> Maximum Pasada 1.6F allowed per crop season: <b>10.5 fluid ounces/acre</b> (0.13 lb. a.i./a)	
<b>Applications</b> Pasada 1.6F may be applied through properly calibrated ground and aerial application equipment. Thorough coverage with direct contact of the spray material to the target pests is required for optimum control. The addition of an organosilicone-based spray adjuvant at a rate not to exceed the adjuvant manufacturer's recommended use rate may improve coverage and control.	
<b>Note</b> Not all crops and/or varieties listed above have been tested for phytotoxic effects. Without specific knowledge about a particular crop and variety, Makhteshim Agan of North America, Inc. strongly recommends that only small areas or numbers of plants of each be treated and evaluated prior to commercial use.	

**BRASSICA (COLE) LEAFY VEGETABLES<sup>1</sup>**

**Crops of Crop Group 5 including:** Broccoli, Broccoli raab (rapini), Brussels sprouts, Cabbage, Cauliflower, Cavalo broccoli, Chinese (gai lon) broccoli, Chinese (bok choy) cabbage, Chinese (napa) cabbage, Chinese mustard (gai choy) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens

Pests Controlled	Rate Fluid Ounces/Acre
Aphids Flea beetles Leafhoppers Whiteflies	3.8
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>7 days</b> Minimum interval between applications: <b>5 days</b> Maximum Pasada 1.6F allowed per crop season: <b>19.2 fluid ounces/acre</b> (0.24 lb. a.i./a) <sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	

**LEAFY GREEN VEGETABLES<sup>1</sup>**

**Crops of Crop Subgroup 4A Plus Watercress including:** Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (rocket), Chervil, Chrysanthemum (edible leaved and garland), Corn salad, Cress (garden), Cress (upland, yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Spinach (including New Zealand and vine (Malabar spinach, Indian spinach)), Watercress (commercial production only. Applications must not be made to native cress growing in streams or other bodies of water), Watercress (upland)

Pests Controlled	Rate Fluid Ounces/Acre
Aphids Flea beetles Leafhoppers Whiteflies	3.8
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>7 days</b> Minimum interval between applications: <b>5 days</b> Maximum Pasada 1.6F allowed per crop season: <b>19.2 fluid ounces/acre</b> (0.24 lb. a.i./a) <b>Applications</b> For applications made to watercress, production fields must be drained of water at least 24 hours prior to application, and water must not be reapplied to the field for a minimum of 24 hours following the application. Applications must be made to fully leafed-up canopies only. <sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	

## LEGUME VEGETABLES<sup>1</sup>

**Crops of Crop Group 6 (except soybean, dry) including:**

**Edible Podded and Succulent Shelled Pea and Bean and Dried Shelled Pea and Bean**

**Bean** (*Lupinus* spp., including grain lupin, sweet lupin, white lupin, and white sweet lupin)

**Bean** (*Phaseolus* spp., including field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean)

**Bean** (*Vigna* spp., including adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, crowder pea, moth bean, mung bean, rice bean, Southern pea, urd bean, yardlong bean)

**Pea** (*Pisum* spp., including dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea)

**Other Beans and Peas** (Broad bean (fava), chickpea (garbanzo bean), guar, jackbean, lablab bean (hyacinth bean), lentil, pigeon pea, soybean (immature seed), sword bean)

<b>Pests Controlled</b>	<b>Rate Fluid Ounces/Acre</b>
Aphids Leafhoppers Whiteflies	3.5
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>7 days</b> Minimum interval between applications: <b>7 days</b> Maximum Pasada 1.6F allowed per crop season: <b>10.5 fluid ounces/acre</b> (0.13 lb. a.i./a) <sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling.	

## ROOT, TUBEROUS, and CORM VEGETABLES<sup>v</sup>

**Crops of Crop Group 1 (except sugarbeet) plus Kava including:** Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Beet (garden)<sup>v</sup>, Burdock (edible)<sup>v</sup>, Canna (edible, Queensland arrowroot), Carrot<sup>2</sup>, Cassava (bitter & sweet)<sup>2</sup>, Celeriac<sup>2</sup>, Chayote (root), Chervil (turnip-rooted)<sup>v</sup>, Chicory<sup>2</sup>, Chufa, Dasheen (taro)<sup>v</sup>, Ginger, Ginseng, Horseradish, Kava<sup>2</sup>,<sup>v</sup>, Leren, Parsley (turnip-rooted), Parsnip<sup>v</sup>, Radish<sup>2</sup>, Oriental radish (diakon)<sup>v</sup>, Rutabaga<sup>2</sup>, Salsify (black)<sup>v</sup>, Salsify (oyster plant), Salsify (Spanish), Skirret, Sweetpotato<sup>v</sup>, Tanier (cocoyam)<sup>v</sup>, Turmeric, Turnip<sup>v</sup>, Yam bean (jicama, manioc pea), Yam (true)<sup>v</sup>.

**(For recommended applications on Potato, see Field Crops section)**

Pests Controlled	Rate Fluid Ounces/Acre
Aphids Flea beetles Leafhoppers Whiteflies	3.5
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>7 days</b> Minimum interval between applications: <b>5 days</b> Maximum Pasada 1.6F allowed per crop season: <b>3.5 fluid ounces/acre</b> (0.044 lb. a.i./a) on Radish; <b>10.5 fluid ounces/acre</b> (0.13 lb. a.i./a) on other crops. Maximum Pasada 1.6F applications per crop season: <b>1</b> on Radish; <b>3</b> on other crops. <sup>1</sup> Not for use on crops grown for seed unless allowed by state-specific supplemental labeling. <sup>2</sup> Tops or greens from these crops may be utilized for food or feed. <sup>3</sup> Use not permitted in California unless otherwise directed by supplemental labeling.	

## STRAWBERRY

Pests Controlled	Rate Fluid Ounces/Acre
Aphids Spittlebugs Whiteflies	3.8
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>7 days</b> Minimum interval between applications: <b>5 days</b> Maximum Pasada 1.6F allowed per crop season: <b>11.4 fluid ounces/acre</b> (0.14 lb. a.i./a) Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging.	

**TREE, BUSH, and VINE CROPS****Recommended Applications – Pasada 1.6F**

Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Pasada 1.6F may not knock down established and heavy insect populations. Two applications may be required to achieve control. Scout fields and re-treat if needed. Pasada 1.6F may be tank mixed with other insecticides as recommended for knock down of pests or for improved control of other pests. Aerial application of Pasada 1.6F may result in slower activity and reduced control relative to results from ground application. For tree and vine crops, recommended application rates are based on full-size mature trees or vines.

Crops contained within certain crop groups recognized by EPA are subject to change. Refer to EPA website ([www.epa.gov](http://www.epa.gov)) for latest crop groups.

**BANANA and PLANTAIN**

<b>Pests Controlled</b>	<b>Rate Fluid Ounces/Acre</b>
Aphids Leafhoppers Thrips	8.0
<p><b>Notes and Restrictions</b>            Pre-Harvest Interval (PHI): <b>0 day</b>            Minimum interval between applications: <b>14 days</b>            Maximum Pasada 1.6F allowed per crop season: <b>40.0 fluid ounces/acre</b> (0.5 lb. a.i./a)</p> <p><b>Applications</b>            Apply specified dosage of Pasada 1.6F as a broadcast or directed spray to infested area ensuring thorough coverage. Pasada 1.6F may be applied through properly calibrated ground or aerial application equipment. Aerial application of Pasada 1.6F may result in slower activity and reduced control relative to results from ground application.</p> <p>Addition of an organosilicone adjuvant at a rate not to exceed 2.0 fluid ounces/100 gallons finished spray solution may improve coverage and pest control.</p>	

**BUSHBERRY**

**Crops of Crop Subgroup 13B including:** Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Juneberry, Lingonberry, Salal

Pests Controlled	Rate Fluid Ounces/Acre
Aphids Leafhoppers/Sharpshooters	3.0 – 4.0
Blueberry maggot Japanese beetle (adults) Thrips (foliage-feeding thrips only)	6.0 – 8.0
<b>Notes and Restrictions</b>	
Pre-Harvest Interval (PHI): <b>3 days</b>	
Minimum interval between applications: <b>7 days</b>	
Maximum Pasada 1.6F allowed per crop season: <b>40.0 fluid ounces/acre</b> (0.5 lb. a.i./a)	
Maximum number of Pasada 1.6F applications per crop season: <b>5</b>	
Minimum application volume (water): 20.0 gpa – ground; 5.0 gpa – aerial.	
Do not apply pre-bloom or during bloom or when bees are actively foraging.	

**CANEBERRY<sup>1</sup>**

**Crops of the Caneberry Crop Subgroup 13A including:**

**Blackberry** (*Rubus* spp. - including Andean blackberry, Arctic blackberry, Bingleberry, Black satin berry, Boysenberry, Brombeere, California blackberry, Chesterberry, Cherokee blackberry, Cheyene blackberry, Common blackberry, Coryberry, Darrowberry, Dewberry, Dirksen thornless berry, Evergreen blackberry, Himalayaberry, Hullberry, Lavacaberry, Loganberry, Lowberry, Lucretiaberry, Mammoth blackberry, Marionberry, Moras, Mures deronce, Nectarberry, Northern dewberry, Olallieberry, Oregon evergreen berry, Phenomenalberry, Rangeberry, Ravenberry, Rosssberry, Shawnee blackberry, Southern dewberry, Tayberry, Youngberry, Zazzamora, and varieties and/or hybrids of these)

**Raspberry** (*Rubus* spp. - including Bababerry, Black raspberry, Blackcap, Caneberry, Framboise, Frambueso, Himbeere, Keriberry, Mayberry, Red raspberry, Thimbleberry, Tulameen, Yellow raspberry, and varieties and/or hybrids of these, and Wild raspberry)

Pests Controlled	Rate Fluid Ounces/Acre
Aphids Leafhoppers Thrips	8.0
<b>Notes and Restrictions</b>	
Pre-Harvest Interval (PHI): <b>3 days</b>	
Minimum interval between applications: <b>7 days</b>	
Maximum Pasada 1.6F allowed per crop season: <b>24.0 fluid ounces/acre</b> (0.3 lb. a.i./a)	
Do not apply pre-bloom or during bloom or when bees are actively foraging.	
<sup>1</sup> Use not permitted in California unless otherwise directed by supplemental labeling.	

**CITRUS**

**Crops of Crop Group 10 including:** Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Satsuma mandarin, White sapote (*Casimiroa* spp.), and other cultivars and/or hybrids of these

<b>Pests Controlled</b>	<b>Rate Fluid Ounces/Acre</b>
Aphids Asian citrus psyllid Blackfly Leafhoppers/Sharpshooters Leafminers Mealybugs Scales Whiteflies	10.0 – 20.0 (dependent on tree size, target pest and infestation pressure)
<b>Pest Suppressed</b>	
Thrips (foliage-feeding thrips only)	10.0 – 20.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>0 day</b> Minimum interval between applications: <b>10 days</b> Maximum Pasada 1.6F allowed per crop season: <b>40.0 fluid ounces/acre</b> (0.5 lb. a.i./a) Do not apply during bloom or within 10 days prior to bloom or when bees are actively foraging. <b>Applications</b> Scales – time applications to the crawler stage. Treat each generation.	

**COFFEE**

<b>Pests Controlled</b>	<b>Rate Fluid Ounces/Acre</b>
Aphids Leafhoppers Whiteflies	8.0
<b>Pest Suppressed</b>	
Scales	8.0
<b>Notes and Restrictions</b>	
Pre-Harvest Interval (PHI): <b>7 days</b>	
Minimum interval between applications: <b>7 days</b>	
Maximum Pasada 1.6F allowed per crop season: <b>40.0 fluid ounces/acre</b> (0.5 lb. a.i./a)	
Do not apply pre-bloom or during bloom or when bees are actively foraging.	
<b>Applications</b>	
Apply specified dosage of Pasada 1.6F as a broadcast or directed spray to infested area ensuring thorough coverage. Pasada 1.6F may be applied through properly calibrated ground or aerial application equipment. Aerial application of Pasada 1.6F may result in slower activity and reduced control relative to results from ground application.	

**GRAPE**

**Including:** American bunch grape, Muscadine grape, and Vinifera grape

<b>Pests Controlled</b>	<b>Rate Fluid Ounces/Acre</b>
Leafhoppers/Sharpshooters Mealybugs	3.0 – 4.0
Grapeleaf skeletonizer	3.8 – 4.0
<b>Notes and Restrictions</b>	
Pre-Harvest Interval (PHI): <b>0 day</b>	
Minimum interval between applications: <b>14 days</b>	
Maximum Pasada 1.6F allowed per crop season: <b>8.0 fluid ounces/acre</b> (0.1 lb. a.i./a)	
Pasada 1.6F may be applied by ground application only.	

**HOP**

<b>Pests Controlled</b>	<b>Rate Fluid Ounces/Acre</b>
Aphids	8.0
<b>Notes and Restrictions</b>	
Pre-Harvest Interval (PHI): <b>28 days</b>	
Minimum interval between applications: <b>21 days</b>	
Maximum Pasada 1.6F allowed per crop season: <b>24.0 fluid ounces/acre</b> (0.3 lb. a.i./a)	

**POME FRUIT**

**Crops of Crop Group 11 including:** Apple, Crabapple, Loquat, Mayhaw, Pear (including Oriental pear), Quince

<b>Pests Controlled</b>	<b>Rate Fluid Ounces/Acre</b>
Leafhoppers	4.0 – 8.0
Aphids (except woolly apple aphid) Apple maggot Leafminers San Jose scale	8.0
<b>FOR PEAR ONLY</b> Mealybugs Pear psylla	20.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>7 days</b> Minimum interval between applications: <b>10 days</b> Maximum Pasada 1.6F allowed per crop season: <b>40.0 fluid ounces/acre</b> (0.5 lb. a.i./a) Do not apply pre-bloom or during bloom or when bees are actively foraging.	
<b>Applications</b> Applications targeting apple maggot should be combined with manufacturer's recommended rate of a sticker, such as Nu-Film 17.	

**POMEGRANATE**

<b>Pests Controlled</b>	<b>Rate Fluid Ounces/Acre</b>
Aphids Leafhoppers/Sharpshooters Whiteflies	8.0
<b>Pest Suppressed</b>	
Scales	8.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>7 days</b> Minimum interval between applications: <b>7 days</b> Maximum Pasada 1.6F allowed per crop season: <b>24.0 fluid ounces/acre</b> (0.3 lb. a.i./a) Do not apply pre-bloom or during bloom or when bees are actively foraging.	

**STONE FRUIT**

**Crops of Crop Group 12 including:** Apricot, Cherry (including sweet and tart), Nectarine, Peach, Plum (including Chickasaw, Damson, and Japanese), Plumcot, Prune (fresh and dried)

<b>Pests Controlled</b>	<b>Rate Fluid Ounces/Acre</b>
Aphids Green June beetle Japanese beetle Leafhoppers/Sharpshooters Plant bugs Rose chafer San Jose scale	4.0 – 8.0
Cherry fruit fly	6.0 - 8.0
<b>Pests Suppressed</b>	
Plum curculio Stink bugs	8.0
<p><b>Notes and Restrictions for Apricot, Nectarine, Peach</b>            Pre-Harvest Interval (PHI): <b>0 day</b>            Minimum interval between applications: <b>7 days</b>            Maximum Pasada 1.6F allowed per crop season: <b>24.0 fluid ounces/acre</b> (0.3 lb. a.i./a)            Minimum application volume (water): 50 gpa – ground application; 25 gpa – aerial application            Do not apply pre-bloom or during bloom or when bees are actively foraging.</p> <p><b>Notes and Restrictions for Cherries, Plums, Plumcot, Prune</b>            Pre-Harvest Interval (PHI): <b>7 days</b>            Minimum interval between applications: <b>10 days</b>            Maximum Pasada 1.6F allowed per crop season: <b>40.0 fluid ounces/acre</b> (0.5 lb. a.i./a)            Minimum application volume (water): 50 gpa – ground application; 25 gpa – aerial application            Do not apply pre-bloom or during bloom or when bees are actively foraging.</p>	

## TREE NUTS

**Crops of Crop Group 14 including:** Almond, Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert, Hickory nut, Macadamia nut, Pecan, Pistachio, Walnut (black and English)

<b>Pests Controlled</b>	<b>Rate Fluid Ounces/Acre</b>
Aphids (except Black pecan aphid) Leafhoppers/Sharpshooters <i>Phylloxera</i> spp. (leaf infestations) Spittlebugs Whiteflies	3.5 – 7.0
Black pecan aphid Mealybugs San Jose scale	8.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>7 days</b> Minimum interval between applications: <b>6 days</b> Maximum Pasada 1.6F allowed per crop season: <b>28.8 fluid ounces/acre</b> (0.36 lb. a.i./a) Minimum application volume (water): 50 gpa – ground application, 25 gpa – aerial application Do not apply pre-bloom or during bloom or when bees are actively foraging.	
<b>Applications</b> Applications for control of San Jose scale should be timed according to crawler stage, treating each successive generation. Two applications on a 10- to 14-day interval may be required to achieve control.	

## TROPICAL FRUIT

**Including:** Acerola, Atemoya, Avocado, Birida, Black sapote, Canistel, Cherimoya, Custard Apple, Feijoa, Jaboticaba, Guava, Llama, Longan, Lychee, Mamey sapote, Mango, Papaya, Passionfruit, Persimmon, Pulasan, Rambutan, Sapodilla, Soursop, Spanish lime, Star apple, Starfruit, Sugar apple, Wax jambu

<b>Pests Controlled</b>	<b>Rate Fluid Ounces/Acre</b>
Aphids Leafhoppers/Sharpshooters Mealybugs Thrips (foliage-feeding thrips only) Whiteflies	8.0
<b>Pest Suppressed</b>	
Scales	8.0
<b>Notes and Restrictions</b> Pre-Harvest Interval (PHI): <b>7 days</b> Minimum interval between applications: <b>10 days</b> Maximum Pasada 1.6F allowed per crop season: <b>40.0 fluid ounces/acre</b> (0.5 lb. a.i./a) Do not apply pre-bloom or during bloom or when bees are actively foraging.	

**OTHER CROPS****Recommended Applications – Pasada 1.6F**

Apply specified rate per acre as a broadcast or directed foliar spray to infested area as pest populations begin to build. Thorough uniform coverage is necessary to achieve optimum control. A spray adjuvant may be used to improve coverage. Pasada 1.6F may not knock down established and heavy insect populations. Two applications may be required to achieve control. Scout fields and re-treat if needed. Pasada 1.6F may be tank mixed with other insecticides as recommended for knockdown of pests or for improved control of other pests.

**CHRISTMAS TREE**

Pests Controlled	Rate Fluid Ounces/Acre
Aphids Adelgids Sawflies	4.0 - 8.0
<b>Notes and Restrictions</b> Minimum interval between applications: <b>7 days</b> Maximum Pasada 1.6F allowed per crop season: <b>40.0 fluid ounces/acre</b> (0.5 lb. a.i./a) <b>Applications</b> Gall-forming adelgids – time applications to coincide with full bud-swell or first bud-break of earliest bud-breaking trees. Once galls form, spraying will be ineffective.	

**POPLAR/COTTONWOOD<sup>v</sup>**

**Including:** Members of the genus *Populus* grown for pulp or timber

Pests Controlled	Rate Fluid Ounces/Acre
Aphids Leaf beetles	4.0 - 8.0
<b>Notes and Restrictions</b> Minimum interval between applications: <b>10 days</b> Maximum Pasada 1.6F allowed per crop season: <b>40.0 fluid ounces/acre</b> (0.5 lb. a.i./a) Do not apply pre-bloom or during bloom or when bees are actively foraging. <sup>v</sup> Use not permitted in California unless otherwise directed by supplemental labeling.	

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry place and in such a manner as to prevent cross-contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to **PRECAUTIONARY STATEMENTS** on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing-type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### IMPORTANT: READ BEFORE USE

#### LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions, disclaimer of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **Conditions, Disclaimer of Warranties and Limitations of Liability.**

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, where in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid, or at Makhteshim Agan of North America, Inc.'s election, the replacement of product.

Pasada and Alias are registered trademarks of Celsius Property B.V.

Admirer Pro, Calypso, Gaucho, Leverage, Provado and Trimax Pro are registered trademarks of Bayer.

Actara, Centric, Cruiser, and Platinum are registered trademarks of the Syngenta Group Company.

Assail and Intruder are registered trademarks of Nippon Soda Company, LTD.

Belay and Clutch are registered trademarks of Arvesta Corporation.

Venom is a registered trademark of Valent USA Corporation.

Bidrin is a registered trademark of AMVAC Chemical Corporation.

Couraze is a trademark of Cheminova, Inc.

Macho and Impulse are trademarks of Albaugh, Inc.

Advise and Gallant are trademarks of Agrilience, LLC.

Widow is a trademark of Loveland Products, Inc.

Nuprid is a trademark of NuFarm America, Inc.



# Pasada<sup>®</sup> 1.6F

FLOWABLE INSECTICIDE

For control of certain insects infesting various crops.

**GROUP 4A INSECTICIDE**

ACTIVE INGREDIENT	% BY WT.
Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]	
-N-nitro-2-imidazolidinimine	17.4%
<b>INERT INGREDIENTS:</b>	<b>82.6%</b>
<b>TOTAL</b>	<b>100.0%</b>

Contains 1.6 pounds of active ingredient per gallon.

**SHAKE WELL BEFORE USING**

**STOP - Read the label before use**  
**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

FIRST AID	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>

EPA Reg. No. 264-763-66222

FIRST AID	
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For medical emergencies involving this product, contact Prostar at 1-877-250-9291.	
<b>NOTE TO PHYSICIAN:</b> No specific antidote is available. Treat the patient symptomatically.	

**For additional precautionary, handling, and use statements, see inside of this booklet.**

  
 Manufactured for:  
**Makhteshim Agan**  
 of North America, Inc.  
 4515 Falls of Neuse Road  
 Suite 300  
 Raleigh, NC 27609

EPA 081307/  
 Notif 092109/Rev B  
 US04321921C

**Net Contents: 1 Gallon**