59807-2_book_art 10/22/14 11:49 AM Page 1

STATE OF HAWAII
Department of Agriculture
ACCEPTED
LICENSE NO.
8282.17

GROUP 21A INSECTICIDE

FujiMite® XLO

MITICIDE/INSECTICIDE

ACTIVE INGREDIENT:

Fenpyroximate: *Tert*-butyl(E)-α-(1,3-dimethyl-5-phenoxypyrazol-4-ylmethyleneaminoxy)-p-toluate **5.0%**

OTHER INGREDIENTS*: **95.0%**

TOTAL **100.0%**

Contains 0.4 lb. active ingredient per U.S. gallon

*Contains petroleum distillates

EPA Reg. No. 71711-40 **EPA Est. No. 67545-AZ-1 (GM) 39578-TX-1 (E) 70815-GA-001 (CB)**

superscript corresponds with lot number

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See inside booklet for First Aid, Precautionary Statements, and Directions for Use

NET CONTENTS: 2.5 gallons

325504
10/14

NICHINO AMERICA®
Nichino America, Inc.
4550 New Linden Hill Road
Wilmington, DE 19808

file: 59807-2_book_art
folder: 59807-2_nichino_208171-26
colors: black, 3025
☐ dp proofed_____

created by: 04-06-09 ek
alt: 10-22-14 ek
size: 6.25" (w) x 7" (h)

FIRST AID	
If inhaled	• Move the person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
If swallowed	• Call a poison control center or doctor immediately for treatment advice. • Do not give any liquid to the person. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-348-5832 for emergency medical treatment information. In case of fire or spills, information may be obtained by calling 1-800-424-9300.	
NOTE TO PHYSICIAN: May pose an aspiration pneumonia hazard. Contains petroleum distillates.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING - AVISO

May be fatal if inhaled. Causes substantial but temporary eye injury. Harmful if swallowed. Avoid contact with skin or clothing. Do not breathe spray mist. Remove and wash contaminated clothing before reuse. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves (such as barrier laminate, nitrile rubber, neoprene rubber, viton, Selection Category E).

For handling activities, wear respirator with an organic vapor removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P, or HE prefilter.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category E on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves (such as barrier laminate, butyl rubber, nitrile rubber, or viton)
- Protective eyewear (such as safety glasses, goggles, or face shield)
- Shoes plus socks

- For handling activities, use either a respirator with an organic vapor removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P, or HE prefilter.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. When handlers use closed systems *or* enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is very highly toxic to fish and aquatic invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having medium to high potential for reaching both surface water and aquatic sediment via runoff for several weeks after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this chemical from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's potential to reach aquatic sediment via runoff.

ENDANGERED SPECIES RESTRICTIONS

This product may pose a hazard to endangered aquatic species. Follow all use directions.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

Minimum Honey Bee Toxicity

Fenpyroximate is practically nontoxic to bees and wasps when fenpyroximate is applied to listed crops according to the label directions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency in your State responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls worn over short-sleeved shirt and short pants
- Socks
- Chemical-resistant footwear
- Protective eyewear (such as safety glasses, goggles, or face shield)
- Chemical-resistant gloves (such as barrier laminate, butyl rubber, nitrile rubber, or viton)

GENERAL INFORMATION

FUJIMITE® XLO miticide/insecticide is used for the control of leafhoppers, mealybugs, mites, psylla, psyllids, and whiteflies.

FUJIMITE XLO miticide/insecticide stops mite feeding immediately after application. **FUJIMITE XLO** miticide/insecticide controls all motile stages of mites by inhibiting cellular respiration in the mitochondrion of cells which results in rapid cessation of all biological activities, including feeding and reproduction. Mortality of mites can be observed within 3 to 7 days after intoxication.

FUJIMITE XLO miticide/insecticide works primarily through contact action, **so thorough spray coverage is necessary**. Mix with sufficient water and apply as a foliar spray to obtain uniform coverage. Dense foliage or excessive growth will often prevent adequate coverage; adjust spray volumes accordingly. Treat plants when pests are immature or at a susceptible stage and populations are building, before crop damage occurs.

TARGET SPECIES		
Apple rust mite (suppression only)	Glassy-winged sharpshooter (suppression only)	Powdery mildew (suppression only)
Asian citrus psyllid	Grape leafhopper	Strawberry spider mite
Broad mite	McDaniel mite	Texas citrus mite
Carmine mite	Mealybug species	Tomato/Potato psyllid
Citrus bud mite	Mint bud mite	Tomato russet mite
Citrus leafminer (suppression only)	Pacific spider mite	Two-spotted spider mite
Citrus red mite	Pear psylla	Variegated leafhopper
Citrus rust mite	Pear rust mite	White apple leafhopper
Citrus thrips (suppression only)	Pecan leaf scorch mite	Whiteflies
Cyclamen mite	Potato leafhopper	Willamette spider mite
European red mite		

APPLICATION DIRECTIONS

- Applications should be made immediately after the spray solution is prepared.
- Apply with properly calibrated spray equipment.
- Apply by ground using the recommended water spray volume found in the **Directions for Use** section of this label.
- Thorough spray coverage is essential for mite and insect control.
- For best results, apply when pest populations are beginning to build, before reaching economic thresholds. Consult your local agricultural advisor or state cooperative extension service for further information.

USE OF ADJUVANTS

When thorough coverage is a concern, it is recommended that a spray adjuvant be used to maximize uniformity of coverage and performance of **FUJIMITE XLO** miticide/insecticide. Use a nonionic activator type wetting, spreading, or penetrating adjuvant or horticultural spray oil adjuvant. Do not use a dormant oil or binder or sticker type adjuvant. Nonionic adjuvants (NIS) should contain at least 75% surfactant. Crop oil concentrates (COC), methylated seed or vegetable oils (MSO), organosilicone products (OS), or blends of these adjuvants should contain at least 15% emulsifier/surfactant. Check compatibility of any adjuvant used with **FUJIMITE XLO** miticide/insecticide before using. Follow the Directions for Use on each adjuvant product label for rates of use and use restrictions.

APPLICATION RESTRICTIONS

- Do not apply within 75 feet of fish-bearing waters.
- Do not use products with the same mode of action in consecutive applications.
- Do not plant rotational crops other than those listed on this label for 30 days following the last application of this product.
- Do not apply through any type of irrigation system.
- Do not apply by Alternate Row Middle (ARM) spray method.

RESISTANCE MANAGEMENT

Repeated use of the same chemistry has been shown to result in the buildup of resistant strains of mites or other insects. Do not use **FUJIMITE XLO** miticide/insecticide in successive miticide applications. Rotate the use of **FUJIMITE XLO** miticide/insecticide with alternate mode of action insecticides:

- Miticides must be rotated with alternate products **for resistance management**.
- **Do not use products with the same mode of action (METI-2 inhibitor/IRAC Group 21A) in consecutive applications**.

Consult your local crop advisor for the most appropriate alternative products. Resistance management strategies recommend that you DO NOT apply rates lower than recommended on the label.

MIXING DIRECTIONS

FUJIMITE XLO Miticide/Insecticide Alone: Shake well before using. Begin with clean equipment. Fill spray tank with ¾ of the amount of water needed for the intended application and then turn on agitation. Pour recommended amount of product on the surface of water in the spray tank. Add the balance of the water to the spray tank with agitation running. Keep agitation running during filling and spraying operations. If spraying must be stopped before emptying the sprayer, resume agitation before spraying the remainder of the load.

FUJIMITE XLO Miticide/Insecticide Tank Mixtures: Shake well before using. Read and follow all label directions for each tankmix product prior to any tank mixing with **FUJIMITE XLO** miticide/insecticide. This product can be mixed with other registered pesticides for use on labeled crops or sites, in accordance with the most restrictive use directions and precautions. Follow all use directions as listed above under **FUJIMITE XLO Miticide/Insecticide Alone** with the following exception: after the **FUJIMITE XLO** miticide/insecticide is thoroughly mixed and the tank is ¾ full, add the recommended amount of wettable powder, soluble powder, flowable, emulsifiable concentrate, or soluble liquid product, **while maintaining agitation**. Then continue adding water to the tank to achieve the desired level, while maintaining agitation.

If you have no experience with the combination you are considering, you should conduct a test to determine physical compatibility. To determine physical compatibility, add the recommended proportions of each chemical with the same proportion of water as will be present in the chemical supply tank into a suitable container; mix thoroughly and allow to stand for five minutes. If the combination remains mixed, or can be readily remixed, the mixture is considered physically compatible.

SPRAY DRIFT MANAGEMENT

Avoid spray drift to all other crops and nontarget areas. Do not apply when weather conditions may cause drift. Do not allow this product to drift onto nontarget areas. Drift may result in illegal residues or injury to adjacent crops and vegetation. Use of larger droplet size will also reduce spray drift.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR. The interaction of equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

Droplet size, boom height, and wind speed are the primary factors determining drift. The specific application conditions required for the use of this product are described below.

Controlling Droplet Size – General Techniques

Volume

Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure

Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.

Nozzle Type

Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Height and Length – Ground

Boom Height

Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

Wind

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. AVOID GUSTY OR WINDLESS CONDITIONS. **Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. This cloud can move in unpredictable directions due to the light and variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

Shielded Sprayers
Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with the uniform deposition of the product.

Air Assisted (Air Blast) Field Crop Sprayers
Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, is configured properly, and that drift is not occurring.

Note: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the application equipment section of this label to determine if use of an air assisted sprayer is recommended.

Air Assisted (Air Blast) Tree And Vine Sprayers
Air assisted tree and vine sprayers carry droplets into the canopy of trees and vines via a radially or laterally directed air stream. These sprayers are not suitable for applying herbicides. In addition to the general drift management practices already described, the following specific practices will further reduce the potential for drift:

- Adjust the deflectors and aiming devices so that spray is only directed into the canopy.
- Block off upward pointed nozzles when there is no overhanging canopy.
- Use only enough air volume to penetrate the canopy and provide good coverage.
- Do not allow spray to go beyond the edge of the cultivated area. Spray the outside row only from outside the planting.

APPLICATION RATE CHART FOR FUJIMITE XLO MITICIDE/INSECTICIDE

Almond; Pistachio		
Pest	**Rate/Acre**	**Use Directions**
Mites (see **Target Species**)	2.0 to 4.0 pints	USE RESTRICTIONS • Apply by ground using a minimum of 100 gallons of water per acre. • Do not apply by air. • Do not apply through any type of irrigation system. • Do not apply more than 8.0 pints per acre per growing season. • Do not make more than 2 applications per growing season. • Allow 14 days between applications. • Preharvest Interval (PHI): 14 days

(continued)

APPLICATION RATE CHART FOR FUJIMITE XLO MITICIDE/INSECTICIDE *(continued)*

Berry, Low Growing (Crop Subgroup 13-07G) - except cranberry
bearberry; bilberry; blueberry, lowbush; cloudberry; lingonberry; muntries; partridgeberry; strawberry; cultivars, varieties, and/or hybrids of these

Pest	Rate/Acre	Use Directions
Mites (see **Target Species**) Whiteflies	2.0 pints	USE RESTRICTIONS • Apply by ground application using a minimum of 25 gallons of water per acre. • Do not apply by air. • Do not apply through any type of irrigation system. • Do not apply more than 4.0 pints per acre per crop cycle. • Do not make more than 2 applications per crop cycle. • Allow 14 days between applications. • Preharvest Interval (PHI): 1 day

NOTE
- Temporary pinking of immature green berries may be observed after a **FUJIMITE XLO** miticide/insecticide application on certain strawberry varieties. This effect is transient and does not affect fruit sizing, color, or quality.
- Avoid puddling of spray solution on plastic mulch as this can potentially result in underside scarring of fruit in direct contact with the plastic.

Citrus Fruit (Crop Group 10-10)
Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; mount white lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin orange, clementine); tangor; trifoliate orange; uniq fruit; cultivars, varieties, and/or hybrids of these

Pest	Rate/Acre	Use Directions
Asian citrus psyllid[1] Citrus thrips (suppression only) Citrus leafminer (supression only) Leafhoppers Mealybugs Other mites (see **Target Species**)	2.0 to 4.0 pints	USE RESTRICTIONS • Apply by ground using a minimum of 100 gallons of water per acre. For full size trees, use a minimum of 200 gallons of water per acre. • Do not apply by air. • Do not apply through any type of irrigation system. • Do not apply to citrus nurseries or citrus in greenhouses. • Do not apply more than 8.0 pints per acre per growing season. • Do not make more than 2 applications per growing season. • Allow 14 days between applications. • Preharvest Interval (PHI): 14 days

[1] For best results, use for control of adults and nymphs present at time of application when newly expanding foliage flush is present.

(continued)

APPLICATION RATE CHART FOR FUJIMITE XLO MITICIDE/INSECTICIDE *(continued)*

Grape
Small Fruit Vine Climbing – except fuzzy kiwifruit (Crop Subgroup 13-07F)
Amur river grape; gooseberry; raisin, table, wine grape; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and/or hybrids of these

Pest	Rate/Acre	Use Directions
Mealybugs Mites (see **Target Species**) Powdery mildew (suppression)	2.0 pints	USE RESTRICTIONS • Apply by ground using a minimum of 50 gallons of water per acre. • For vines with a heavy canopy, or in high pressure situations, higher water volumes are recommended. If lower water volume amounts are used, tractor speed must be reduced to ensure complete coverage. • Do not apply by air. • Do not apply through any type of irrigation system. • Do not apply more than 2.0 pints per acre per growing season. • Do not make more than 2 applications per growing season. • Allow 14 days between applications. • Preharvest Interval (PHI): 14 days
Willamette spider mite	1.5 to 2.0 pints	
Leafhoppers	1.0 to 2.0 pints [1]	

[1] Use higher rate for dense foliage. Best control of leafhoppers is achieved by applications when majority of the population is in an immature development stage.

Hops

Pest	Rate/Acre	Use Directions
Mites (see **Target Species**)	2.0 to 3.0 pints	USE RESTRICTIONS • Apply by ground using a minimum of 100 gallons of water per acre. • Do not apply by air. • Do not apply through any type of irrigation system. • Do not apply more than 3.0 pints of product per acre per growing season. • Do not apply more than 1 application per growing season. • Preharvest Interval (PHI): 15 days RECOMMENDATIONS • For best results, apply before mite populations exceed 5 mites per leaf.

NOTE
Leaf yellowing may occur when **FUJIMITE XLO** miticide/insecticide is combined with spray oil in excess of 1% of the spray volume. If this symptom occurs, it is usually more pronounced on newly expanding leaves. This symptom may occur in plants under stress and is worsened by certain conditions including the following:
• High Temperatures (air temperatures exceeding 90˚F at the time of application or within a few days after application).
• Wet soil conditions and high humidity (rainy, misty, or foggy weather within a few days after application).
• Storm damage (including hail and wind).

(continued)

APPLICATION RATE CHART FOR FUJIMITE XLO MITICIDE/INSECTICIDE *(continued)*

Mint peppermint; spearmint		
Pest	**Rate/Acre**	**Use Directions**
Mites (see **Target Species**)	1.0 to 2.0 pints	USE RESTRICTIONS • Apply by ground using a minimum of 25 gallons of water per acre. • Do not apply by air. • Do not apply through any type of irrigation system. • Do not apply more than 4.0 pints per acre per growing season. • Do not make more than 2 applications per growing season. • Allow 7 days between applications. • Preharvest Interval (PHI): 1 day

Nonbearing Deciduous Fruit, Nut Trees, and Vines		
Pest	**Rate/Acre**	**Use Directions**
Leafhoppers Mealybugs Mites (see **Target Species**)	2.0 pints	USE RESTRICTIONS • Apply by ground using a minimum of 75 gallons of water per acre. • Do not apply by air. • Do not apply through any type of irrigation system. • Do not apply to citrus nurseries or citrus in greenhouses. • Do not apply more than 2.0 pints per acre per growing season. • Do not make more than 1 application per growing season. • Do not harvest edible crops for 12 months following application unless the crop is listed on the label.

(continued)

APPLICATION RATE CHART FOR FUJIMITE XLO MITICIDE/INSECTICIDE *(continued)*

Pome Fruit (Crop Group 11-10)
apple; azarole; crabapple; loquat; mayhaw; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these

Pest	Rate/Acre	Use Directions
Leafhoppers Mealybugs Mites (see **Target Species**) Pear psylla	2.0 pints	USE RESTRICTIONS • Apply by ground using a minimum of 100 gallons of water per acre. • Do not apply by air. • Do not apply through any type of irrigation system. • Do not apply by Alternate Row Middle (ARM) spray method. • Do not apply more than 2.0 pints per acre per growing season. • Do not make more than 1 application per growing season. • Preharvest Interval (PHI): 14 days

Stone Fruit (Crop Group 12-12)
apricot; apricot, Japanese; capulin; cherry, black; cherry, nanking; cherry, sweet; cherry, tart; jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these

Pest	Rate/Acre	Use Directions
Mites (see **Target Species**) Leafhoppers	2.0 pints	USE RESTRICTIONS • Apply by ground using a minimum of 80 gallons of water per acre. • Do not apply by air. • Do not apply through any type of irrigation system. • Do not apply more than 4.0 pints per acre per growing season. • Do not make more than 2 applications per year. • Allow 14 days between applications. • Preharvest Interval (PHI): 7 days

Tree Nut (Crop Group 14) - excluding almond and pistachio
beech nut; Brazil nut; butternut; cashew; chestnut; chinquapin; filbert (hazelnut); hickory nut; macadamia nut; pecan; walnut, black and English

Pest	Rate/Acre	Use Directions
Mites (see **Target Species**)	2.0 to 4.0 pints	USE RESTRICTIONS • Apply by ground using a minimum of 100 gallons of water per acre. • Do not apply by air. • Do not apply through any type of irrigation system. • Do not apply more than 4.0 pints per acre per growing season. • Do not make more than 2 applications per growing season. • Allow 14 days between applications. • Preharvest Interval (PHI): 14 days

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Nonrefillable plastic containers - for liquid dilutables less than 5 gallons

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable plastic containers - for any dilutable pesticide greater than 5 gallons

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Refillable plastic containers

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Return to point-of-sale or offer for recycling if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by State and local authorities.

IMPORTANT: READ BEFORE USE

By using this product, user or buyer accepts the following conditions, warranty, disclaimer of warranties, and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be accurate and must be followed carefully. However, because of extreme weather and soil conditions, use methods and other factors beyond the control of Nichino America, Inc. (NAI), it is impossible for NAI to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. To the extent consistent with applicable law, all such risks are assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of NAI is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, NAI disclaims any liability whatsoever for incidental or consequential damages, including, but not limited to, liability arising out of breach of contract, express or implied warranty (including warranties of merchantability and fitness for a particular purpose), tort, negligence, strict liability, or otherwise.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT THE ELECTION OF NICHINO AMERICA, THE REPLACEMENT OF PRODUCT.

© 2014 Nichino America, Inc. FujiMite and Nichino America logo are registered trademarks of Nichino America, Inc.

Formulated and Packaged in U.S.A. for
Nichino America, Inc.
4550 New Linden Hill Road
Wilmington, DE 19808
888-740-7700

SL-179 071113-2
11-13-2013

NOTES

NOTES

59808-2_base_art 10/13/14 9:02 AM Page 1

GROUP **21A** INSECTICIDE

FujiMite® XLO

MITICIDE/INSECTICIDE

ACTIVE INGREDIENT:

Fenpyroximate: *Tert*-butyl(E)-α-(1,3-dimethyl-5-phenoxypyrazol-4-ylmethyleneaminoxy)-p-toluate **5.0%**

OTHER INGREDIENTS*: **95.0%**

TOTAL **100.0%**

Contains 0.4 lb. active ingredient per U.S. gallon

*Contains petroleum distillates

EPA Reg. No. 71711-40 **EPA Est. No. 67545-AZ-1 (GM) 39578-TX-1 (E) 70815-GA-001 (CB)**

superscript corresponds with lot number

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See attached booklet for First Aid, Precautionary Statements, and Directions for Use

NET CONTENTS: 2.5 gallons

325504
10/14

NICHINO AMERICA®
Nichino America, Inc.
4550 New Linden Hill Road
Wilmington, DE 19808

file: 59808-2_base_art
folder: 59808-2_nichino_208171-26
colors: black, 3025
☐ dp proofed_____

created by: 04-06-09 ek
alt: 10-13-14 ek
size: 7" (w) x 7" (h)