



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2009-2011** LIC. NO.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2012-2014** LIC. NO.



Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD **2015-2017** LIC. NO.

8283.7

front label

ESL012109N REV031309

Clear Choice™

Lawn Friendly Weed Remover†

NEW

MICROTECHNOLOGY
Improves penetration of reduced active ingredient formulation!®

Step 1: Shake well

Step 2: Apply

Step 3: Let dry

Ready to use! No mixing required!†

**Kills listed broadleaf weeds in lawns -
right down to the root.**

Won't harm lawns when used as directed.†

KILLS: Dandelions, Plantains, Clover, Chickweed, Bedstraw, Black Medic, English Daisy, Curled Dock, Heal-all, Ground Ivy (Creeping Charlie), Knotweed, Ragweed, Shepherd's Purse, plus over 60 other lawn weeds.

www.todaysclearchoice.com

000795-B/1

† Important: Do not apply to Bentgrass and Floratam. /
No se aplique al Bentgrass y a Floratam. See full list of
controlled weeds and use directions on peel away label.


® When compared to other pesticides using these
active ingredients.

Formulated in Canada for: Petro-Canada
385 Southdown Road
Mississauga, Ontario L5K 1A8

EPA Reg. No. 69526-12

EPA Est. No. 79693-CAN-001

PEEL HERE >



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS.

- Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

ENVIRONMENTAL HAZARDS

- This pesticide is toxic to fish and aquatic invertebrates. This product may adversely affect non-target plants. Do not apply to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply to areas where surface water is present, or to intertidal areas below the mean high water mark, except as noted on appropriate labels. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas, and to non-target plants. Do not apply when windy. To prevent product run-off, do not water to the point of runoff, or apply when raining or when rain is expected that day. Do not contaminate water when disposing of equipment wash waters or rinsate.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read and follow all label directions.

GENERAL PRECAUTIONS AND RESTRICTIONS

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply near lakes, streams, rivers or ponds.
- Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not apply in windy conditions.
- Do not apply as a fine mist because of potential injury to desirable plants.

WHEN TO USE

- Apply when weeds are young and actively growing.
- Do not water within 24 hours after spraying.
- Spray when daytime temperatures are above 45° F and below 90° F.
- Do not apply when the lawn is under severe stress from drought, insect attack or high temperatures.
- Do not use more product than specified on the label.
- Newly seeded lawns may be treated after the third mowing.
- If the lawn is dry, water 24 hours before using this product.

WHERE TO USE

This product is used as a spot treatment to control broad leaf weeds on a variety of lawns, in yards, vacant lots and other grassy areas.

- **IMPORTANT:** Do not use this product on Bentgrass and Floratam, a variety of St. Augustine grass. Certain hybrid bermudagrasses vary in tolerance to this product. Temporary discoloration may occur on St. Augustine and Centipede.

LIST OF WEEDS: This product controls the following weeds: Bedstraw, Bindweed (field), Birdsfoot trefoil, Black medic, Blackseed plantain, Blueweed, Brambles, Broad-leaved plantain, Buckhorn plantain, Bull thistle, Burdock, Buttercup, Canada thistle, Carolina geranium, Chickweed, Chicory, Clover, Cocklebur, Common chickweed, Cudweed, Daisy fleabane, Dandelion, Devil's paint brush (hawkweed), Dock (curled), English daisy, Evening primrose, Fall dandelion (fall hawkbit), False dandelion, Fleabane, Florida pusley, Frenchweed poison oak, Grass leaved stichwort, Ground ivy (creeping Charlie), Hawkweed, Heal-all, Henbit, Honeysuckle, Jimsonweed, Knotweed, Kochia, Kudzu, Lambsquarters, Little starwort, Mallow, Morningglory, Mouse-ear chickweed, Mustards,

LIST OF WEEDS (cont'd): Narrow-leaved plantain, Oxalis (yellow woodsorrel), Ox-eye daisy, Pennywort, Peppergrass, Pigweed, Pineapple weed, Plantains (narrow, buckhorn, broadleaf), Poison ivy, Poorjoe, Povertyweed, Prostrate pigweed, Purslane, Ragweed, Russian thistle, Sandwort (thyme-leaved), Sheep sorrel, Shepherd's purse, Smartweed (green), Soliva, Sow thistle, Speedwell (purslane leaved), Spotted spurge, Spreadwell (annual), Spurge, Stitchwort, Stonecrop (mossy), Sumac, Veronica (thyme-leaved), Vervain, Vetch, Violet, Wild aster, Wild blackberry, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild raspberry, Yarrow, Yellow hawkweed

HOW TO USE

- This product is ready-to-use. Do not mix or dilute.
- Shake well before use.
- Open spray nozzle.
- Adjust nozzle to provide desired spray pattern.
- Hold spray about 12 inches from weed.
- Aim at center of weed and spray to lightly cover. Do not overspray.
- Rotate nozzle to closed position.
- Weeds begin to die after 12 hours. It may take up to 4 weeks for a weed to completely die.
- If necessary, repeat application after 8 weeks.

STORAGE AND DISPOSAL

- Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.
- Pesticide Storage: Keep out of reach of children and animals. Keep from freezing. Keep pesticide in original container. Do not put into food or drink containers. Store in a cool, dry place, in a locked storage area.
- Pesticide and Container Disposal: If empty: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If recycling unavailable, place container in trash.
- If partly filled: Call your local solid waste agency for disposal instructions.
- Never place unused product down any indoor or outdoor drain.

CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

It is impossible to eliminate all risks inherently associated with the use of this product. Injury to desirable plants, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Petro-Canada, the manufacturer, or seller. All such risks shall be assumed by the buyer. To the extent consistent with applicable law, neither Petro-Canada, the manufacturer nor seller shall be liable for consequential, special or indirect damages resulting from the use or handling of this product. Except as provided by applicable law or except as warranted by this label, Petro-Canada, the manufacturer or seller makes no warranties, guarantees, or representations of any kind, either express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. Except as provided by applicable law, buyer's or user's exclusive remedy, and Petro-Canada's, the manufacturer's or seller's total liability, shall be for damages not exceeding the cost of the product.

Formulated in Canada for: Petro-Canada
385 Southdown Road, Mississauga, Ontario L5K 1A8

EPA Reg. No. 69526-12
EPA Est. No. 79693-CAN-001