

Use this in hospitals, medical and dental offices, nursing homes, day care centers, nurseries, restaurants, bars, cafeterias, institutional kitchens, food processing plants, food storage areas, convenience stores, institutional facilities, public rest rooms, institutions, schools, colleges, universities, athletic facilities, locker rooms, dressing rooms, shower and bath areas, exercise facilities, campgrounds, playgrounds, public facilities, transportation terminals, hotels and motels.

WHERE TO USE: on washable hard nonporous surfaces: floors, walls, cabinets, counters, sinks, appliances, nonwooden picnic tables, bathtubs, shower stalls, shower walls, glass shower doors tubs and glazed tiles, urinals, toilets, toilet bowls, toilet bowl surfaces, chrome fixtures, other bathroom fixtures, and hard nonporous surfaces made of metal, stainless steel, chrome, glazed porcelain, glazed ceramic, sealed stone, fiberglass and plastic, enameled surfaces, finished woodwork, vinyl and plastic upholstery. Test a small area first. Do not use on PVD finished surfaces. Note: Some older tubs or sinks may be etched or discolored by Betco® Sure Bet™ II. Test on a small hidden area first. Do not use on decorative brass fixtures, polished aluminum, natural or synthetic marble. Do not place container on Formica®.

This product is a Hospital disinfectant, bactericidal at 1:32 dilution (4 oz. per gallon of water) according to the AOAC Use-Dilution Test method, modified in the presence of 400 ppm hard water plus 5% organic serum against: Pseudomonas aeruginosa, Salmonella enterica, Staphylococcus aureus.

PELIGRO:

Si no sabe el ingles, solicite instrucciones a su supervisor antes de usar este producto.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PREPARATION OF DISINFECTANT USE-SOLUTION: Add 4 oz. per gallon of water (for 400 ppm water hardness).

DISINFECTION DIRECTIONS: Apply use-solution with a cloth, sponge, or sprayer or by soaking until surfaces are thoroughly wet. Treated surfaces must remain wet for 10 minutes. Wipe dry with a clean cloth or sponge or let air dry. For sprayer applications, use a coarse spray device. Spray with quick full strokes 6 - 8 inches from surface. Rub with a brush, sponge or cloth. Do not breathe spray. After use, close nozzle. For heavily soiled areas, a preliminary cleaning is required. Prepare a fresh solution daily or more often if use solution becomes diluted or soiled. Rinse all surfaces that come in contact with food such as countertops, appliances, tables and stovetops with potable water before reuse. This product must not be used to clean utensils, glassware and dishes. This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that [1] is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or [2] contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

***VIRUCIDAL DIRECTIONS:** Follow Disinfection Directions. Treated surfaces must remain wet for 1 minute. Use a 30-minute contact time against Foot and Mouth Disease. Allow surfaces to air dry.

MOLD AND MILDEW CONTROL DIRECTIONS: Add 3 - 4 oz. per gallon of water to inhibit the growth of mold and mildew and their odors on hard, nonporous surfaces. Thoroughly wet all treated surfaces completely. Let air dry. Repeat application weekly or when growth or odor reappears.

***KILLS HIV-1 ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 ON SURFACES/ OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Clean up must always be done wearing protective gloves, gowns, masks and eye protection. **Cleaning Procedure:** Blood and other body fluids containing HIV must be thoroughly cleaned from surfaces and objects before application of Betco Sure Bet II.

(M)SDS No. 314



RLB2066

©2014 Betco Corporation
1001 Brown Avenue | Toledo, Ohio 43607
Made in U.S.A. | All Rights Reserved.

888-GO BETCO (888-462-3826) | Betco.com



1131414

LICENSED

PERIOD 2015-2017 LIC. NO. 8324.30

Contact Time: Leave surface wet for 1 minute. Use a 30-minute contact time against Foot and Mouth Disease. Use a 10-minute contact time for disinfection against all other bacteria claimed.

Disposal of Infectious Materials: Blood, body fluids, cleaning materials and clothing must be autoclaved and disposed of according to local regulations for infectious waste disposal.

HEAVY DUTY CLEANER DIRECTIONS: Use 8 oz. per gallon of water for heavy duty cleaning.

TO CLEAN AND DEODORIZE: Use 1:32 dilution. Apply to wet the surface thoroughly. Let stand for at least one minute, rinse with water or allow to air dry.

**PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals**

DANGER

Corrosive. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin, or on clothing. Wear protective eyewear (goggles, safety glasses or face shield), protective clothing and protective gloves (rubber or chemical resistant) when handling. Do not breathe spray. May be fatal if swallowed. Avoid contamination of food. Wash thoroughly with soap and water after handling and before eating, drinking, using tobacco, chewing gum or using the toilet. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Open dumping is prohibited. Store in original container in areas inaccessible to children.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Fill container ¼ full with water and reclose the container. Agitate vigorously, and dispose of rinsate consistent with pesticide disposal instructions. Repeat two more times. Then offer for recycling if available or puncture and dispose in sanitary landfill or by other procedures approved by state and local authorities. Follow pesticide disposal instructions for rinsate. If not triple rinsed, these containers are acute hazardous wastes and must be disposed in accordance with local, state, and federal regulations.



Sure Bet™ II

One-Step Acid Cleaner, Disinfectant, *Virucide, Mildewstat, Deodorizer with Organic Soil Tolerance

***Virucidal against HIV-1 (AIDS Virus) and Herpes Simplex Virus Types 1 & 2**

Active Ingredients:

Octyl decyl dimethyl ammonium chloride	0.75%
Diocetyl dimethyl ammonium chloride	0.30%
Didecyl dimethyl ammonium chloride	0.45%
Alkyl (C ₁₄ 50%, C ₁₂ 40%, C ₁₆ 10%) dimethyl benzyl ammonium chloride	1.00%
Inert Ingredients	97.50%
Total:	100.00%

**BATHROOM
CLEANER
314**

NET CONTENTS: 1 U.S. GALLON (3.78 L)

**KEEP OUT OF REACH
OF CHILDREN
DANGER
PELIGRO**

See side panel for Precautionary Statements and First Aid.

EPA EST. NO. 4170-OH-1
EPA REG. NO. 6836-86-4170