Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2014-2016 LIC. NO.

8334.2

"KEEP IT CLEAN!"

MONK™

Use MONK Disinfectant Wipes in Schools, Restaurants, Hospitals, Dental Offices, USDA inspected food processing facilities, Institutional Facilities, Athletic and Exercise Facilities, Health Clubs, Prisons, Public Restrooms, Showers, Bathrooms, and Kitchens.

DISINFECTANT WIPES

One Step Cleaner / Disinfectant / Deodorizer

KEEP OUT OF REACH OF CHILDREN

CAUTION

- Cleans and disinfects in one easy step
- Does not contain bleach or ammonia
- Effective in the presence of 5% serum contamination
- Kills bacteria****
- Strong wipe for heavy duty cleaning
- These wipes will not harm most surfaces

MONK DISINFECTANT WIPES KILL:

- Staphylococcus aureus (Staph)
- Staphylococcus aureus Methicillin-Resistant [MRSA] [CA-MRSA]
- HIV-1 [AIDS Virus]
- Avian Influenza [H3N2] [H5N1]

Active Ingredients:	
Octyl decyl dimethyl ammonium chloride	0.069%
Dioctyl dimethyl ammonium chloride	0.028%
Didecyl dimethyl ammonium chloride	0.042%
Alkyl (C_{14}, 50%; C_{12}, 40%; C_{16}, 10%) dimethyl benzyl ammonium chloride	0.093%
Other Ingredients:	99.768%
Total	*100.000%

* Includes weight of absorbent wipe.

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2011-2013 LIC. NO.

Antibacterial Formula Starts Killing Bacteria on Contact

LARGE 7" X 7" 700 TOWELETTES

Department of Agriculture
STATE OF HAWAII

LICENSED

PERIOD 2008-2010 LIC. NO.

See back panel for Precautionary Statements and First Aid EPA Reg. No. 6836-313-70930 EPA Est. No. 70930-PA-001 Item #69700

10.75” width

ight

Use Monk Disinfectant Wipes to clean and disinfect non-porous personal protection equipment, protective headgear, hard hats, half mask respirators, full face breathing apparatus, gas masks, goggles, spectacles, face shields, silicone rubber or PVC hearing protectors and vinyl covered ear muffs. Rinse all equipment that comes in contact with skin before reuse with clean warm water , and allow to air dry.

TO OPEN PACKAGE: Remove cover. Pull up wipe corner from center of roll. Thread through hole in the cover. Replace cover. Open center cap on cover to remove wipe. When not in use, close center cap to prevent moisture loss.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

TO CLEAN AND DEODORIZE: Wipe surface with wipe. Let air dry. Discard used wipe in trash. Do not flush in toilet.

TO CLEAN AND DISINFECT (DISINFECTION / VIRUCIDAL* / FUNGICIDAL / MOLD AND MILDEW CONTROL DIRECTIONS):
Wipe surface with wipe until surface is visibly wet. Allow surface to remain wet for 10 minutes. Let air dry. If surfaces are extremely dirty, clean first with another wipe before disinfecting. Discard used wipe in trash. Do not flush. Rinse food contact surfaces with clean, potable water after disinfecting. Do not use to disinfect dishes, glassware or utensils. Do not use as a diaper wipe or for personal cleansing. (Note: Do not use on unpainted wood, natural marble or brass.)

For fungicidal activity, MONK Disinfectant Wipes are an effective fungicide against Trichophyton mentagrophytes [the athlete's foot fungus] in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities. MONK Disinfectant Wipes is a one-step fungicide.

For mold and mildew, MONK Disinfectant Wipes will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, nonporous surfaces. Follow disinfection instructions. Repeat treatment every seven days, or more often if new growth appears.

*KILLS HIV AND HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces / objects with blood or body fluids, and in which the surfaces / objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) or Hepatitis B Virus (HBV) or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OR HBV OR HCV ON SURFACES / OBJECTS SOILED WITH BLOOD / BODY FLUIDS.
Personal protection: Clean-up must always be done wearing protective gloves, gowns, masks and eye protection.

Cleaning procedure: Blood and other body fluids containing HIV or HBV or HCV must be thoroughly cleaned from surfaces and objects before application of MONK Disinfectant Wipes. MONK Disinfectant Wipes can be used for this purpose.

*This product is for use on hard nonporous surfaces not conducive to treatment by immersion or excess liquid against Avian Influenza A.

Contact time: Leave surface wet for 1 minute (60 seconds) for HIV-1 and 10 minutes for HBV and HCV. Use a 10-minute contact time for disinfection against all other viruses, bacteria and fungi claimed. Disposal of infectious material: Blood, body fluids, cleaning materials and clothing should be autoclaved and disposed of according to local regulations for infectious waste disposal.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
Caution. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID
If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor or going for treatment.

STORAGE AND DISPOSAL:

Store product in original container in areas inaccessible to children. Keep securely closed. Refill empty container only with MONK Disinfectant Wipes. Discard empty container in trash or recycle.

MONK Disinfectant Wipes are effective against:
******[Bacteria]**
Community Associated Methicillin-Resistant Staphylcoccus aureus [CA-MRSA] NRS384, USA300
Community Associated Methicillin-Resistant Staphylcoccus aureus [CA-MRSA] NRS123, USA400
Staphylococcus aureus (Staph)
Salmonella enterica (Salmonella)
Pseudomonas aeruginosa [Pseudomonas]
Campylobacter jejuni [Campylobacter]
Acinetobacter baumannii
Brevibacterium ammoniagenes [Brevibacterium]
Campylobacter jejuni
Enterobacter aerogenes
Enterococcus faecalis [Enterococcus]
Enterococcus faecalis- Vancomycin Resistant [VRE]
Escherichia coli [E. coli]
ESBL Escherichia coli - [Extended spectrum beta-lactamase producing E. coli]
Klebsiella pneumoniae [Klebsiella]
Legionella pneumophilia
Pseudomonas cepacia [Pseudomonas]
Salmonella schottmuelleri [Salmonella]
Salmonella typhi [Salmonella]
Serratia marcescens [Serratia]
Shigella dysenteriae [Shigella]
Staphylococcus aureus - Methicillin-Resistant [MRSA]
Staphylococcus aureus - Multi-Drug Resistant
Staphylococcus aureus-Vancomycin Intermediate Resistant - [VISA]
Streptococcus pyogenes [Strep] [a cause of scarlet fever]
Vibrio cholerae
[Viruses:]
*Rotavirus (Diarrhea-causing virus)
*Hepatitis B Virus [HBV]
*Hepatitis C Virus [HCV]
*Herpes Simplex Virus Type 1
*Herpes Simplex Virus Type 2
*HIV-1 [AIDS Virus]
*Human Coronavirus
*Influenza Type A / Brazil [Influenza]
*Norwalk Virus [Norovirus]
*Respiratory Syncytial Virus [RSV]
*SARS Associated Coronavirus [SARS] [cause of Severe Acute Respiratory Syndrome]
*Vaccinia [Pox Virus]
[Animal Viruses:]
Avian Influenza [H5N1]
Avian Influenza [H3N2]
Avian Infectious Bronchitis Virus
Canine Distemper Virus
Feline Calicivirus
Newcastle's Disease Virus
Pseudorabies Virus
[Fungi:]
Aspergillus niger
Trichophyton mentagrophytes [the Athlete's Foot Fungus] [A cause of Ringworm of the foot]

National Towelette Company
1726 Woodhaven Drive
Bensalem, PA 19020

Phone: 215-245-7300