



STATE OF HAWAII
Department of Agriculture

ACCEPTED

LICENSE NO. **8334.5**

FLEX[®] Wipes DISINFECTANT WIPES

Use **FLEX WIPES Disinfectant Wipes** in Schools, Restaurants, Hospitals, Dental Offices, USDA inspected food processing facilities, Institutional Facilities, Exercise Facilities, Health Clubs, Prisons, Public Restrooms, Showers, Bathrooms, and Kitchens.

Kills 99.9% of bacteria in 15 seconds!

- *Effective in the presence of 5% serum contamination*
- *Kills Cold and Flu Virus***
- *Kills Mold & Mildew*
- *These wipes will not harm most surfaces*
- *Contains No Phosphate*
- *Does not contain bleach or ammonia*

A **ONE-STEP** disinfectant cleaner that is effective against a broad spectrum of bacteria when used as directed

ELIMINATES Pseudomonas, MRSA, CA-MRSA, E.coli, Campylobacter, VRE, Salmonella in 4 minutes!

800
LARGE 8" X 6"
TOWELETTES

Active Ingredients:	
Octyl decyl dimethyl ammonium chloride	0.0909%
Dioctyl dimethyl ammonium chloride	0.0364%
Didecyl dimethyl ammonium chloride	0.0545%
Alkyl (C ₁₄ , 50%; C ₁₂ , 40%; C ₁₆ , 10%) dimethyl benzyl ammonium chloride	0.1212%
Other Ingredients:	99.6970%
Total	100.0000% ¹

¹ Includes weight of absorbent wipe.

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

**On hard non-porous surfaces

EPA Reg. No. 6836-340-70930

EPA Est. No. 70930-PA-001

Item #10800

Net Wt: 11 lbs 12 ounces

See back panel for Precautionary Statements and First Aid

Use FLEX WIPES Disinfectant Wipes on washable hard, nonporous surfaces of: Chairs, Counters, Diaper Changing Tables, Doorknobs, Equipment tables, Faucets, Floors, Glass, Showers, Sinks, Tanning Beds, Telephones, Resuscitators, Stands and other hard nonporous surfaces made of: acrylic, metal, plastic and upholstery (vinyl and plastic).

TO OPEN PACKAGE: Open Lid. To start wipes, pull up on wipe and move wipe into the small dispensing opening and pull sheet up to remove wipe. The next wipe will pop up automatically. Close lid securely between uses.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use as a diaper wipe or for personal cleansing. This is not a baby wipe! Do not flush in toilet.

For plastic and painted surfaces, spot test on an inconspicuous area before use. FLEX WIPES Disinfectant Wipes will not harm most surfaces: including acrylic, sealed fiberglass and vinyl. Do not use on unpainted wood. [Note: Not recommended for unpainted wood, natural marble or brass.]

TO CLEAN AND DEODORIZE: Wipe surface with wipe. Let air dry. Discard used wipe in trash. Do not flush in toilet.

DISINFECTION / VIRUCIDAL* / MOLD AND MILDEW CONTROL DIRECTIONS:

Wipe surface with wipe until surface is visibly wet. Use enough wipes to thoroughly wet surfaces. Allow surface to remain wet for 4 minutes. For Candida albicans, treated surfaces must remain wet for 8 minutes. For Norwalk Virus [Feline Calicivirus], Norovirus, and Rotavirus, treated surfaces must remain wet for 10 minutes. Let air dry. If surfaces are extremely dirty, clean first with another wipe before disinfecting. Discard used wipe in trash. Do not flush.

Rinse food contact surfaces with clean, potable water after disinfecting. Do not use to disinfect dishes, glassware or utensils. **FOR MOLD AND MILDEW** FLEX WIPES Disinfectant Wipes will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, nonporous surfaces. Follow disinfection instructions. Repeat treatment every seven days, or more often if new growth appears.

SANITIZING:

To sanitize hard, non-porous, non-food contact surfaces; wipe surface, using enough wipes to allow treated surface to remain wet for 15 seconds for Staphylococcus aureus [Staph] (ATCC 6538), Klebsiella pneumoniae (ATCC 4352), and Campylobacter jejuni (ATCC 29428). Let surface dry. For highly soiled surfaces, clean excess dirt first.

***KILLS HIV AND HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces / objects with blood or body fluids, and in which the surfaces / objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) or Hepatitis B Virus (HBV) or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OR HBV OR HCV ON SURFACES / OBJECTS SOILED WITH BLOOD / BODY FLUIDS.

Personal protection: Clean-up must always be done wearing protective gloves, gowns, masks and eye protection.

Cleaning procedure: Blood and other body fluids containing HIV or HBV or HCV must be thoroughly cleaned from surfaces and objects before application of FLEX WIPES Disinfectant Wipes. FLEX WIPES Disinfectant Wipes can be used for this purpose.

Contact time: Leave surface wet for 1 minute for HIV-1. Use appropriate contact time as listed on label for all other viruses, bacteria and fungi claimed.

Disposal of infectious material: Blood, body fluids, cleaning materials and clothing must be disposed of according to local regulations for infectious waste disposal.

This product is for use on hard nonporous surfaces not conducive to treatment by immersion or excess liquid against Avian Influenza A.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.

Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor or going for treatment.

FLEX WIPES Disinfectant Wipes are effective against:

Bacteria

- Staphylococcus aureus - Community Associated Methicillin-Resistant [CA-MRSA] [NRS384] [USA300]
- Staphylococcus aureus - Community Associated Methicillin-Resistant [CA-MRSA] [NRS123] [USA400]
- Staphylococcus aureus (Staph)
- Salmonella enterica (Salmonella)
- Enterococcus faecalis- Vancomycin Resistant [VRE]
- Escherichia coli [E. coli]
- ESBL Escherichia coli - [Extended spectrum beta-lactamase producing E. coli]
- Klebsiella pneumoniae [Klebsiella]
- Legionella pneumophila
- Burkholderia cepacia
- Salmonella schottmuelleri [Salmonella]
- Salmonella typhi [Salmonella]
- Serratia marcescens [Serratia]
- Shigella dysenteriae [Shigella]
- Staphylococcus aureus-Vancomycin Intermediate Resistant - [VISA]
- Streptococcus pyogenes [Strep] [a cause of scarlet fever]
- Vibrio cholerae

Viruses:

- *Rotavirus
- *Hepatitis B Virus [HBV]
- *Hepatitis C Virus [HCV]
- *Herpes Simplex Virus Type 1
- *Herpes Simplex Virus Type 2
- *HIV-1 [AIDS Virus]
- *Human Coronavirus
- *Influenza Type A Virus / Brazil [Influenza]
- *Norwalk Virus [Norovirus]
- *Respiratory Syncytial Virus [RSV]
- *SARS Associated Coronavirus [SARS] [cause of Severe Acute Respiratory Syndrome]
- *Vaccinia Virus [Pox Virus]

Animal Viruses:

- Avian Influenza A [H5N1]
- Avian Influenza A [H3N2]
- Canine Distemper Virus
- Feline Calicivirus
- Newcastle's Disease Virus
- Pseudorabies Virus

Fungi:

- Candida albicans

STORAGE AND DISPOSAL:

Store in original container in areas inaccessible to children. Keep securely closed. Refillable container. Refill empty container only with FLEX WIPES Disinfectant Wipes. Do not reuse this container except as described. If not refilling, discard empty container in trash or recycle.

National Towelette Company
1726 Woodhaven Drive
Bensalem, PA 19020
Phone: 215-245-7300