



LICENSED

LICENSED

PERIOD 2011-2013 LIC. NO.

PERIOD 2014-2016 LIC. NO.

8378.22



**RESTRICTED USE PESTICIDE**  
DUE TO AVIAN AND AQUATIC TOXICITY  
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

# DIAZINON AG600 WBC Insecticide

**Water-Based Concentrate**  
For control of certain insects on fruits, nuts, vegetables, and  
ornamentals grown outdoors in nurseries.

<b>ACTIVE INGREDIENT:</b>		<b>% BY WT.</b>
Diazinon: O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate .....		56.0%
<b>INERT INGREDIENTS:</b> .....		44.0%
	<b>TOTAL</b>	<b>100.0%</b>

Shake contents well before using.  
**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

EPA REG. NO. 66222-103-34704

EPA EST. NO. 34704-MS-002

NET CONTENTS 2½ GALS. (9.46 L)

060711 V1D 01G12

FORMULATED FOR  
LOVELAND PRODUCTS, INC., P.O. BOX 1286, GREELEY, COLORADO 80632-1286

<b>FIRST AID</b>	
This product contains diazinon, an organophosphate insecticide that inhibits cholinesterase.	
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious or convulsing person.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<p><b>FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-866-944-8565.</b></p> <p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p> <p><b>NOTE TO PHYSICIAN:</b> This product contains diazinon, an organophosphate that inhibits cholinesterase. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal, but should be administered only in conjunction with atropine.</p>	

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing vapor or spray mist. Do not contaminate food or feed. Food utensils, such as teaspoons or measuring cups, must not be used for food purposes after use in measuring pesticides. Keep out of reach of domestic animals. Do not use on humans, household pets, or livestock. Do not permit children or pets to contact treated areas until spray has dried. Do not contaminate ornamental fishponds.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of any water-proof material. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.

#### Mixers, loaders, applicators, and other handlers using engineering controls must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves if mixing or loading
- Chemical resistant apron if mixing or loading

See engineering controls for additional requirements.

#### Handlers performing tasks, such as cleaning equipment or spill clean-up for which engineering controls are not feasible must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical resistant gloves
- Chemical resistant footwear plus socks, chemical resistant apron if exposed to the concentrate
- Chemical resistant apron if exposed to the concentrate,
- A respirator with an organic-vapor removing cartridge with a prefilter approved for pesticides (MSHA / NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA / NIOSH approval number prefix TC-14G) or a NIOSH-approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P, or HE prefilter. If mixing this product with oil, do not use N type pre filter.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

#### ENGINEERING CONTROLS

Mixers and loaders must use a closed system that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides that provides dermal and inhalation protection [40 CFR 170.240 (d)(4)], and must

- wear the personal protective equipment required for mixers/loaders using engineering controls;
- wear protective eyewear if the system operates under pressure; and
- be provided and have immediately available for use in an emergency, such as a broken package, spill, or equipment breakdown, chemical resistant footwear and the respirator specified in the PPE section of this labeling for handlers not using engineering controls.

Applicators using motorized ground equipment must use an enclosed cab that meets the definition in the Worker Protection Standard for Agricultural Pesticides [40 CFR 170.240 (d)(5)] for dermal protection. Flaggers supporting aerial applications to lettuce must use an enclosed cab that meets the definition in the Worker Protection Standard for Agricultural Pesticides [40 CFR 170.240 (d)(5)] for dermal protection.

In addition, applicators must:

- wear the personal protective equipment required in the PPE section of this labeling for handlers using engineering controls;
- either wear the respirator specified for handlers not using engineering controls or use an enclosed cab that is declared in writing by the manufacturer or by a government agency to provide at least as much respiratory protection as the respirator specified for handlers not using engineering controls;
- be provided and have immediately available for use in an emergency when they

must exit the cab in the treated area, coveralls, chemical resistant footwear, and if not already using one, the respirator specified for handlers not using engineering controls;

- take off any PPE that was worn in the treated area before reentering the cab, and
- store all such PPE in a chemical-resistant container, such as a plastic bag, to prevent contamination of the inside of the cab.

**Exception:** For application to apples and lettuce, see directions for use for a special exception to these engineering controls requirements.

Pilots must use an enclosed cockpit in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)]. When entering or leaving an aircraft contaminated with pesticide residues, pilots must wear chemical resistant gloves and must store used gloves in a chemical resistant container, such as a plastic bag, to prevent contamination of the inside of the cockpit.

**Note:** Aerial applications are permitted only on lettuce.

#### USER SAFETY RECOMMENDATIONS

##### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Do not exceed maximum permitted label rates. Rates above those recommended significantly increase potential hazards to birds, especially waterfowl. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwater. This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Reformulation or repackaging of this product is prohibited.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI). **The REI for each crop is listed in the directions for use associated with each crop.**

**Exception:** If the product is soil-injected or soil-incorporated, the Worker Protection Standard under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls worn over long sleeve shirt and long pants,
- Chemical-resistant gloves made of any waterproof material,
- Chemical-resistant footwear plus socks and
- Chemical resistant headgear if overhead exposures.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

#### WORK SAFETY RULES

**Repeated exposures to cholinesterase inhibitors such as are contained in this product may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor.** When handling Diazinon AG600 WBC, do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. See FIRST AID section of this label. When handling this product, wear the personal protective equipment listed on this label.

#### GENERAL INFORMATION

Diazinon AG600 WBC is an emulsifiable solution which, when diluted with water according to the directions, is intended for the control of insect pests of agricultural crops and other plants. Do not use in greenhouses. Do not use in barns or dwellings. Do not use on livestock. Aerial application is prohibited except as allowed on the label for use on lettuce.

In the directions for use which follow, best control is obtained when application is made when infestation is noticed.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, since crop or plant injury may result. Failure to follow the directions for use and precautions on this label may result in poor insect control, crop injury, or illegal residues.

Do not use water from treated cranberry beds for drinking purposes or to irrigate crops other than those appearing on EPA approved diazinon labels.

**Note:** The California Department of Pesticide Regulation has set a 5-day reentry period for diazinon on peaches and nectarines. Five days should elapse between the time of application and the time a worker may enter the field to engage in any activity requiring substantial contact with treated foliage. When a mixture of two or more organophosphate pesticides are applied in combination, the interval should be prolonged by adding to the largest applicable interval an additional 50% of that interval.

When an adjuvant is to be used with this product, Loveland Products, Inc. suggests the use of a Chemical Producers and Distributors Association certified adjuvant.

### SPRAY EQUIPMENT

Applications of Diazinon AG600 WBC using sufficient water volume to provide thorough and uniform coverage generally provide the most effective pest control.

To avoid spray drift, do not apply when conditions favor drift from beyond the target area. Avoid overlap because crop injury may occur.

Equip sprayers with nozzles that provide accurate and uniform application. Be certain that nozzles are the same size and uniformly spaced across the boom. Calibrate sprayer before use.

For more information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult your local or state agricultural authorities.

### MIXING PROCEDURES

Thoroughly clean spray equipment before using this product. Prepare only the amount of spray mixture needed for the immediate application. Vigorous agitation is necessary for proper dispersal of this product. Maintain maximum agitation throughout the spraying operation. Do not let the spray mixture stand in the spray tank overnight. Flush the spray equipment thoroughly following each use.

#### 1. Diazinon AG600 WBC Alone:

Add ½ of the required amount of water to the mix tank. With the agitator running, add the Diazinon AG600 WBC to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after the Diazinon AG600 WBC has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

#### 2. Diazinon AG600 WBC in Tank-Mixtures:

**Note:** Do not mix Diazinon AG600 WBC with any formulation of Captan or Captec® because crop injury may occur. Diazinon is physically compatible with most insecticide and fungicide products; if you are not certain of the physical compatibility of Diazinon AG600 WBC with other tank mix partners, please contact your local Loveland Products, Inc. sales representative. When using Diazinon AG600 WBC in tank mixtures, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank mix products' labels. Confirm that the tank mixture is safe to the crop by spraying a small area first, and evaluating crop safety after an appropriate period. Do not treat larger areas until crop safety has been confirmed.

Add ½ of the required amount of water to the mix tank. Start the agitator running before adding any tank-mix partners. Add tank-mix partners in this order: all products in water-soluble packaging, wettable powders, wettable granules (dry flowables), liquid flowables, liquids, and emulsifiable concentrates. Always allow each tank-mix partner to fully disperse before adding the next product. Maintain sufficient agitation while adding the remainder of the water and until all of the mixture has been applied.

### PREHARVEST INTERVAL

The required days to wait between the last application and harvest are given in ( ) after each crop name.

### FRUIT AND NUT CROPS

Diazinon AG600 WBC may be applied using ground application equipment as specified in the following table. This table indicates the minimum amount of water that can be used for application. The rate of Diazinon AG600 WBC for concentrated spray applied to fruit trees by ground equipment is based on a dilute full cover spray applied with conventional ground equipment at a rate of 300-400 gals. of water per acre. The specific amount of product to be used appears under the separate directions for use for each crop. If the rate calls for 1 pt. of Diazinon AG600 WBC in 100 gals. of water, this amounts to 3-4 pts. of product per acre when applied at the usual dilute ground spray volumes of 300 - 400 gals. of water per acre. In no case should the amount of product used per acre exceed the maximum amount per acre per application that is specified in the **COMMENTS** section for individual crops.

**Note:** To protect bees, do not apply this product to trees when trees or substantial numbers of weeds in the orchard are in bloom.

Diazinon AG600 WBC may be used during dormancy on deciduous trees and vines. When mixing with spray oils, follow oil manufacturer's use directions.

CROP	GROUND APPLICATION MINIMUM GALS./ACRE	
	Dilute	Concentrate
Almonds*	100	20
Apples	100	20
Apricots	100	20
Blackberries**	100	20
Blueberries	100	20
Boysenberries**	100	20
Cherries	100	20
Cranberries	15	-
Dewberries**	100	20
Loganberries**	100	20
Nectarines	100	20
Peaches	100	20
Pears	100	20
Plums	100	20
Prunes	100	20
Raspberries**	100	20
Strawberries	100	20

\*CA Only

\*\*CA, OH, OR, or WA Only

FRUIT AND NUT CROPS			
CROP	INSECT	RATE	COMMENTS
ALMONDS (California Only)	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite eggs, Brown Mite eggs, Twig Borer, Apple Aphid eggs, Black Cherry Aphid eggs, Mealy Plum Aphid eggs, Mealybugs	12 ¾ - 19 fl. oz. + 2 - 3 gals. dormant oil or 1-1 ½ gals. superior type oil in 100 gals. of water	Apply during the dormant season only.
	<ul style="list-style-type: none"> <li>• Do not apply more than 76 ½ fl. oz. of diazinon or 9 gals. of oil per acre.</li> <li>• Make a maximum of one application per year during the dormant season only.</li> <li>• Apply every other year unless pest infestations can be controlled only with consecutive annual treatments.</li> <li>• The REI is 7 days.</li> </ul>		
APPLES (21)	Woolly Apple Aphid	12 ¾ fl. oz. in 100 gals. of water	Apply when infestations occur. Slight russeting may occur on some varieties of apples such as Golden Delicious.
	San Jose Scale	12 ¾ fl. oz. in 100 gals. of water	Apply as a dormant or delayed-dormant spray.
<ul style="list-style-type: none"> <li>• A maximum of two applications are allowed per year: 1) either one as a dormant application and one foliar application, or 2) two in-season foliar applications regardless of target pest.</li> <li>• Minimum retreatment interval is 14 days.</li> <li>• EXCEPTION for Airblast Applications to Apples: When application using enclosed-cab airblast equipment is not feasible in apple orchards, applicators are permitted to use open-cab airblast equipment, PROVIDED they are wearing chemical-resistant headgear AND the personal protective equipment specified in this labeling for handlers not using engineering controls.</li> <li>• The REI is 4 days.</li> </ul>			
APRICOTS (21)	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite eggs, Brown Mite eggs, Twig Borer, Apple Aphid eggs, Black Cherry Aphid eggs, Mealy Plum Aphid eggs, Mealybugs	12 ¾ fl. oz. in 100 gals. of water	Apply as a dormant spray only once per year. When using spray oils, follow spray oil manufacturer's use directions.
	Aphids, Brown Mite (Clover Mite), Two-spotted Spider Mite, Olive Scale Crawler, San Jose Scale Crawler	12 ¾ fl. oz. in 100 gals. of water	Apply when infestation is noticed.
	Olive Scale Crawler	6 ½ fl. oz. + 1 ½ gals. of light medium horticultural oil per 100 gals. of water	Apply when crawlers are present.
	Apricot Mealybug	12 ¾ fl. oz. per 100 gals. of water	Apply as a cover spray from petal fall to June.
<ul style="list-style-type: none"> <li>• Do not apply more than 51 fl. ozs. of product per acre per application.</li> <li>• A maximum of two applications are allowed per year: 1) a maximum of one as a dormant application and 2) a maximum of one as an in-season foliar application regardless of target pest.</li> <li>• For in-season applications, apply every other year unless pest infestations can be controlled only with consecutive annual treatments.</li> <li>• The REI is 4 days.</li> </ul>			
BLUEBERRIES (7)	Cranberry Fruitworm, Cherry Fruitworm, Blueberry Maggot, Aphids, Thrips, Two-spotted Spider Mite	12 ¾ fl. oz./A	Apply when infestation occurs.
	Fire Ants	12 ¾ fl. oz./A	To aid in the control of fire ants, slowly apply 1 gal. of diluted mixture over and 6 inches around each fire ant mound. Apply gently to avoid disturbing ants. Use equipment capable of delivering the diluted product as a gentle rain. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness. For best results, apply in cool weather (65-80°F) or in early morning or late evening hours.
<ul style="list-style-type: none"> <li>• Do not apply more than 12 ¾ fl. oz. of product per acre per application.</li> <li>• A maximum of two applications are allowed per year: 1) a maximum of one as an in-season foliar application regardless of target pest and 2) a maximum of one for control of fire ants.</li> <li>• The REI is 5 days.</li> </ul>			

6

**FRUIT AND NUT CROPS cont'd.:**

CROP	INSECT	RATE	COMMENTS
CANEberries (Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries) (California, Ohio, Oregon, and Washington Only) (7)	Raspberry Fruitworm	25 ½ fl. oz./A in 100 gals. of water	Apply when insects occur. Apply in a minimum of 100 gals. of water per acre (200 gals. per acre maximum). Make application when blossom buds separate.
	Raspberry Crown Borer (Raspberry Root Borer)	51 fl. oz./A	Apply a single application in a minimum of 100 gals. of water per acre as a drench to the crown and lower canes in the spring before buds break.
	<ul style="list-style-type: none"> <li>Do not apply more than 51 fl. oz. of product per acre.</li> <li>Make a maximum of one application per year regardless of the target pest.</li> <li>The REI is 5 days.</li> </ul>		
CHERRIES (21)	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite eggs, Brown Mite eggs, Twig Borer, Apple Aphid eggs, Black Cherry Aphid eggs, Mealy Plum Aphid eggs, Mealybugs	12 ¾ fl. oz./A in 100 gals. of water	Apply as a dormant spray. When using spray oils, follow spray oil manufacturer's use directions. Apply only once during the dormant season.
	Eyespotted Bud Moth, Fruittree Leafroller, Leafhoppers, Black Cherry Aphid, San Jose Scale Crawlers	12 ¾ fl. oz. in 100 gals. of water	Apply when infestation occurs.
	Cherry Fruit Fly	6 ½ - 12 ¾ fl. oz. in 100 gals. of water	Apply when adult flies begin to emerge.
	Cherry Rust Mite	6 ½ - 12 ¾ fl. oz. in 100 gals. of water	Apply to trees after harvest.
	<ul style="list-style-type: none"> <li>Do not apply more than 51 fl. oz. of product per acre per application.</li> <li>A maximum of two applications are allowed per year: 1) a maximum of one as a dormant application and 2) a maximum of one as an in-season foliar application regardless of target pest.</li> <li>For in-season applications, apply every other year unless pest infestations can be controlled only with consecutive annual treatments.</li> <li>The REI is 4 days.</li> </ul>		
CRANBERRIES (7)	Blackheaded Fireworm ( <i>Rhopobota neavana</i> )	51 fl. oz./A	Apply to larval stage. Pheromone trap captures and sweep monitoring may be used to optimize treatment timing. Repeat application after 14 days if necessary. A maximum of 3 applications per year is permitted regardless of target pest.
	Cranberry Tipworm	51 fl. oz./A	Presence of damage is not an indicator of active infestation. For best results, target early-stage larvae. Repeat application after 14 days if necessary. A maximum of 3 applications per season is permitted.
	Cranberry Fruitworm	51 - 76 ½ fl. oz./A	Cranberry growth stage and/or berry inspection for eggs may be used to optimize treatment timing; consult your local extension agent or pest management adviser. Repeat application after 14 days if necessary. A maximum of 3 applications per year is permitted regardless of target pest.
	<ul style="list-style-type: none"> <li>For all pests, use in a minimum of 15 gals. of water per acre for ground application and up to 400 gals. of water per acre for chemigation.</li> <li>Do not make more than 3 applications per year regardless of the target pest.</li> <li>Do not use water from irrigated or flooded cranberry beds for drinking purposes or to irrigate crops other than those appearing on EPA-approved diazinon labels.</li> <li>The REI is 5 days.</li> </ul>		

**FRUIT AND NUT CROPS cont'd.:**

CROP	INSECT	RATE	COMMENTS
NECTARINES (21)	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite eggs, Brown Mite eggs, Twig Borer, Apple Aphid eggs, Black Cherry Aphid eggs, Mealy Plum Aphid eggs, Mealybugs	12 ¾ fl. oz. in 100 gals. of water	Apply as a dormant spray. When using spray oils, follow spray oil manufacturer's use directions. Apply only once during the dormant season.
	Aphids, Brown Mites (Clover Mites), Two-Spotted Spider Mites, Olive Scale Crawler, San Jose Scale Crawler, Peach Twig Borer, Oriental Fruit Moth	12 ¾ fl. oz. in 100 gals. of water	Apply when infestation occurs.
<ul style="list-style-type: none"> <li>• Do not apply more than 51 fl. oz. of product per acre per application.</li> <li>• A maximum of two applications are allowed per year: 1) a maximum of one as a dormant application and 2) a maximum of one as an in-season foliar application regardless of target pest.</li> <li>• For in-season applications, apply every other year unless pest infestations can be controlled only with consecutive annual treatments.</li> <li>• The REI is 4 days.</li> </ul>			
PEACHES (21)	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite eggs, Brown Mite eggs, Twig Borer, Apple Aphid eggs, Black Cherry Aphid eggs, Mealy Plum Aphid eggs, Mealybugs	12 ¾ fl. oz. in 100 gals. of water	Apply as a dormant spray. When using spray oils, follow spray oil manufacturer's use directions. Do not apply more than 51 oz. of product or 6 gals. of oil per acre. Apply only once during the dormant season.
	Aphids, Brown Mites (Clover Mites), Two-Spotted Spider Mite, Olive Scale Crawler, San Jose Scale Crawler	12 ¾ fl. oz. in 100 gals. of water	Apply when infestation occurs.
	Oriental Fruit Moth	12 ¾ fl. oz. in 100 gals. of water	Apply at 100% petal fall.
	Peach Twig Borer, Parlatoria Scale	12 ¾ fl. oz. in 100 gals. of water	Apply when infestation occurs.
	Leafhoppers	12 ¾ fl. oz. in 100 gals. of water	Apply when infestation occurs.
	White Peach Scale	19-25 ½ fl. oz. in 100 gals. of water	Apply post harvest timed to coincide with peak crawler and immature scale activity.
<ul style="list-style-type: none"> <li>• Do not apply more than 51 fl. oz. of product per acre per application.</li> <li>• A maximum of two applications are allowed per year: 1) a maximum of one as a dormant application and 2) a maximum of one as an in-season foliar application regardless of target pest.</li> <li>• For in-season applications, apply every other year unless pest infestations can be controlled only with consecutive annual treatments.</li> <li>• The REI is 4 days.</li> </ul>			
PEARS (21)	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite eggs, Brown Mite eggs, Pear Leaf Blister Mite eggs, Twig Borer, Apple Aphid eggs, Black Cherry Aphid eggs, Mealy Plum Aphid eggs, Mealybugs	12 ¾ fl. oz. + 2-3 gals. dormant oil or 1-1 ½ gals. superior type oil in 100 gals. of water	Apply as a dormant spray. Do not apply more than 51 oz. of product or 6 gals. of oil per acre. Apply only once during the dormant season.
	Pear Psylla	6 ½ -12 ¾ fl. oz. in 100 gals. of water	Apply as a cover spray at pre-pink stage.
	Tentiform Leafminer, Mealybugs, Codling Moths, Rosy Apple Aphids, Green Apple Aphids, Woolly Apple Aphids, Spider Mites	12 ¾ fl. oz. in 100 gals. of water	Apply as a cover spray. Note: Mites will be suppressed if diazinon is used in this regular cover spray program.
	Apple Maggot	12 ¾ fl. oz. in 100 gals. of water	Apply as a cover spray when flies are active and laying eggs.
	San Jose Scale Crawlers, Forbes Scale Crawlers	12 ¾ fl. oz. in 100 gals. of water	Apply when crawlers first appear.
<ul style="list-style-type: none"> <li>• Do not apply more than 51 fl. oz. of product per acre per application.</li> <li>• A maximum of two applications are allowed per year: 1) a maximum of one as a dormant application and 2) a maximum of one as an in-season foliar application regardless of target pest.</li> <li>• For in-season applications, apply every other year unless pest infestations can be controlled only with consecutive annual treatments.</li> <li>• The REI is 4 days.</li> </ul>			

8

**FRUIT AND NUT CROPS cont'd.:**

<b>CROP</b>	<b>INSECT</b>	<b>RATE</b>	<b>COMMENTS</b>
<b>PLUMS (21)</b>	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite eggs, Brown Mite eggs, Twig Borer, Apple Aphid eggs, Black Cherry Aphid eggs, Mealy Plum Aphid eggs, Mealybugs	12 ¾ fl. oz. in 100 gals. of water	Apply as a dormant spray. When using spray oils, follow spray oil manufacturer's use directions. Apply only once during the dormant season.
	Brown Mite (Clover Mite), European Red Mite	12 ¾ fl. oz. in 100 gals. of water	Apply when infestation occurs.
	Leafcurl Plum Aphid, Mealy Plum Aphid, Thistle Aphid	6 ½ - 12 ¾ fl. oz. in 100 gals. of water	
	<ul style="list-style-type: none"> <li>• Do not apply more than 51 fl. oz. of product per acre per application.</li> <li>• A maximum of two applications are allowed per year: 1) a maximum of one as a dormant application and 2) a maximum of one as an in-season foliar application regardless of target pest.</li> <li>• For in-season applications, apply every other year unless pest infestations can be controlled only with consecutive annual treatments.</li> <li>• The REI is 4 days.</li> </ul>		
<b>PRUNES (21)</b>	San Jose Scale, Parlatoria Scale, Black Scale, Brown Scale, Apricot Scale, European Red Mite eggs, Brown Mite eggs, Twig Borer, Apple Aphid eggs, Black Cherry Aphid eggs, Mealy Plum Aphid eggs, Mealybugs	12 ¾ fl. oz. in 100 gals. of water	Apply as a dormant spray. When using spray oils, follow spray oil manufacturer's use directions. Apply only once during the dormant season.
	Brown Mite (Clover Mite), European Red Mite	12 ¾ fl. oz. in 100 gals. of water	Apply when infestation occurs.
	Leafcurl Plum Aphid, Mealy Plum Aphid, Thistle Aphid	6 ½-12 ¾ fl. oz. in 100 gals. of water	Apply when infestation occurs.
	<ul style="list-style-type: none"> <li>• Do not apply more than 51 fl. oz. of product per acre per application.</li> <li>• A maximum of two applications are allowed per year: 1) a maximum of one as a dormant application and 2) a maximum of one as an in-season foliar application regardless of target pest.</li> <li>• For in-season applications, apply every other year unless pest infestations can be controlled only with consecutive annual treatments.</li> <li>• The REI is 4 days.</li> </ul>		
<b>STRAWBERRIES (5)</b>	Aphids, Two-spotted Spider Mite, Strawberry Leafroller	12 ¾ fl. oz. in 100 gals. of water	Apply when pests occur.
	Cyclamen Mite	25 ½ fl. oz. in 100 gals. of water per acre	Direct spray to the plant crown. Plants should be sufficiently agitated to assure thorough coverage of the foliage.
	Mole Crickets	25 ½ fl. oz.	Broadcast in sufficient water to obtain even coverage when insects are present. Apply 1 - 2 days before transplanting and immediately incorporate into the top 1 - 2 inches of soil.
	<ul style="list-style-type: none"> <li>• Do not apply more than 25 ½ fl. oz. of product per acre per application.</li> <li>• Make a maximum of one foliar application per crop and a maximum of one soil application per crop.</li> <li>• The REI is 3 days.</li> </ul>		

**VEGETABLE CROPS**

For foliar or soil applications, Diazinon AG600 WBC should always be applied as a spray in sufficient water to assure thorough coverage of the foliage or soil. Diazinon AG600 WBC may be applied using ground or aerial (lettuce only) application equipment as specified in the following table. Aerial application is prohibited except as allowed on the label for use on lettuce. This table indicates the minimum amount of water that can be used for the application. The specific amount of product to be used appears under the separate directions for use for each crop.

Crop	Ground Application	Min. Gals.	Aerial Application
	Min. GPA	1000 Sq. Ft. (Ground)	Min. GPA
Beets, Red (Table)	10	2	-
Broccoli	10	2	-
Brussels Sprouts	10	2	-
Cabbage	10	2	-
Cantaloupes	5	1	-
Carrots	10	2	-
Casaba Melons	5	1	-
Cauliflower	10	2	-
Collards	10	2	-
Crenshaw Melons	5	1	-
Endive (Escarole)	10	2	-
Ginseng	10	2	-
Honeydew Melons	5	1	-
Kale	10	2	-
Lettuce, Head	10	2	5
Lettuce, Leaf	10	2	5
Muskmelons	5	1	-
Mustard Greens	10	2	-
Onions (Bulb and Green)	10	2	-
Persian Melons	5	1	-
Radishes	10	2	-
Rutabagas	10	2	-
Spinach	10	2	-
Tomatoes	10	2	-
Watermelons	5	1	-

Observe stated time intervals between last application and harvest as well as intervals between and maximum numbers of applications per season. Do not apply in greenhouses.

NOTES: (1) Soil Incorporation: Following application of this product to control vegetable soil insects, immediately incorporate the product into the soil to the recommended depth (see directions for individual crops) using a rotary hoe, cultivator, disk, harrow, or other suitable means. For control of surface cutworms, incorporate 2-3 inches; for subterranean cutworms, incorporate 3-6 inches. (2) Diazinon will not control organophosphate-resistant leafminers.

#### VEGETABLE CROPS

CROP	INSECT	RATE	COMMENTS
BEETS, RED (Table)	Cutworms	51 - 102 fl. oz/A	Broadcast just before planting and immediately incorporate into the (Table) soil. See Soil Incorporation note.
	Mole Crickets	25 ½ fl. oz/A	One to two days before planting, broadcast and immediately incorporate into the top 1 - 2 inches of soil.
	Wireworms	76 ½ - 102 fl. oz/A	Broadcast just before planting and immediately incorporate into the top 4 - 8 inches of soil.
<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>The REI is 3 days.</li> </ul>			

10

**VEGETABLE CROPS cont'd.:**

<b>CROP</b>	<b>INSECT</b>	<b>RATE</b>	<b>COMMENTS</b>
<b>BROCCOLI</b>	Root Maggots	51- 76 ½ fl. oz/A	Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil.
		3 ¼ - 6 ½ fl. oz. per 50 gals. of water	In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply recommended rate by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.
	Cutworms	51- 102 fl. oz/A	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	25 ½ fl. oz/A	One to two days before planting, broadcast and immediately incorporate into the top 1 - 2 inches of soil.
	Wireworms	76 ½ - 102 fl. oz/A	Broadcast just before planting and immediately incorporate into the top 4 - 8 inches of soil.
<ul style="list-style-type: none"> <li>• Do not make more than one soil application per year regardless of target pest.</li> <li>• The REI is 4 days.</li> </ul>			
<b>BRUSSELS SPROUTS</b>	Root Maggots	51 - 76 ½ fl. oz/A	Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil.
		3 ¼ - 6 ½ fl. oz. per 50 gals. of water	In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply recommended rate by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.
	Cutworms	51-102 fl. oz/A	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	25 ½ fl. oz/A	One to two days before planting, broadcast and immediately incorporate into the top 1 - 2 inches of soil.
	Wireworms	76 ½ - 102 fl. oz/A	Broadcast just before planting and immediately incorporate into the top 4 - 8 inches of soil.
<ul style="list-style-type: none"> <li>• Do not make more than one soil application per year regardless of target pest.</li> <li>• The REI is 4 days.</li> </ul>			
<b>CABBAGE</b>	Root Maggots	51 - 76 ½ fl. oz/A	Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil.
		3 ¼ - 6 ½ fl. oz. in 50 gals. of water	In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply recommended rate by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.
	Cutworms	51- 102 fl. oz/A	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	25 ½ fl. oz/A	One to two days before planting, broadcast and immediately incorporate into the top 1 - 2 inches of soil.
	Wireworms	76 ½ - 102 fl. oz/A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
<ul style="list-style-type: none"> <li>• Do not make more than one soil application per year regardless of target pest.</li> <li>• The REI is 4 days.</li> </ul>			

**VEGETABLE CROPS cont'd.:**

<b>CROP</b>	<b>INSECT</b>	<b>RATE</b>	<b>COMMENTS</b>
<b>CARROTS</b>	Cutworms	51 - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	25 ½ fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1 - 2 inches of soil.
	Wireworms	76 ½ - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>The REI is 3 days.</li> </ul>		
<b>CAULIFLOWER</b>	Root Maggots	51 - 76 ½ fl. oz./A	Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil.
		3 ¼ - 6 ½ fl. oz. in 50 gals. of water	In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply recommended rate by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.
	Cutworms	51 - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	25 ½ fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	76 ½ - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>The REI is 4 days.</li> </ul>		
<b>COLLARDS</b>	Cutworms	51 - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	25 ½ fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1 - 2 inches of soil.
	Wireworms	76 ½ - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>The REI is 4 days.</li> </ul>		
<b>ENDIVE (Escarole)</b>	Cutworms	51 - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	25 ½ fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	76 ½ - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>The REI is 4 days.</li> </ul>		
<b>GINSENG (30)</b>	Leafhoppers, Aphids, Lygus Bugs, Flea Beetles, Jumping Plant Lice	9 ½ - 12 ¾ fl. oz./A	Spray when insects first appear. For heavy insect infestations, use 12 ¾ fl. oz. per acre. Apply by ground equipment using 10-100 gals. of water per acre.
	<ul style="list-style-type: none"> <li>Do not apply more than one application per year.</li> <li>Do not apply during flowering of three- or four-year old crops.</li> <li>Do not graze treated areas or feed treated forage to livestock.</li> <li>The REI is 3 days.</li> </ul>		

12

**VEGETABLE CROPS cont'd.:**

<b>CROP</b>	<b>INSECT</b>	<b>RATE</b>	<b>COMMENTS</b>
<b>KALE</b>	Cutworms	51 - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	25 ½ fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	76 ½ - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>The REI is 4 days.</li> </ul>		
<b>LETTUCE</b> (Head and Leaf) (14)	Cutworms	51 fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	25 ½ fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	51 fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminer	6 ½ - 12 ¾ fl. oz./A	Apply when insects occur.
<ul style="list-style-type: none"> <li>Do not make more than one soil application per crop regardless of target pest.</li> <li>Do not make more than one foliar application per crop regardless of target pest.</li> <li>Aerial application is permitted on lettuce.</li> <li>EXCEPTION: Applicators applying to lettuce are permitted to use open-cab equipment provided they are wearing chemical-resistant head gear AND the personal protective equipment specified in this labeling for handlers not using engineering controls.</li> <li>The REI is 3 days.</li> </ul>			
<b>MELONS</b> (Cantaloupes, Casabas, Crenshaws, Honeydews, Muskmelons, Persians and hybrids of these) and <b>WATERMELONS</b> (3)	Cutworms	51 - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Wireworms	76 ½ - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>The REI is 3 days.</li> </ul>			
<b>HONEYDEW MELONS</b>	Aphids, Dipterous Leafminer, Thrips, Striped Cucumber Beetle, Melonworms, Leafhoppers, Spider mites	6 ½ - 19 fl. oz./A	Apply when infestation occurs.
	<ul style="list-style-type: none"> <li>Do not make more than one foliar application per year regardless of target pest.</li> <li>The REI is 3 days.</li> </ul>		
<b>MUSTARD GREENS</b>	Cutworms	51 - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	25 ½ fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	76 ½ - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>The REI is 4 days.</li> </ul>			
<b>ONIONS</b> (Bulb and Green)	Onion Maggots	51 - 102 fl. oz./A	Broadcast just before planting and mix into the top 3-4 inches of soil. Note: Diazinon will not control organophosphate-resistant onion maggots.
	Wireworms	76 ½ - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
<ul style="list-style-type: none"> <li>Do not make more than one soil application per year regardless of target pest.</li> <li>The REI is 3 days.</li> </ul>			

**VEGETABLE CROPS cont'd.:**

<b>CROP</b>	<b>INSECT</b>	<b>RATE</b>	<b>COMMENTS</b>
<b>RADISHES</b>	Cutworms	51 - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	25 ½ fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	76 ½ - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
<ul style="list-style-type: none"> <li>• Do not make more than one soil application per year regardless of target pest.</li> <li>• The REI is 3 days.</li> </ul>			
<b>RUTABAGAS</b>	Cabbage Maggot	51- 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the soil to a depth of 4 inches. See Soil Incorporation note.
	<ul style="list-style-type: none"> <li>• Do not make more than one soil application per year.</li> <li>• The REI is 4 days.</li> </ul>		
<b>SPINACH</b>	Cutworms	51- 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Wireworms	76 ½ - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	<ul style="list-style-type: none"> <li>• Do not make more than one soil application per year regardless of target pest.</li> <li>• The REI is 3 days.</li> </ul>		
<b>TOMATOES</b>	Cutworms	51-102 fl. oz./A	Broadcast just before planting and immediately incorporate into the soil. See Soil Incorporation note.
	Mole Crickets	25 ½ fl. oz./A	One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	76 ½ - 102 fl. oz./A	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	<ul style="list-style-type: none"> <li>• Do not make more than one soil application per growing season regardless of target pest.</li> <li>• The REI is 2 days.</li> </ul>		

**ORNAMENTAL STATEMENT**

Before treating a large number of ornamental plants with Diazinon AG600 WBC alone or as a tank mixture with any other material, make a test application on a few plants and observe for 7 to 10 days prior to treating large areas to reduce the possibility of plant injury.

INSECT CONTROL ON ORNAMENTALS GROWN OUTDOORS IN NURSERIES				
CROP	PEST	RATE	COMMENTS	
ARBORVITAE, AZALEA, BALSAM FIR (Maine Only), BIRCH, BOXWOOD, CAMELLIA, CARNATION, CHRYSANTHEMUM, CRABAPPLE, DOGWOOD, DOUGLAS FIR, ELM, EUONYMUS, GLADIOLI, HAWTHORN, HOLLY, HONEY LOCUST, HONEYSUCKLE, JUNIPER, LILAC, LILY, LOCUST, MAPLE, MARIGOLD, OAK, PANSY, PETUNIA, PINE, ORNAMENTAL PLUM, POPLAR, RHODODENDRON, ROSE, SCOTCH PINE, SPIREA, SPRUCE, SYCAMORE, WAX PLANT (Hoya), WILLOW, YELLOWWOOD, and YEW	Aphids, Bagworms, Carnation Bud Mite, Carnation Shoot Mite, Clover Mite, Cyclamen Mite, Dipterous Leafminer, European Pine Shoot Moth, European Red Mite, Flea Beetles, Holly Bud Moth, Leafhoppers, Obscure Root Weevil, Omnivorous Leaf-tier, Privet Mite, Scale Crawlers (Cottony-cushion, Lecanium, Pine Needle, San Jose, Soft), Thrips, Two-spotted Spider Mite, Juniper Webworm, Whiteflies	½ fl. oz. per 3 gals. water or 12 ¾ fl. oz. per 100 gals. of water	Apply the recommended rates. Spray when listed pests first appear. Try to spray underside of leaves and penetrate dense foliage.	
	Apple-and-Thorn Skeletonizer, Cotoneaster Webworm, European Pine Sawfly, Fall Webworm, Hemlock Chermes, Oak Looper, Obliquebanded Leafroller, Pear Slug, Tent Caterpillar	1 ¼ fl. oz. per 3 gals. of water or 25.6 fl. oz. per 100 gals. of water		
	Mimosa Webworm	2 ½ teaspoons in 3 gals. of water or 13 ¼ fl. oz. in 100 gals. of water	Apply as a thorough foliar spray when adults first appear.	
<ul style="list-style-type: none"> <li>Do not use on Ferns, Poinsettia, Hibiscus, Papaya, Pilea, and Gardenia, since injury to the plants may occur.</li> <li>The maximum application rate on all ornamentals is 25.6 ounces per acre.</li> <li>Make a maximum of one application per crop cycle.</li> <li>The REI is 7 days for flowers and other commercial ornamentals grown for cutting.</li> <li>The REI is 2 days for all other commercial ornamentals.</li> <li>Application is permitted only on commercial ornamentals grown outdoors in nurseries.</li> </ul>				

### CHEMIGATION (APPLICATION THROUGH AN IRRIGATION SYSTEM)

Diazinon AG600 WBC may be applied alone or in combination with other pesticides registered for application through sprinkler (including center pivot, lateral move, end tow side (wheel) roll, traveler, big gun, solid set, or hand move) or drip (trickle) irrigation systems (including surface and subsurface). Do not apply this product through any other type of irrigation system. Diazinon is physically compatible with most insecticide and fungicide products; if you are not certain of the physical compatibility of Diazinon AG600 WBC with other tank mix partners, please contact your local Loveland Products, Inc. sales representative. **Note: Do not mix Diazinon AG600 WBC with any formulation of Captan or Captec® because crop injury may occur.**

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect the irrigation system used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments if the need arises.

#### Chemigation systems connected to public water systems

**Note:** Loveland Products, Inc. does not encourage connecting chemigation sys-

tems to public water supplies. The following information is provided for users who have diligently considered all other application and water supply options before electing to make such a connection.

1. Public water systems means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from a point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

#### Sprinkler and Drip (trickle) Chemigation

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not use sprinkler chemigation when wind speed favors drift beyond the area intended for treatment.

#### Operating Instructions

1. Determine the acreage to be treated by the system.
2. Measure the appropriate amount of Diazinon AG600 WBC needed to treat the acreage to be treated. Refer to the specific directions for use for the application rate and the amount of water per acre.
3. Add the pre-measured Diazinon AG600 WBC to the chemigation system supply tank and dilute with water if needed. Proper agitation is needed if Diazinon AG600 WBC is tank mixed with other pesticides registered for sprinkler or drip chemigation.
4. Start the irrigation system and bring the system up to full pressure, making certain that all sprinkler heads are functioning properly.
5. Engage the chemigation injection or venturi system to add the Diazinon AG600 WBC to the irrigation lines.
6. Continue to run irrigation system for a few minutes after supply tank is empty to allow material in the irrigation lines to reach the sprinkler heads furthest from the chemigation injection point.

**Note:** When treatment with Diazinon AG600 WBC has been completed, further field irrigation over the treated area should be avoided for 48 hours to prevent washing the product off of the crop.

Apply Diazinon AG600 WBC only in sufficient irrigation water to achieve thorough plant coverage. Excess water will wash the product from the plant resulting in poor control.

To avoid a hazard to animals, wells or adjoining crops, do not allow irrigation water to collect or runoff.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**PESTICIDE STORAGE:** If freezing occurs, place the product in a warm room (68 °F or above) and allow to thaw. Roll or shake container frequently until thawing is complete.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. To avoid harming aquatic organisms in rivers and other surface waters, do not pour the concentrate, spray mixture, or rinse water into sanitary drains (for example, toilets, floor drains, and sinks) or into storm water sewers (for example street drains). If pesticide, spray mixture, or rinsate cannot be disposed of by use according to the label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative

#### Storage & Disposal cont'd.:

at the nearest EPA Regional Office for guidance.

#### CONTAINER DISPOSAL:

**Nonrefillable Container (five gallons or less):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then puncture and dispose of in a sanitary landfill, incinerate or burn, if allowed by state and local authorities. Stay out of smoke from burning container.

**Nonrefillable Container (greater than five gallons):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then puncture and dispose of in a sanitary landfill, incinerate or burn, if allowed by state and local authorities. Stay out of smoke from burning container.

**Refillable Container: (greater than 55 gallons):** Refillable container. Refill this container with diazinon only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC – 1-800-424-9300.**

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes.

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY BEFORE BUYING OR USING THIS PRODUCT,** read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not

specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Captec is a registered trademark of MICRO FLO COMPANY LLC

**SUPPLEMENTAL LABEL  
DIAZINON AG600 WBC INSECTICIDE  
EPA REG. NO. 66222-103-34704**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL AND PAMPHLET. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

**FOR PROTECTION OF AQUATIC SPECIES IN COUNTIES AND PORTIONS OF COUNTIES LOCATED IN THE SACRAMENTO AND SAN JOAQUIN VALLEYS**

**BELOW 1000 FEET ELEVATION FROM POTENTIAL CONTAMINATION IN SURFACE WATERS OF DIAZINON APPLIED DURING THE DORMANT SPRAY PERIOD, THE FOLLOWING ADDITIONAL LABEL REQUIREMENTS APPLY:**

- Dormant applications on orchard crops are restricted to ground application equipment only.
- Do not apply within 100 feet upslope of "sensitive aquatic sites" such as any irrigation ditch, drainage canal or body of water that may drain into a river or tributary unless a suitable method is used to contain or divert runoff waters. Waters that are contained or diverted must be held for a minimum of 72 hours before release into a sensitive aquatic site.
- Maintain a vegetative buffer strip a minimum of 10 feet wide from the edge of a field that is adjacent to and within 100 feet of sensitive aquatic sites.
- Do not apply this product to orchards when soil moisture is at field capacity and/or when a storm event likely to produce runoff from the treated orchard is forecasted by NOAA/NWS (National Weather Service) to occur within 48 hours following application.
- Make dormant applications only when insect scouting information or the recommendation of a Pest Control Advisor indicate treatment is required. (See UC IPM Guidelines for San Jose Scale in stone fruits and almonds and aphids in stone fruits. Use the prune dormant spur sampling program to determine need for a dormant treatment in that crop).
- Apply only when wind speed is 3-10 mph at the application site as measured by an anemometer outside of the orchard on the side nearest and upwind from a sensitive site.
- When sensitive aquatic sites are downwind from orchards, spray the first three rows nearest the sensitive aquatic sites only when the wind is blowing away from the sites. The row at the edge of the field next to sensitive aquatic sites must be sprayed with the outside nozzles turned off. Spray must not be directed higher than the tree canopy and spray must be directed away from sensitive aquatic sites.
- The Stewardship Bulletin "Orchard Practices for Protecting Surface Water" must be available to handlers and equipment operators at the application site during all application activities. (Note: Bulletin available through CURES.)

**THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION. PLEASE REFER TO CONTAINER LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.**